Similarities and differences of dietary and other determinants of iodine status in pregnant women from three European birth cohorts

Mariana Dineva, Margaret P. Rayman, Deborah Levie, Mònica Guxens, Robin P. Peeters, Jesus Vioque, Llúcia González, Mercedes Espada, Jesús Ibarluzea, Jordi Sunyer, Tim I.M. Korevaar, Sarah C. Bath

Affiliations

Department of Nutritional Sciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford, United Kingdom (MD, MPR, and SCB); The Generation R Study Group, Erasmus University Medical Centre, Rotterdam, The Netherlands (DL and TIMK); Department of Internal Medicine, Academic Centre For Thyroid Diseases, Erasmus University Medical Centre, Rotterdam, The Netherlands (DL, RPP, and TIMK); Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Centre–Sophia Children’s Hospital, Rotterdam, The Netherlands (DL and MG); ISGlobal, Barcelona, Spain (DL, MG, and JS); Pompeu Fabra University, Barcelona, Spain (DL, MG, and JS); Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Instituto de Salud Carlos III, Madrid, Spain (DL, MG, JV, JI, and JS); Nutritional Epidemiology Unit, Miguel Hernández University, ISABIAL-FISABIO, Alicante, Spain (JV); Epidemiology and Environmental Health Joint Research Unit, FISABIO-Universitat Jaume I-Universitat de València, Valencia, Spain (LG); Predepartamental Unit of Medicine, University Jaume I, Castelló, Spain (LG); Departamento de Salud del Gobierno Vasco, Public Health Laboratory of Bilbao, Basque Government, Derio, Spain (ME); BIODONOSTIA Health Research Institute, Donostia – San Sebastián, Spain (ME and JI); Departamento de Salud del Gobierno Vasco, Subdirección de Salud Pública de Guipúzcoa, Donostia – San Sebastián, Spain (JI);

Facultad de Psicología, University of the Basque Country UPV/EHU, Donostia - San Sebastián, Spain (JD); and Hospital del Mar Research Institute (IMIM), Barcelona, Spain (JS).

Corresponding author

Sarah C. Bath, Department of Nutritional Sciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford GU2 7XH, United Kingdom. E-mail: s.bath@surrey.ac.uk. Telephone: +44 (0)1483 683631.

Acknowledgements

EUthyroid project: This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 634453.

ALSPAC: We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The UK Medical Research Council and Wellcome (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and Dr Bath will serve as a guarantor for the contents of this paper. ALSPAC data collection is funded from a wide range of sources, a comprehensive list of which is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). The existing iodine measurements in ALSPAC were funded from (i) the NUTRIMENTHE project, which received a research grant from the European Community's 7th Framework Programme (FP7/2008–2013) under grant agreement 212652; and (ii) a Ph.D. studentship that was funded by Wassen International and the Waterloo Foundation (2009–2012).

We would like to thank Dr Pauline Emmett for helping with the dietary analysis of the ALSPAC food frequency questionnaire.

Generation R: The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, Rotterdam, and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond (STAR-MDC), Rotterdam. The Generation R Study is supported by the Erasmus Medical Center, Rotterdam, the Erasmus University Rotterdam, the Netherlands Organization for Health Research and Development (ZonMw), the Netherlands Organization for Scientific Research (NWO), the Ministry of Health, Welfare and Sport. A grant from the Sophia Children's Hospital Research Funds supports the neurodevelopmental work on thyroid; Robin P. Peeters is supported by a clinical fellowship from ZonMw, project number 90700412.

INMA: This study was funded by grants from UE (FP7-ENV-2011 cod 282957 and HEALTH.2010.2.4.5-1) and Spain: Instituto de Salud Carlos III (Red INMA G03/176; CB06/02/0041; FIS-FEDER: PI041436, PI05/1079, PI06/0867, PI081151, FIS-and PS09/00090, PI11/01007, PI11/02591, PI11/02038, PI13/1944, PI13/2032, PI14/00891, PI14/01687, and PI16/1288; Miguel Servet-FEDER CP11/00178, CP15/00025, and CPII16/00051, MS13/00054), Generalitat Valenciana: FISABIO (UGP 15-230, UGP-15-244, and UGP-15-249), Generalitat de Catalunya-CIRIT 1999SGR 00241, Fundació La marató de TV3 (090430), Department of Health of the Basque Government (2005111093 and 2009111069), and the Provincial Government of Gipuzkoa (DFG06/004 and DFG08/001).

Abbreviations

ALSPAC, the Avon Longitudinal Study of Parents and Children

BMI, body mass index

- 75 CI, confidence interval
- 76 CRM, certified reference material
- 77 EXP, exponentiation
- 78 FFQ, food frequency questionnaire
- 79 INMA, INfancia y Medio Ambiente
- 80 IQ, intelligence quotient
- 81 RSD, relative standard deviation
- 82 SD, standard deviation
- 83 SEE, standard error of estimate
- 84 SFFQ, semi-quantitative food frequency questionnaire
- 85 UI/Creat, urinary iodine-to-creatinine ratio
- 86 UIC, urinary iodine concentration
- 87 UK, United Kingdom
- 88 WHO, World Health Organisation

89 Abstract

90 **Purpose:** As a component of thyroid hormones, adequate iodine intake is essential during
91 pregnancy for fetal neurodevelopment. Across Europe, iodine deficiency is common in
92 pregnancy but data are lacking on the predictors of iodine status at this life stage. We
93 therefore aimed to explore determinants of iodine status during pregnancy in three European
94 populations of differing iodine status.

95 **Methods:** Data were from 6566 pregnant women from three prospective population-based
96 birth cohorts from the United Kingdom (ALSPAC, n=2852), Spain (INMA, n=1460), and the
97 Netherlands (Generation R, n=2254). Urinary iodine-to-creatinine ratio (UI/Creat, $\mu\text{g/g}$) was
98 measured in spot-urine samples in pregnancy (≤ 18 -weeks gestation). Maternal dietary intake,
99 categorised by food groups (g/day) was estimated from food-frequency questionnaires
100 (FFQs). Multivariable regression models used dietary variables (energy-adjusted) and
101 maternal characteristics as predictors of iodine status.

102 **Results:** Median UI/Creat in pregnant women of ALSPAC, INMA and Generation R was
103 121, 151, and 210 $\mu\text{g/g}$, respectively. Maternal age was positively associated with UI/Creat in
104 all cohorts ($P < 0.001$), while UI/Creat varied by ethnicity only in Generation R ($P < 0.05$). Of
105 the dietary predictors, intake of milk and dairy products (per 100 g/day) was positively
106 associated with UI/Creat in all cohorts [ALSPAC ($B = 3.73$, $P < 0.0001$); INMA ($B = 6.92$,
107 $P = 0.002$); Generation R ($B = 2.34$, $P = 0.001$)]. Cohort-specific dietary determinants positively
108 associated with UI/Creat included fish and shellfish in ALSPAC and INMA, and eggs and
109 cereal/cereal products in Generation R.

110 **Conclusions:** The cohort-specific dietary determinants probably reflect not only dietary
111 habits but iodine-fortification policies, hence public-health interventions to improve iodine
112 intake in pregnancy need to be country-specific.

113 **Keywords:** iodine, pregnancy, diet, determinants, milk and dairy products, ALSPAC

114 **Introduction**

115 Iodine is an essential component of the thyroid hormones which are important for optimal
116 fetal and early postnatal neurodevelopment [1, 2]. Mild-to-moderate maternal iodine
117 deficiency in early pregnancy has been associated with suboptimal offspring cognitive
118 outcomes [3–7]. The early stages of pregnancy mark the beginning of crucial fetal brain
119 development processes such as neuron proliferation, migration and differentiation [8].
120 Though these early processes are thyroid hormone-dependent, the fetal thyroid gland is not
121 fully functional until 18-20 weeks, highlighting the importance of thyroid hormone supply
122 from the mother. In this critical early period, the mother therefore needs sufficient iodine
123 intake to maintain optimal thyroid function [1]. As a result of the increased demand for
124 thyroid hormones and other physiological changes associated with pregnancy, pregnant
125 women have a higher iodine requirement than the general population, putting them at greater
126 risk of deficiency [9, 10].

127 As more than 90% of the dietary iodine absorbed is excreted by the kidneys, urinary
128 iodine concentration (UIC) is considered to be a good estimate of recent iodine intake at the
129 population level [11]. In pregnant populations, iodine sufficiency is defined by a median UIC
130 in the range of 150-249 $\mu\text{g/L}$, corresponding to the iodine intake of 250 $\mu\text{g/day}$ recommended
131 by the World Health Organisation (WHO) [12].

132 Determining the main food sources of iodine in pregnancy is essential so that
133 pregnant women can be given information on how to achieve adequate iodine nutrition.
134 Although good food sources of iodine, such as milk, eggs, fish, and, in some countries,
135 iodised salt, are well-known, the consumption patterns of these foods vary between and
136 within populations, as does their iodine content (i.e., as a result of seasonal variation,
137 agricultural practices, and differences in iodine content of soil and water) [13, 14].
138 Consequently, some variation in the importance of different iodine food sources to population

iodine status is expected between countries, hence universal dietary recommendations to increase iodine intake are unlikely to be appropriate.

Considering the negative consequences of iodine deficiency in pregnancy and the fact that many pregnant women worldwide are still iodine-deficient [15], achieving adequate iodine status in the pregnant population is of public-health importance. Data are, however, lacking on the determinants of iodine status in pregnancy in both deficient and sufficient areas; knowledge of such factors would help to identify women likely to have low iodine status.

This study aimed to explore the determinants of iodine status during early pregnancy in three European populations of differing iodine status. The objectives of the study were: (i) to establish whether iodine status during early pregnancy is associated with maternal socio-demographic, anthropometric, lifestyle factors, and pregnancy characteristics; (ii) to determine how maternal iodine status is influenced by dietary intake during pregnancy; (iii) to identify similarities and differences in the main determinants of iodine status between deficient and sufficient pregnant populations.

Methods

Study population

Data from three prospective population-based birth cohorts were used: (i) the Avon Longitudinal Study of Parents and Children (ALSPAC) in the United Kingdom (UK) [16, 17], (ii) Generation R in the Netherlands [18], and (iii) INfancia y Medio Ambiente (INMA) in Spain [19]. In ALSPAC, 14 541 pregnant women living in the former Avon area in the South-West of England, with an expected delivery date between 1st April 1991 and 31st December 1992, were recruited. The study website contains details of all the data that are available through a fully searchable data dictionary and variable search tool [20]. In

Generation R, 9 778 mothers residing in Rotterdam with an expected delivery date from April 2002 to January 2006 were enrolled in the study. In total, 2 150 pregnant women were recruited as part of the INMA Project from three regions in Spain (Valencia, Sabadell and Gipuzkoa), in the period of November 2003 to January 2008.

Ethics

Ethical approval was obtained prior to recruitment from a number of bodies: the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees (ALSPAC), the Medical Ethical Committee of the Erasmus Medical Centre (Generation R), the Ethical Committee of the Municipal Institute of Medical Investigation and the ethical committees of the hospitals involved in the studies (INMA). All participating women provided informed consent.

Selection criteria for the current study

Women were selected for the current study if they had at least one pre-existing measure of urinary iodine concentration in pregnancy [4, 21, 22] or urine samples available for iodine measurement, provided that the child had a measure of intelligence quotient (IQ) for ALSPAC and Generation R. In INMA, iodine was measured in all women with additional urine samples available, irrespective of child-IQ data.

Women with multiple pregnancies, in-vitro fertilisation, known thyroid disease, and/or use of thyroid-related medication were excluded (Fig. 1). We restricted analyses to measures from early pregnancy, the most critical time for iodine-dependent brain development [2, 23], hence for this study, only samples collected ≤ 18 weeks' gestation were included.

Iodine measurements

Urinary iodine concentration (UIC, $\mu\text{g/L}$) was measured in spot-urine samples collected at a median (25-75th percentile) gestational age of 11.0 (8.0-15.0) weeks in ALSPAC, 13.1 (12.1-14.6) weeks in Generation R, and 13.0 (12.4-13.9) weeks in INMA. Gestational week was established using ultrasound examination (Generation R and INMA) or the date of the last menstrual period (ALSPAC). To adjust for variation in intra- and inter-individual daily hydration status [24–27], iodine concentration was corrected by dividing by urinary creatinine concentration to give the iodine-to-creatinine ratio (UI/Creat, $\mu\text{g/g}$). Correcting UIC by use of urinary creatinine concentration reduces intra-individual variation [25], especially in cohorts of the same sex and age-range [i.e., our cohorts were all women of childbearing age (15-44 years)]. The iodine-to-creatinine ratio has previously been used in all three cohorts [4, 7, 21].

In each cohort we reported both the median (25-75th percentile) UIC ($\mu\text{g/L}$) and UI/Creat ($\mu\text{g/g}$). The percentage with UI/Creat $<150 \mu\text{g/g}$ was also reported; this cut-off was informed by the WHO threshold for adequacy in pregnancy (a median UIC $\geq 150 \mu\text{g/L}$) [12] and when corrected for creatinine concentration, has been used in previous research [4, 6, 7].

Laboratory analysis

Urinary creatinine concentration was determined by the Jaffe rate method in all cohorts. Urinary iodine concentration was determined as previously described in detail [4, 22]; a brief description follows.

ALSPAC: Urinary iodine concentration was measured as ^{127}I at the Trace Element Unit, Southampton General Hospital, on a dynamic reaction cell inductively-coupled plasma mass spectrometer (NexION 300D Perkin-Elmer, Beaconsfield). The accuracy of the results was verified using the certified reference material (CRM) Seronorm urine Levels 1 and 2

(Nycomed, Norway), and accuracy was also monitored by measurement of EQUIP samples at regular intervals throughout the analysis. Within-run precision gave a relative standard deviation (RSD) of 0.8% at 42 µg/L, 2.5% at 84 µg/L, 0.6% at 149 µg/L, and 2.0% at 297 µg/L. Between-run precision was 8.7% RSD at 42 µg/L, 6.5% at 84 µg/L, 7.2% at 149 µg/L, and 6.8% at 297 µg/L.

Generation R: Urinary iodine concentration was measured in Radboud University Medical Centre, Nijmegen, the Netherlands, by the Sandell-Kolthoff method. Iodine calibration was performed using the CRM Seronorm urine Levels 1 and 2 (Nycomed, Norway) and four EQUIP samples that were certified for urinary iodine concentration (Centers for Disease Control and Prevention, USA). At a level of 216 µg/L the within-assay precision was 5.1% RSD and the between-assay precision was 14.3% RSD (n= 30).

INMA: Urinary iodine concentration was measured at the Public Health Laboratory Standards, Basque Government Department of Health, Spain (*Acreditación LE/1108 with ISO 15189 for Clinical Laboratories, National Accreditation Entity*), using a paired-ion reversed-phase, high-performance liquid chromatography with electrochemical detection at a silver working electrode (Waters Chromatography, Milford, MA). The accuracy of the results was verified using the CRM Seronorm urine Levels 1 and 2 (Nycomed, Norway) and internal quality control samples. Within-run precision was 4.5% RSD at 50 µg/L, 3.2% at 100 µg/L, and 2.0% at 300 µg/L. Between-run precision was 7.9% RSD at 50 µg/L, 3.5% at 100 µg/L, and 2.5 % at 300 µg/L.

In ALSPAC, there was concern that some urine samples had been contaminated by the use of iodine-containing test strips [28], and therefore samples with UIC >500 µg/L and/or UI/Creat >700 µg/g were excluded (n=412, 11.8%). These cut-offs were based on previous information from ALSPAC and from other studies of UK pregnant women [4, 29,

30]. Some ALSPAC women had multiple urine samples and in cases where one sample was contaminated, the results from the next available uncontaminated sample were used (n=50).

Dietary assessment

Maternal diet in all cohorts was assessed using a food frequency questionnaire (FFQ). This was administered in late pregnancy in ALSPAC (at 32 weeks) and in early pregnancy in Generation R and INMA (at the same time as urine sampling). The FFQ was unquantified in ALSPAC, but was semi-quantitative (SFFQ) in Generation R and INMA. The FFQ was self-administered in ALSPAC and Generation R, and administered by trained interviewers in INMA.

ALSPAC: Detailed information about the design of the questionnaire can be found elsewhere [31, 32]. Briefly, women were asked to indicate how frequently “nowadays” they consumed 43 food groups and individual foods with five predefined frequency categories, ranging from “never or rarely” to “more than once a day”. Portion sizes were not included in the questionnaire.

Generation R: SFFQ in Generation R represented an adapted version of a validated SFFQ in elderly subjects [33]. In summary, it contained a list of 293 foods and asked about frequency of consumption in the past three months, mostly therefore reflecting the first trimester. Questions about portion size, estimated using food photographs or Dutch household measures, methods of preparation, and additions to foods were also included [3, 34]. The food list had previously been reduced to 17 main food groups [35], based on the European Prospective Investigation into Cancer and Nutrition SOFT classification (EPIC-SOFT) [36].

INMA: SFFQ was based on an expanded adaptation of a 61-item SFFQ by Willet and colleagues [37] that was developed and validated [38, 39]. To summarise, women were asked

how often, on average, they had consumed a specified standard portion of 100 food items in early pregnancy (since their last menstrual period until the time of the interview at ~12 weeks), using nine frequency categories, ranging from “never or less than once a month” to “six or more times per day”.

For the current study, in each cohort, food intake (g/day) was calculated by multiplying the frequency of consumption by an average standard portion (for the ALSPAC FFQ) or a specified portion (for the SFFQ in Generation R and INMA) of that food. Foods were then classified into food groups and the amounts of individual food items consumed were summed accordingly. To facilitate comparison between cohorts, the classification of food groups used in Generation R [36] was used as a template. This required the formation of new food groups in ALSPAC, while the pre-existing food groups were used in Generation R and INMA, with some minor modifications to make them comparable. The definitions of food groups in the individual cohorts are shown in Supplemental Table 1. Separate information about the consumption of table salt and use of iodine-fortified table salt was collected only in the INMA cohort.

Iodine supplement use

Detailed data on the use of potassium iodide and/or iodine-containing multivitamin/mineral supplements during pregnancy were collected only in INMA. For this analysis, we used data on iodine supplements from pre-pregnancy until the end of the first trimester, expressed as mean iodine intake from supplements. In our ALSPAC sample, only two women took a kelp supplement and one took a potassium-iodide supplement; they were excluded from the analysis. Data on the use of iodine-containing multivitamin supplements from pre-conception until enrolment (at time of urine sample collection) were available only for a sub-set of our

287 Generation R sample (n=381); women who took an iodine-containing supplement (n=345)
 288 were excluded in sensitivity analyses.

289

290 *Maternal and pregnancy characteristics*

291 Information on pregnancy characteristics, anthropometrics, socio-demographic and
 292 environmental exposures was collected by means of questionnaires or extracted from
 293 obstetric medical records. Discrete variables were re-categorised, where possible, to facilitate
 294 comparisons between cohorts. Exposure factors in these analyses can be classified into three
 295 groups: (i) maternal factors: maternal age (ALSPAC: at last menstrual period; Generation R,
 296 INMA: at urine collection); pre-pregnancy body mass index (BMI, kg/m²); ethnicity
 297 (ALSPAC: White/non-white; Generation R: Dutch/non-Dutch, where non-Dutch =
 298 Indonesian, Cape Verdian, Moroccan, Dutch Antilles, Surinamese, Turkish, other non-
 299 Western, Asian, other Western; INMA: Spanish/non-Spanish, where non-Spanish = Latin
 300 American, European, others); parity (zero, one, two or more); smoking status (never smoked,
 301 smoked initially or until pregnancy was known, continued smoking); and alcohol
 302 consumption during pregnancy (yes/no); (ii) markers of socio-economic status: education
 303 level (ALSPAC: low = no qualification, certificate of secondary education, or vocational;
 304 medium = O level or A level; high = a degree; Generation R: low = no education or primary;
 305 medium = secondary phase 1 and 2; high = higher phase 1 and 2; INMA: low = no education,
 306 unfinished primary, or primary; medium = secondary; high = university degree); monthly net
 307 household income (Generation R: low <€1200, medium = €1200-2200, high >€2200); home
 308 ownership (ALSPAC: owned/mortgaged, private/other rented, council rented); crowding
 309 index (ALSPAC: ≤1 person per room, +1 person per room); family adversity index
 310 [ALSPAC: 18-item measure of hardships during pregnancy [40], categorised into: no
 311 adversities, mild (0-2), severe (≥3) [41]]; life event score (ALSPAC: exposure to stressful

situations during pregnancy); marital status (ALSPAC: never married, married, other; Generation R: unmarried/married); and living with father's child/partner (ALSPAC, INMA: yes/no); (iii) child factors: child's sex (male/female).

Statistical analyses

To study the associations of maternal characteristics and diet with iodine status, we used UI/Creat as a measure of individual iodine status. UI/Creat was not normally distributed but right-skewed. To meet parametric-test assumptions, UI/Creat was transformed using the natural logarithm. Outliers were assessed by visual inspection of box-plots. We assessed non-linearity of the associations of each continuous independent variable with UI/Creat by adding their squared term to the regression models, and also by plotting each potential determinant variable against UI/Creat and comparing the fit (R^2) of a linear vs. quadratic function through the data points.

Multiple linear regression models with log-transformed UI/Creat as the dependent variable were performed for each cohort using two models: Model 1 included maternal and pregnancy factors, markers of socio-economic status and child's sex as independent variables; Model 2 included variables from Model 1 plus dietary intake of food groups.

Analyses of the dietary influences on UI/Creat (Model 2) were also adjusted for estimated energy intake (kcal/day). Effect estimates (unstandardised B coefficients) for food groups were expressed per 100 g and also per portion (g). The increase in the geometric mean of UI/Creat per 100 g and per portion increase in intake and its 95% confidence interval (CI) were calculated by back-transformation [exponentiation (EXP)] of the effect estimates and 95% CIs from logarithmic scale. The following formulae were used for the back-transformations: *Effect estimate per 100 g* = $EXP(intercept + B*100) - EXP(intercept)$; *Lower 95% CI* = $EXP[intercept + (B*100) - (1.96*standard\ error\ of\ estimate\ (SEE)*100)]$

337 $-EXP(intercept)$; Upper 95% CI = $EXP[intercept + (B*100) + (1.96*SEE*100)] - EXP$
 338 $(intercept)$. For calculations per portion size, the multiplication by 100 in the formulae is
 339 replaced with the portion size (g) for each food group accordingly. When calculating the
 340 effect estimates from Model 2, all categorical covariates were set to their reference group and
 341 the continuous covariates gestational week, maternal age, BMI and energy intake were
 342 centred to their means.

343 We conducted four types of sensitivity analyses: (i) under-reporting: to account for
 344 potential under-reporting of energy intake in Model 2, in all cohorts, we excluded women
 345 with energy intakes below the 5th percentile; (ii) iodine supplements: we adjusted Model 2 for
 346 iodine-supplement use in INMA (as the only cohort with complete iodine-supplement data)
 347 and excluded iodine-supplement-users in Generation R (as data were available only for a sub-
 348 set of the sample); (iii) gestational age: as the median gestational week at urine sampling was
 349 later in INMA and Generation R than in ALSPAC, we performed sensitivity analyses to test
 350 the associations with iodine status in the first trimester, using samples collected up to 13
 351 weeks; (iv) covariate creatinine-adjustment method: since age, BMI and ethnicity are known
 352 predictors of urinary creatinine concentration [42] and urinary creatinine can also vary during
 353 gestation [43], this could result in spurious associations of these variables with the ratio of
 354 UI/Creat; we therefore performed sensitivity analyses for these variables (BMI, age,
 355 ethnicity, and gestational age) using Model 1 and 2 with the (natural) log-transformed UIC
 356 ($\mu\text{g/L}$) unadjusted for creatinine as our dependent variable and added creatinine concentration
 357 as a separate independent variable to the models. This method has been recommended
 358 previously [42] and ensures that UIC is adjusted for dilution by creatinine concentration
 359 while the associations of the other variables with UIC are independent of urinary creatinine.

360 All analyses were conducted using multiply-imputed data in order to account for
 361 missing data on socio-demographic variables. Multiple imputation was performed using the

automatic method in SPSS. A total of 20 datasets were generated and analysed using standard multiple imputation procedures [44]. Detailed information about the imputed variables is provided in the Electronic Supplementary Material. Missing FFQ data were not imputed, as diet has a wide inter-person variability hence imputation of dietary data would not be sufficiently accurate. All statistical analyses were performed with IBM SPSS Statistics version 24.0 (IBM Corp., Armonk, NY, USA). Values were considered statistically significant at $P < 0.05$.

Results

Sample characteristics

After exclusions, the final study population comprised 6566 pregnant women: 2852 from ALSPAC, 2254 from Generation R, and 1460 from INMA (Fig. 1). Descriptive statistics of mothers by cohort are shown in Table 1. Maternal age varied across cohorts, with women in ALSPAC having a lower mean $\pm$ standard deviation (SD) age than women in INMA [28.7 (± 4.5) vs. 31.4 (± 4.1) years, respectively]. Median BMI was within the healthy range in all cohorts (Table 1). The majority of mothers defined themselves as White in ALSPAC (98.2%), Spanish in INMA (91.8%), while slightly more than half of the women in Generation R said they were non-Dutch (51.4%). Most women were nulliparous and non-smokers, with similar proportions between cohorts.

Iodine status

The median UI/Creat (25-75th percentile) was 121 (81-193) $\mu\text{g/g}$ in women from the UK (62.8% $< 150 \mu\text{g/g}$), 151 (96-255) $\mu\text{g/g}$ in women from Spain (49.5% $< 150 \mu\text{g/g}$), and 210 (140-303) $\mu\text{g/g}$ in women from the Netherlands (28.8% $< 150 \mu\text{g/g}$) (Table 2).

Association with socio-demographic and lifestyle factors

In multivariable models (adjusted Models 1 and 2), gestational week of urine sample, maternal age, and BMI were associated with UI/Creat in all three cohorts. Gestational week at urine sampling (≤ 18 weeks) was positively associated with UI/Creat in ALSPAC ($B=0.051$, $P<0.0001$) but not in the other two cohorts (Table 3). However, in sensitivity analyses restricted to samples collected up to 13 weeks, there was an association of gestational week with UI/Creat in all three cohorts (ALSPAC: $B=0.029$, $P<0.001$, $n=1951$; Generation R: $B=0.031$, $P=0.049$, $n=1094$; INMA: $B=0.079$, $P=0.045$, $n=747$) with a larger effect size in Generation R and INMA than in the analyses up to 18 weeks. In the sensitivity analysis using covariate creatinine-adjustment, the results for the association of gestational week up to 18 weeks with UIC did not substantially differ from those with UI/Creat in ALSPAC and INMA, while in Generation R the effect size was higher, reaching statistical significance (Supplemental Table 3).

There was a positive association of maternal age with UI/Creat (ALSPAC: $B=0.014$, $P<0.0001$; Generation R: $B=0.018$, $P<0.0001$; INMA: $B=0.020$, $P=0.0001$). After further adjusting for maternal diet and energy intake estimated from the FFQs (Model 2), the effect size of age was attenuated by 16 - 30 % across cohorts, but remained statistically significant (Table 3). The positive association with age remained in all cohorts, except in Model 2 for ALSPAC, when using the covariate creatinine-adjustment method of UIC (Supplemental Table 3).

There was a negative association of BMI with UI/Creat (ALSPAC: $B= -0.013$, $P=0.0001$; Generation R: $B= -0.011$, $P<0.0001$; INMA: $B= -0.013$, $P=0.005$), which, after adjustment for maternal diet and energy intake remained statistically significant (Table 3, Model 2). However, BMI was not associated with UIC ($\mu\text{g/L}$) with the covariate creatinine-

adjustment method in Generation R and INMA but remained negatively associated with UIC in ALSPAC, though with a lower effect size (Supplemental Table 3).

Cohort-specific socio-demographic and lifestyle factors were identified as determinants of iodine status. In Generation R, maternal ethnicity and smoking were associated with UI/Creat (Table 3). Compared to the Dutch women, Moroccan, Turkish and other non-Western women had a higher UI/Creat, whereas Surinamese and those from the Dutch Antilles had a lower UI/Creat. Some of these effects were attenuated after accounting for maternal diet in Model 2 (Table 3). Similarly to UI/Creat, UIC (with covariate creatinine-adjustment) also differed by ethnicity; Moroccan, Turkish and other non-Western women still had a significantly higher UIC than the Dutch, while the UICs of Surinamese and Dutch Antilles women did not significantly differ from those of the Dutch women (Supplemental Table 3). Generation R women who reported smoking *vs.* those who never smoked had a lower UI/Creat, which remained statistically significant after adjustment for maternal diet (Table 3, Model 1 and 2). In ALSPAC, family adversity index (severe *vs.* none; $B = -0.100$, $P = 0.016$), and marital status (married *vs.* never-married; $B = 0.095$, $P = 0.015$) were associated with UI/Creat, even after adjusting for maternal diet (Table 3). Results for all predictors included in the multivariable models are presented in Supplemental Table 2.

Dietary influences on iodine status

As not all women with urinary iodine measurements before 18 weeks had also completed an FFQ, numbers for these analyses were lower for all cohorts (Fig. 1): ALSPAC ($n = 2710$), Generation R ($n = 1580$), INMA ($n = 1446$). Descriptive statistics of dietary intakes of food groups for pregnant women in each cohort are presented in Supplemental Table 4.

“Milk and dairy products” was the only food group positively associated with UI/Creat in all three cohorts: ALSPAC ($B = 3.73$, $P < 0.0001$); Generation R ($B = 2.34$,

$P=0.001$); INMA ($B=6.92$, $P=0.002$) (Supplemental Table 5). Based on the fully-adjusted models, one-portion increase in consumption of milk and dairy products (e.g., a glass of milk, 200 g) [45] was associated with a 5-14 $\mu\text{g/g}$ increase in the geometric mean of UI/Creat (Table 4). Intake of “fish and shellfish” was positively associated with UI/Creat in pregnant women in Spain (INMA) and the UK (ALSPAC), though the effect size per 100 g was more than five times larger in INMA women ($B=28.04$, $P=0.029$ vs. $B=5.10$, $P=0.017$).

Cohort-specific dietary determinants were also identified (Table 4). In ALSPAC, consumption of fruit ($P=0.012$), cakes and confectionery ($P=0.028$) were positively associated with UI/Creat. Intake of cereals and cereal products ($P<0.0001$), eggs ($P=0.0001$), nuts and seeds ($P=0.001$), and added fat ($P=0.022$) were all positively associated with UI/Creat in Generation R. Higher salt intake (including iodised salt) (per 1 g) was associated with higher UI/Creat ($P=0.003$), while increasing meat intake was associated with lower UI/Creat ($P=0.002$) in INMA.

Excluding women with extremely low energy intakes ($< 5^{\text{th}}$ percentile) did not considerably change our results, except for small changes in INMA where the association between fish intake and UI/Creat was borderline significant but the effect size remained relatively unchanged (*data not shown*). In INMA, mean daily iodine-supplement intake was an important determinant of UI/Creat (Standardised $B=0.235$, $P<0.0001$). Adjusting for iodine-supplement use in INMA attenuated the result for maternal age ($B=0.012$, $P=0.018$) but did not change the rest of the results substantially (*data not shown*). In Generation R, the exclusion of iodine-supplement users resulted in higher effect estimates for some ethnic groups (e.g., Moroccan, Turkish and other Western); the associations between intake of non-alcoholic and alcoholic beverages and UI/Creat reached statistical significance but the effect sizes remained unchanged; the rest of the results remained relatively unchanged (*data not shown*).

Discussion

In this study of pregnant women, several dietary (milk and dairy products) and maternal factors (maternal age, BMI, gestational week) were associated with UI/Creat across all cohorts in adjusted models. Furthermore, important cohort-specific dietary determinants were identified, such as fish intake in ALSPAC (UK), egg and cereal/cereal product intake in Generation R (Netherlands), and fish, salt, and meat intake in INMA (Spain).

The population of pregnant women from the Netherlands was iodine-sufficient, while populations from the UK and Spain were mildly-to-moderately deficient. It should be noted that the ALSPAC median-UIC value, although from samples that are nearly 25 years old, is almost identical to the UK value from the 2017 Global Scorecard for Iodine Nutrition in pregnant women i.e., 95 $\mu\text{g/L}$ vs. 99 $\mu\text{g/L}$ [15].

The differences in iodine status between the countries may be partly explained by differing use of iodised salt. The iodised salt penetration rate in households and the food industry (e.g., bread-making) in the Netherlands has been estimated as 60% and 70%, respectively, while a 16% penetration rate has been reported in Spanish households [46]. By contrast, iodisation of salt was never common in the UK [47], and even nowadays its availability is very limited (21.5%); furthermore, the iodine concentration of the major UK brand is low [48].

Association with socio-demographic and lifestyle factors

Gestational week of urine samples was positively associated with a UI/Creat in ALSPAC but not in the other cohorts. This may be because ALSPAC samples were collected at an earlier gestational age than in the other cohorts; if the greatest increase in urinary iodine excretion is in the first weeks of pregnancy, the effect of gestational age may be attenuated in later

samples (i.e., up to 18 weeks as in our study). Indeed in sensitivity analyses, when we restricted analysis to samples collected up to 13 weeks, the effect size was higher in the other two cohorts. The positive association between gestational week of urine sample with UI/Creat in early pregnancy could be attributed to an increase in glomerular filtration rate and subsequent renal iodine loss in early pregnancy [49, 50], though data are conflicting as studies report both an increase and a decrease in the UI/Creat [51–54]. As urinary iodine excretion in early vs. late pregnancy stages might be higher as a result of increased renal clearance of iodine, using early measurement to assess status might overestimate true intake [55]. Creatinine clearance has also been shown to vary during gestation [43], which could have biased our results when using UI/Creat. However, gestational week was also positively associated with UIC ($\mu\text{g/L}$) alone when including creatinine separately in the model.

In all three cohorts, BMI was negatively associated with UI/Creat. In a study in non-pregnant adults [56], creatinine clearance was positively associated with BMI, independent of adiposity, and a positive association with lean thigh-tissue area was reported; this suggests that the association of creatinine with BMI might be explained by lean body mass. The negative association between BMI and UI/Creat that we report may therefore be partly explained by the use of creatinine adjustment and may highlight potential issues with the use of this measure of iodine status [42, 57]. Indeed in sensitivity analysis, we did not find an association of BMI with UIC alone in two of the cohorts when we adjusted for creatinine by including it as a covariate in the model instead of using the ratio of UI/Creat.

There was a positive association of maternal age with UI/Creat in all cohorts. Age was still significantly associated with UI/Creat after adjustment for dietary intake though the effect size was attenuated; it was further reduced when iodine-supplement use in INMA was accounted for, suggesting that some of the effect of age could be explained by diet and supplement use in older women. Similarly to BMI, age is a known predictor of urinary

creatinine [42]. However, in sensitivity analysis when adjusting for creatinine separately in the model, unlike BMI, age remained positively associated with UIC but with a lower effect size.

The iodine status of pregnant women from the Netherlands varied by ethnic origin, with higher iodine status in Moroccan, Turkish and other non-Western women, and lower iodine status in Surinamese and Dutch Antilles women, even after adjusting for socio-demographic factors. Variation in diet may partially explain these differences as some of the effect estimates were attenuated when adjusting for dietary intake. Alternatively, there may be genetic variability in iodine or creatinine clearance [42, 58]. Indeed, we found that when using UIC and adjusting for urinary creatinine separately in the model, UIC varied similarly to UI/Creat between the ethnic groups though some of the associations were attenuated and were no longer significant (e.g., for Surinamese and Dutch Antilles women). It should be noted that numbers in some of these ethnic-group categories were relatively small ($n < 100$). Ethnicity was not significantly associated with iodine status in ALSPAC or INMA, but this may reflect the small sample sizes of other ethnic groups in these cohorts. Ethnic differences in iodine status could help to identify subgroups at high-risk for iodine insufficiency; in countries with a large proportion of diverse ethnic groups, culturally-specific approaches to improve dietary adequacy may be more suitable than a single solution for the whole population.

Dietary influences on iodine status

The only food group that was positively associated with UI/Creat in all three cohorts was “milk and dairy products”, demonstrating their significant role as an important dietary determinant of iodine status in pregnancy. This finding is consistent with the results of other

studies in European pregnant women, i.e., Norway [59, 60], Iceland [61], Italy [62], Spain [63], and the UK [52, 64], as well as studies of pregnant women in Australia [65].

Based on our model, a portion of “milk and dairy products” equivalent to a glass of milk (200 g) was associated with 5 to 14 $\mu\text{g/g}$ increase in UI/Creat across cohorts. The effect sizes for milk differed between cohorts; the highest effect size was in Spain, while the lowest was in the Netherlands. This is in line with the milk-iodine concentration in each country [i.e., $\sim 26 \mu\text{g}$ in Spain [66], $\sim 15 \mu\text{g}$ in the Netherlands [67], and $15 \mu\text{g}$ in the UK in 1990/1991 [68]]. The results might be different if repeated now as the iodine concentration in UK milk is higher than estimated when ALSPAC women were recruited in 1990/1991, i.e. 427 vs. $150 \mu\text{g/kg}$ [68, 69].

Consumption of eggs and fish were positively associated with UI/Creat, though not consistently across the cohorts; the association with egg intake was statistically significant only in the Netherlands, while intake of fish and shellfish was associated with UI/Creat only in Spain and the UK. The effect size for eggs was higher than that expected, given their iodine content, but similar to values reported previously [60]. These higher values may reflect the consumption of eggs with salt in the Netherlands, which is likely to be iodised. An average portion of fish (120 g) was associated with some 6 to 34 $\mu\text{g/g}$ increase in UI/Creat, across cohorts. Variation in the effect-size could partly reflect the variability in average fish consumption, particularly of white fish, which is a good iodine source [70] (e.g., pregnant women in Spain consumed a higher amount of white fish daily than did women in the UK). The wide CIs around the estimates probably relate to the variability in fish-iodine concentration [14] [i.e., the proportion of oily fish in the food group has a much lower iodine concentration [70]] and to the irregular nature of fish consumption which may not be captured by a spot-UI/Creat.

Consumption of cereals and cereal products was a statistically significant determinant of UI/Creat only in the Netherlands; this association is probably driven by consumption of bread which is made with iodised salt in the Netherlands [71], whereas without iodised salt, bread has a low iodine concentration and was not found to be a predictor of iodine status in the UK or Spanish cohorts. Intake of iodised table salt was measured only in pregnant women from Spain; 1 g of salt was associated with around 32 µg/g increase in UI/Creat. Both of these results suggest that iodised salt consumed either discretionarily (e.g., as table salt in Spain), or as part of processed foods (e.g., in bread in the Netherlands) is an important dietary determinant of iodine status. We observed the highest UI/Creat in the Netherlands, followed by Spain, and the lowest in the UK, suggesting that iodised-salt use might be a key dietary factor with large influence on iodine status.

Surprisingly, meat intake was negatively associated with UI/Creat in the Spanish cohort, even after controlling for socio-demographic variables. Higher urinary creatinine concentrations have been reported in individuals with a high-meat diet, which may account for the lower ratio [72, 73].

Some of our other food-group associations are difficult to explain and may be chance findings e.g. the positive association with fruit, and cakes/confectionary in the UK. However, fruit intake was also associated with urinary iodine excretion in Norwegian pregnant women [59] and the finding might warrant further investigation.

Although we have observed significant increases in UI/Creat, the effect sizes were relatively small and the total explained variance in UI/Creat (R^2) was low. This is probably explained by the large day-to-day variability in iodine intake which cannot be captured in a single spot-urine sample and the measurement errors of dietary assessment methods in capturing habitual iodine intake. Moreover, as a result of the physiological changes occurring

during gestation (e.g., increased renal iodine clearance), pregnancy may not represent a steady state of iodine metabolism (intake vs. excretion) [10].

Strengths and limitations

Strengths of our study include the large sample size, the inclusion of pregnant women from three geographically and culturally different populations. Furthermore, we explored associations with the entire range of food groups, rather than focusing on a few groups or isolated foods as in previous studies [62, 63]. However, our study also has a number of limitations. Firstly, the use of an FFQ for measuring diet and the use of spot-urine samples for estimating individual iodine intake have their methodological disadvantages [24–26, 74]. Although we only had spot-urine samples which might not reflect individual iodine intake or status, we used UI/Creat which has been shown as a valid alternative to the 24-hour urinary iodine excretion when used in homogenous population groups [75]. Although urinary iodine concentration was measured in three different laboratories using different assays, each laboratory ensured accuracy by use of certified reference materials. Exploring associations between data from an FFQ and spot-UI/Creat can also be problematic, as an FFQ is designed to measure habitual diet [76], whereas a spot-urine sample reflects iodine intake in the last 24-48 hours [11]; this may be reflected in the more consistent association between UI/Creat and intake of daily food items (e.g., milk) than those infrequently consumed (e.g., fish). However, our large cohort sample size would have helped to overcome this limitation to some extent. Secondly, it should be noted that in ALSPAC there was a time difference in the administration of the FFQ (at 32 weeks) and urine-sample collection (≤ 18 weeks), hence the FFQ might not reflect diet during early pregnancy. Thirdly, although we harmonised the classification of foods into food groups, there were some differences in the foods included in each group which may explain variation in effect sizes between cohorts (i.e., in ALSPAC the

dairy food group included milk and cheese, while in INMA and Generation R, ice cream, yoghurt, cream were also included). Fourthly, we had incomplete, or no, data on iodine supplement use in two of the cohorts (Generation R and ALSPAC), which could be an important determinant that we were unable to evaluate; however this is unlikely to be a limitation in ALSPAC as it is unlikely that women would have taken an iodine-containing supplement in the early 1990s. Lastly, we measured urinary iodine only in women with available child IQ data in two of the cohorts which might have created bias.

Conclusions

Various maternal characteristics and dietary habits were associated with UI/Creat in pregnancy, some of which were population-specific. For that reason, universal interventions and dietary recommendations to improve the iodine intake of pregnant women might not be appropriate; a country-specific approach needs to be adopted. Between countries, but also within countries with a large proportion of different ethnic groups, culturally-specific recommendations are probably necessary. Achieving and maintaining iodine sufficiency in populations requires monitoring dietary determinants of iodine status so that appropriate action can be taken, where necessary.

Ethical standards

All studies in these analysis have been approved by the appropriate ethics committees and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. Detailed information is included as part of the main text (see Methods, sub-section Ethics).

633 **Conflict of interest**

634 The authors declare that they have no conflict of interest

635

636 **Authors' contributions**

637 MD performed statistical analyses, interpreted the data, and wrote the first version of the

638 manuscript. MPR, SCB, RPP, TIMK, MG, and DL contributed to statistical analyses,

639 interpretation of the data, and writing of the manuscript. JV, LG, ME, JI, and JS helped with

640 writing of the manuscript. All authors read and approved the final manuscript.

References

1. de Escobar GM, Obregón MJ, del Rey FE (2004) Maternal thyroid hormones early in pregnancy and fetal brain development. *Best Pract Res Clin Endocrinol Metab* 18:225–248 . doi: 10.1016/j.beem.2004.03.012
2. Glinioer D (2007) The importance of iodine nutrition during pregnancy. *Public Health Nutr* 10:1542–1546 . doi: 10.1017/S1368980007360886
3. van Mil NH, Tiemeier H, Bongers-Schokking JJ, Ghassabian A, Hofman A, Hooijkaas H, Jaddoe VW V., de Muinck Keizer-Schrama SM, Steegers EAP, Visser TJ, Visser W, Ross HA, Verhulst FC, de Rijke YB, Steegers-Theunissen RPM (2012) Low Urinary Iodine Excretion during Early Pregnancy Is Associated with Alterations in Executive Functioning in Children. *J Nutr* 142:2167–2174 . doi: 10.3945/jn.112.161950
4. Bath SC, Steer CD, Golding J, Emmett P, Rayman MP (2013) Effect of inadequate iodine status in UK pregnant women on cognitive outcomes in their children: Results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Lancet* 382:331–337 . doi: 10.1016/S0140-6736(13)60436-5
5. Hynes KL, Otahal P, Hay I, Burgess JR (2013) Mild Iodine Deficiency During Pregnancy Is Associated With Reduced Educational Outcomes in the Offspring: 9-Year Follow-up of the Gestational Iodine Cohort. *J Clin Endocrinol Metab* 98:1954–1962 . doi: 10.1210/jc.2012-4249
6. Hynes K, Otahal P, Burgess J, Oddy W, Hay I (2017) Reduced Educational Outcomes Persist into Adolescence Following Mild Iodine Deficiency in Utero, Despite Adequacy in Childhood: 15-Year Follow-Up of the Gestational Iodine Cohort Investigating Auditory Processing Speed and Working Memory. *Nutrients* 9:1354 . doi: 10.3390/nu9121354
7. Murcia M, Espada M, Julvez J, Llop S, Lopez-Espinosa M-J, Vioque J, Basterrechea M,

- 666 Riaño I, González L, Alvarez-Pedrerol M, Tardón A, Ibarluzea J, Rebagliato M (2017)
 667 Iodine intake from supplements and diet during pregnancy and child cognitive and motor
 668 development: the INMA Mother and Child Cohort Study. *J Epidemiol Community*
 669 *Health* 0:1–7 . doi: 10.1136/jech-2017-209830
- 670 8. Williams GR (2008) Neurodevelopmental and Neurophysiological Actions of Thyroid
 671 Hormone. *J Neuroendocrinol* 20:784–794 . doi: 10.1111/j.1365-2826.2008.01733.x
- 672 9. Glinoe D (2004) The regulation of thyroid function during normal pregnancy:
 673 importance of the iodine nutrition status. *Best Pract Res Clin Endocrinol Metab* 18:133–
 674 152 . doi: 10.1016/J.BEEM.2004.03.001
- 675 10. Delange F (2007) Iodine requirements during pregnancy, lactation and the neonatal
 676 period and indicators of optimal iodine nutrition. *Public Health Nutr* 10:1571–1580 .
 677 doi: 10.1017/S1368980007360941
- 678 11. Zimmermann MB (2008) Methods to assess iron and iodine status. *Br J Nutr* 99:S2–S9
 679 . doi: 10.1017/S000711450800679X
- 680 12. World Health Organisation (2007) Assessment of iodine deficiency disorders and
 681 monitoring their elimination. A guide for programme managers.
 682 http://apps.who.int/iris/bitstream/10665/43781/1/9789241595827_eng.pdf. Accessed
 683 20 Sep 2017
- 684 13. Rasmussen LB, Andersen S, Ovesen L, Laurberg P (2009) Iodine Intake and Food
 685 Choice. In: Preedy VR, Burrow GN, Watson RR (eds) *Comprehensive Handbook of*
 686 *Iodine: Nutritional, Biochemical, Pathological and Therapeutic Aspects*. Elsevier,
 687 London, pp 331–337
- 688 14. Haldimann M, Alt A, Blanc A, Blondeau K (2005) Iodine content of food groups. *J Food*
 689 *Compos Anal* 18:461–471 . doi: 10.1016/j.jfca.2004.06.003
- 690 15. Iodine Global Network (2017) Global Scorecard of Iodine Nutrition in 2017 in the

general population and in pregnant women (PW).

http://www.ign.org/cm_data/IGN_Global_Scorecard_AllPop_and_PW_May2017.pdf.

Accessed 24 Sep 2017

16. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, Molloy L, Ness A, Ring S, Smith GD (2013) Cohort profile: The “Children of the 90s”-The index offspring of the avon longitudinal study of parents and children. *Int J Epidemiol* 42:111–127 . doi: 10.1093/ije/dys064
17. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey smith G, Henderson J, Macleod J, Molloy L, Ness A, Ring S, Nelson SM, Lawlor DA (2013) Cohort profile: The avon longitudinal study of parents and children: ALSPAC mothers cohort. *Int J Epidemiol* 42:97–110 . doi: 10.1093/ije/dys066
18. Kooijman MN, Kruithof CJ, van Duijn CM, Duijts L, Franco OH, van IJzendoorn MH, de Jongste JC, Klaver CCW, van der Lugt A, Mackenbach JP, Moll HA, Peeters RP, Raat H, Rings EHHM, Rivadeneira F, van der Schroeff MP, Steegers EAP, Tiemeier H, Uitterlinden AG, Verhulst FC, Wolvius E, Felix JF, Jaddoe VWV (2016) The Generation R Study: design and cohort update 2017. *Eur J Epidemiol* 31:1243–1264 . doi: 10.1007/s10654-016-0224-9
19. Guxens M, Ballester F, Espada M, Fernández MF, Grimalt JO, Ibarluzea J, Olea N, Rebagliato M, Tardón A, Torrent M, Vioque J, Vrijheid M, Sunyer J (2012) Cohort Profile: The INMA—INfancia y Medio Ambiente—(Environment and Childhood) Project. *Int J Epidemiol* 41:930–940 . doi: 10.1093/ije/dyr054
20. ALSPAC (2018) Data Dictionary. <http://www.bristol.ac.uk/alspac/researchers/our-data/>. Accessed 20 Mar 2018
21. Ghassabian A, Steenweg-de Graaff J, Peeters RP, Ross HA, Jaddoe VW, Hofman A, Verhulst FC, White T, Tiemeier H (2014) Maternal urinary iodine concentration in

pregnancy and children's cognition: results from a population-based birth cohort in an iodine-sufficient area. *BMJ Open* 4:e005520–e005520 . doi: 10.1136/bmjopen-2014-005520

22. Murcia M, Rebagliato M, Espada M, Vioque J, Santa Marina L, Alvarez-Pedrerol M, Lopez-Espinosa M-J, León G, Iñiguez C, Basterrechea M, Guxens M, Lertxundi A, Perales A, Ballester F, Sunyer J, INMA Study Group (2010) Iodine intake in a population of pregnant women: INMA mother and child cohort study, Spain. *J Epidemiol Community Health* 64:1094–9 . doi: 10.1136/jech.2009.092593
23. Stiles J, Jernigan TL (2010) The basics of brain development. *Neuropsychol Rev* 20:327–48 . doi: 10.1007/s11065-010-9148-4
24. Vejbjerg P, Knudsen N, Perrild H, Laurberg P, Andersen S, Rasmussen LB, Ovesen L, Jørgensen T (2009) Estimation of Iodine Intake from Various Urinary Iodine Measurements in Population Studies. *Thyroid* 19:1281–1286 . doi: 10.1089/thy.2009.0094
25. König F, Andersson M, Hotz K, Aeberli I, Zimmermann MB (2011) Ten repeat collections for urinary iodine from spot samples or 24-hour samples are needed to reliably estimate individual iodine status in women. *J Nutr* 141:2049–54 . doi: 10.3945/jn.111.144071
26. Andersen S, Karmisholt J, Pedersen KM, Laurberg P (2008) Reliability of studies of iodine intake and recommendations for number of samples in groups and in individuals. *Br J Nutr* 99:813–8 . doi: 10.1017/S0007114507842292
27. Rasmussen LB, Ovesen L, Christiansen E (1999) Day-to-day and within-day variation in urinary iodine excretion. *Eur J Clin Nutr* 53:401–7
28. Pearce EN, Lazarus JH, Smyth PP, He X, Smith DF, Pino S, Braverman LE (2009) Urine Test Strips as a Source of Iodine Contamination. *Thyroid* 19: . doi:

- 741 10.1089=thy.2009.0120
- 742 29. Pearce EN, Lazarus JH, Smyth PPA, He X, Dall’Amico D, Parkes AB, Burns R, Smith
 743 DF, Maina A, Bestwick JP, Jooman M, Leung AM, Braverman LE (2010) Perchlorate
 744 and Thiocyanate Exposure and Thyroid Function in First-Trimester Pregnant Women. *J*
 745 *Clin Endocrinol Metab* 95:3207–3215 . doi: 10.1210/jc.2010-0014
- 746 30. Bath SC, Pop VJM, Furmidge-Owen VL, Broeren MAC, Rayman MP (2017)
 747 Thyroglobulin as a Functional Biomarker of Iodine Status in a Cohort Study of Pregnant
 748 Women in the United Kingdom. *Thyroid* 27:426–433 . doi: 10.1089/thy.2016.0322
- 749 31. Emmett P, Symes C, Braddon F, Heaton K (1992) Validation of a new questionnaire for
 750 assessing habitual intakes of starch, non-starch poly-saccharides, sugars and alcohol. *J*
 751 *Hum Nutr Diet* 5:245–254
- 752 32. Rogers I, Emmett P, ALSPAC Study Team (1998) Diet during pregnancy in a population
 753 of pregnant women in South West England. *Eur J Clin Nutr* 52:246–250
- 754 33. Klipstein-Grobusch K, den Breeijen JH, Goldbohm R a, Geleijnse JM, Hofman A,
 755 Grobbee DE, Witteman JC (1998) Dietary assessment in the elderly: validation of a
 756 semiquantitative food frequency questionnaire. *Eur J Clin Nutr* 52:588–596 . doi:
 757 10.1038/sj.ejcn.1600611
- 758 34. Timmermans S, Steegers-Theunissen RPM, Vujkovic M, Bakker R, Den Breeijen H,
 759 Raat H, Russcher H, Lindemans J, Hofman A, Jaddoe VW V, Steegers EAP (2011)
 760 Major dietary patterns and blood pressure patterns during pregnancy: The Generation R
 761 Study. *Am J Obstet Gynecol* 205:337.e1-337.e12 . doi: 10.1016/j.ajog.2011.05.013
- 762 35. Steenweg-de Graaff J, Tiemeier H, Steegers-Theunissen RPM, Hofman A, Jaddoe VW
 763 V, Verhulst FC, Roza SJ (2014) Maternal dietary patterns during pregnancy and child
 764 internalising and externalising problems. The Generation R Study. *Clin Nutr* 33:115–
 765 121 . doi: 10.1016/j.clnu.2013.03.002

- 766 36. Slimani N, Fahey M, Welch A, Wirfält E, Stripp C, Bergström E, Linseisen J, Schulze
 767 M, Bamia C, Chloptsios Y, Veglia F, Panico S, Bueno-de-Mesquita H, Ocké M, Brustad
 768 M, Lund E, González C, Barcos A, Berglund G, Winkvist A, Mulligan A, Appleby P,
 769 Overvad K, Tjønneland A, Clavel-Chapelon F, Kesse E, Ferrari P, Van Staveren W,
 770 Riboli E (2002) Diversity of dietary patterns observed in the European Prospective
 771 Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* 5:1311 . doi:
 772 10.1079/PHN2002407
- 773 37. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH,
 774 Speizer FE (1985) Reproducibility and validity of a semiquantitative food frequency
 775 questionnaire. *Am J Epidemiol* 122:51–65 . doi: 10.1093/acprof
- 776 38. Vioque J, Weinbrenner T, Asensio L, Castelló A, Young IS, Fletcher A (2007) Plasma
 777 concentrations of carotenoids and vitamin C are better correlated with dietary intake in
 778 normal weight than overweight and obese elderly subjects. *Br J Nutr* 97:977 . doi:
 779 10.1017/S0007114507659017
- 780 39. Vioque J, Navarrete-Muñoz E-M, Gimenez-Monzó D, García-de-la-Hera M, Granado
 781 F, Young IS, Ramón R, Ballester F, Murcia M, Rebagliato M, Iñiguez C (2013)
 782 Reproducibility and validity of a food frequency questionnaire among pregnant women
 783 in a Mediterranean area. *Nutr J* 12:26 . doi: 10.1186/1475-2891-12-26
- 784 40. Bowen E, Heron J, Waylen A, Wolke D (2005) Domestic violence risk during and after
 785 pregnancy: findings from a British longitudinal study. *BJOG An Int J Obstet Gynaecol*
 786 112:1083–1089 . doi: 10.1111/j.1471-0528.2005.00653.x
- 787 41. Winsper C, Zanarini M, Wolke D (2012) Prospective study of family adversity and
 788 maladaptive parenting in childhood and borderline personality disorder symptoms in a
 789 non-clinical population at 11 years. *Psychol Med* 42:2405–2420 . doi:
 790 10.1017/S0033291712000542

- 791 42. Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL (2005) Urinary
792 Creatinine Concentrations in the U.S. Population: Implications for Urinary Biologic
793 Monitoring Measurements. *Environ Health Perspect* 113:192–200 . doi:
794 10.1289/ehp.7337
- 795 43. Baba Y, Furuta I, Zhai T, Ohkuchi A, Yamada T, Takahashi K, Matsubara S, Minakami
796 H (2017) Effect of urine creatinine level during pregnancy on dipstick test. *J Obstet*
797 *Gynaecol Res* 43:967–973 . doi: 10.1111/jog.13327
- 798 44. Sterne JAC, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, Wood AM,
799 Carpenter JR (2009) Multiple imputation for missing data in epidemiological and
800 clinical research: potential and pitfalls. *BMJ* 338:b2393 . doi: 10.1136/BMJ.B2393
- 801 45. Crawley H, Mills A, Patel S, Great Britain. Ministry of Agriculture F and F (1993) Food
802 portion sizes. H.M.S.O, London
- 803 46. World Health Organisation (2007) Iodine deficiency in Europe : a continuing public
804 health problem. <http://apps.who.int/iris/handle/10665/43398>. Accessed 1 Oct 2017
- 805 47. Phillips DI (1997) Iodine, milk, and the elimination of endemic goitre in Britain: the
806 story of an accidental public health triumph. *J Epidemiol Community Health* 51:391–3
- 807 48. Bath SC, Button S, Rayman MP (2014) Availability of iodised table salt in the UK – is
808 it likely to influence population iodine intake? *Public Health Nutr* 17:450–454 . doi:
809 10.1017/S1368980012005496
- 810 49. Glinioer D (1997) The Regulation of Thyroid Function in Pregnancy: Pathways of
811 Endocrine Adaptation from Physiology to Pathology. *Endocr Rev* 18:404–433 . doi:
812 10.1210/edrv.18.3.0300
- 813 50. Cheung KL, Lafayette RA (2013) Renal physiology of pregnancy. *Adv Chronic Kidney*
814 *Dis* 20:209–14 . doi: 10.1053/j.ackd.2013.01.012
- 815 51. Fuse Y, Shishiba Y, Irie M (2013) Gestational changes of thyroid function and urinary

- 816 iodine in thyroid antibody-negative Japanese women. *Endocr J* 60:1095–1106 . doi:
817 10.1507/endocrj.EJ13-0184
- 818 52. Bath SC, Furnidge-Owen VL, Redman CW, Rayman MP (2015) Gestational changes
819 in iodine status in a cohort study of pregnant women from the United Kingdom: season
820 as an effect modifier. *Am J Clin Nutr* 101:1180–7 . doi: 10.3945/ajcn.114.105536
- 821 53. Li C, Peng S, Zhang X, Xie X, Wang D, Mao J, Teng X, Shan Z, Teng W (2016) The
822 urine iodine to creatinine as an optimal index of iodine during pregnancy in an iodine
823 adequate area in China. *J Clin Endocrinol Metab* 101:1290–1298 . doi: 10.1210/jc.2015-
824 3519
- 825 54. Brander L, Als C, Buess H, Haldimann F, Harder M, Hänggi W, Herrmann U, Lauber
826 K, Niederer U, Zürcher T, Bürgi U, Gerber H (2003) Urinary iodine concentration
827 during pregnancy in an area of unstable dietary iodine intake in Switzerland. *J*
828 *Endocrinol Invest* 26:389–396 . doi: 10.1007/BF03345192
- 829 55. Stilwell G, Reynolds PJ, Parameswaran V, Blizzard L, Greenaway TM, Burgess JR
830 (2008) The Influence of Gestational Stage on Urinary Iodine Excretion in Pregnancy. *J*
831 *Clin Endocrinol Metab* 93:1737–1742 . doi: 10.1210/jc.2007-1715
- 832 56. Gerchman F, Tong J, Utzschneider KM, Zraika S, Udayasankar J, McNeely MJ, Carr
833 DB, Leonetti DL, Young BA, de Boer IH, Boyko EJ, Fujimoto WY, Kahn SE (2009)
834 Body Mass Index Is Associated with Increased Creatinine Clearance by a Mechanism
835 Independent of Body Fat Distribution. *J Clin Endocrinol Metab* 94:3781–3788 . doi:
836 10.1210/jc.2008-2508
- 837 57. O’Brien KM, Upson K, Cook NR, Weinberg CR (2016) Environmental Chemicals in
838 Urine and Blood: Improving Methods for Creatinine and Lipid Adjustment. *Environ*
839 *Health Perspect* 124:220–227
- 840 58. James GD, Sealey JE, Alderman M, Ljungman S, Mueller FB, Pecker MS, Laragh JH

- (1988) A Longitudinal Study of Urinary Creatinine and Creatinine Clearance in Normal Subjects: Race, Sex, and Age Differences. *Am J Hypertens* 1:124–131 . doi: 10.1093/ajh/1.2.124
59. Brantsæter AL, Haugen M, Julshamn K, Alexander J, Meltzer HM (2009) Evaluation of urinary iodine excretion as a biomarker for intake of milk and dairy products in pregnant women in the Norwegian Mother and Child Cohort Study (MoBa). *Eur J Clin Nutr* 63:347–354 . doi: 10.1038/sj.ejcn.1602952
60. Brantsæter AL, Abel MH, Haugen M, Meltzer HM (2013) Risk of suboptimal iodine intake in pregnant norwegian women. *Nutrients* 5:424–440 . doi: 10.3390/nu5020424
61. Gunnarsdottir I, Gustavsdottir AG, Steingrimsdottir L, Maage A, Johannesson AJ, Thorsdottir I (2013) Iodine status of pregnant women in a population changing from high to lower fish and milk consumption. *Public Health Nutr* 16:325–329 . doi: 10.1017/S1368980012001358
62. Mian C, Vitaliano P, Pozza D, Barollo S, Pitton M, Callegari G, Di Gianantonio E, Casaro A, Acamulli DN, Busnardo B, Mantero F, Girelli ME (2009) Iodine status in pregnancy: Role of dietary habits and geographical origin. *Clin Endocrinol (Oxf)* 70:776–780 . doi: 10.1111/j.1365-2265.2008.03416.x
63. Torres MT, Francés L, Vila L, Manresa JM, Falguera G, Prieto G, Casamitjana R, Toran P (2017) Iodine nutritional status of women in their first trimester of pregnancy in Catalonia. *BMC Pregnancy Childbirth* 17:249 . doi: 10.1186/s12884-017-1423-4
64. Bath SC, Walter A, Taylor A, Wright J, Rayman MP (2014) Iodine deficiency in pregnant women living in the South East of the UK: the influence of diet and nutritional supplements on iodine status. *Br J Nutr* 111:1622–31 . doi: 10.1017/S0007114513004030
65. Blumenthal N, Byth K, Eastman CJ (2012) Iodine Intake and Thyroid Function in

- Pregnant Women in a Private Clinical Practice in Northwestern Sydney before
Mandatory Fortification of Bread with Iodised Salt. *J Thyroid Res* 2012:1–6 . doi:
10.1155/2012/798963
66. Soriguer F, Gutierrez-Repiso C, Gonzalez-Romero S, Oliveira G, Garriga MJ, Velasco
I, Santiago P, de Escobar GM, Garcia-Fuentes E, Iodine Deficiency Disorders Group of
Spanish Society of Endocrinology and Nutrition (2011) Iodine concentration in cow's
milk and its relation with urinary iodine concentrations in the population. *Clin Nutr*
30:44–48 . doi: 10.1016/j.clnu.2010.07.001
67. National Institute for Public Health and the Environment (RIVM) (2016) Dutch Food
Composition Database (NEVO) online version 2016/5.0. In: RIVM, Bilthoven.
<http://nevo-online.rivm.nl/>. Accessed 19 Jan 2018
68. Lee SM, Lewis J, Buss DH, Holcombe GD, Lawrance PR (1994) Iodine in British foods
and diets. *Br J Nutr* 72:435 . doi: 10.1079/BJN19940045
69. Stevenson MC, Drake C, Givens DI (2018) Further studies on the iodine concentration
of conventional, organic and UHT semi-skimmed milk at retail in the UK. *Food Chem*
239:551–555 . doi: 10.1016/J.FOODCHEM.2017.06.135
70. Dahl L, Johansson L, Julshamn K, Meltzer HM (2004) The iodine content of Norwegian
foods and diets. *Public Health Nutr* 7:569–576 . doi: 10.1079/PHN2003554
71. Verkaik-Kloosterman J, Buurma-Rethans E, Dekkers A (2012) The iodine intake of
children and adults in the Netherlands: Results of the Dutch National Food
Consumption Survey 2007-2010 - RIVM.
http://www.rivm.nl/en/Documents_and_publications/Scientific/Reports/2012/april/The_iodine_intake_of_children_and_adults_in_the_Netherlands_Results_of_the_Dutch_National_Food_Consumption_Survey_2007_2010. Accessed 1 Oct 2017
72. Cross AJ, Major JM, Sinha R (2011) Urinary biomarkers of meat consumption. *Cancer*

- 891 Epidemiol biomarkers Prev 20:1107–11 . doi: 10.1158/1055-9965.EPI-11-0048
- 892 73. Lykken GI, Jacob RA, Munoz JM, Sandstead HH (1980) A mathematical model of
893 creatine metabolism in normal males--comparison between theory and experiment. Am
894 J Clin Nutr 33:2674–85
- 895 74. Shim J-S, Oh K, Kim HC (2014) Dietary assessment methods in epidemiologic studies.
896 Epidemiol Health 36:e2014009 . doi: 10.4178/epih/e2014009
- 897 75. Knudsen N, Christiansen E, Brandt-Christensen M, Nygaard B, Perrild H (2000) Age-
898 and sex-adjusted iodineacreatinine ratio. A new standard in epidemiological surveys?
899 Evaluation of three different estimates of iodine excretion based on casual urine samples
900 and comparison to 24 h values. Eur J Clin Nutr 54:361–363
- 901 76. Willett W (2012) Nutritional epidemiology, Third edit. Oxford University Press,
902 Oxford; New York
- 903

904 **Table 1** Descriptive statistics ^a of the study population by cohort

Sample characteristics	ALSPAC (n=2852)	Generation R (n=2254)	INMA (n=1460)
Maternal factors			
Age ^{b,c} (years), mean ($\pm$ SD)	28.7 ($\pm$ 4.5)	29.9 ($\pm$ 5.0)	31.4 ($\pm$ 4.1)
Pre-pregnancy BMI (kg/m ²), median (IQRs) ^c	22.3 (20.5 - 24.6)	23.5 (21.5 - 26.4)	22.5 (20.8 - 25.0)
Ethnicity ^d , n (%)			
Reference group	2800 (98.2)	1095 (48.6)	1340 (91.8)
Non-white	52 (1.8)	N/A	N/A
Non-Dutch	N/A	1159 (51.4) ^e	N/A
Non-Spanish	N/A	N/A	120 (8.2)
Parity ^c , n (%)			
0	1354 (47.5)	1279 (56.7)	806 (55.2)
1	965 (33.8)	665 (29.5)	544 (37.3)
≥ 2	533 (18.7)	310 (13.8)	110 (7.5)
Smoking status, n (%)			
Never smoked	2169 (76.1)	1671 (74.2)	1020 (69.9)
Stopped smoking	333 (11.7)	211 (9.3)	187 (12.8)
Continued smoking	350 (12.2)	372 (16.5)	253 (17.3)
Alcohol consumption, n (%)			
No	1350 (47.4)	1458 (64.7)	1330 (91.1)
Yes	1502 (52.6)	796 (35.3)	130 (8.9)
Markers of socio-economic status			
Education level, n (%)			
Low	576 (20.2)	247 (11.0)	337 (23.1)
Medium	1780 (62.4)	995 (44.1)	581 (39.8)
High	496 (17.4)	1012 (44.9)	542 (37.1)
Net household income (€ per month), n (%)			
Low < €1200	N/A	492 (21.8)	N/A
Medium €1200-2200	N/A	597 (26.5)	N/A
High > €2200	N/A	1165 (51.7)	N/A
Home ownership, n (%)			
Owned/mortgaged	2425 (85.0)	N/A	N/A
Private/other rented	236 (8.3)	N/A	N/A
Council rented	191 (6.7)	N/A	N/A
Crowding index, n (%)			
≤ 1 person per room	2747 (96.3)	N/A	N/A
+ 1 person per room	105 (3.7)	N/A	N/A
Family adversity index, n (%)			
None 0	1395 (48.9)	N/A	N/A
Mild 1-2	1124 (39.4)	N/A	N/A
Severe > 3	333 (11.7)	N/A	N/A

Table 1 Descriptive statistics ^a of the study population by cohort (*continued*)

Sample characteristics	ALSPAC (n=2852)	Generation R (n=2254)	INMA (n=1460)
Markers of socio-economic status			
Life event score, median (IQRs)	3.0 (2.0 - 5.0)	N/A	N/A
Marital status, n (%)			
Never-married	355 (12.5)	1136 (50.4)	N/A
Married	2357 (82.6)	1118 (49.6)	N/A
Other ^f	140 (4.9)	N/A	N/A
Living with a partner, n (%)			
Yes	2720 (95.4)	N/A	1445 (99.0)
No	132 (4.6)	N/A	15 (1.0)
Child factors			
Child's sex ^c , n (%)			
Male	1405 (49.3)	1147 (50.9)	737 (50.5)
Female	1447 (50.7)	1107 (49.1)	723 (49.5)

^a Data presented as mean ($\pm$ SD) for continuous normally distributed variables, median (IQRs) for continuous non-normally distributed variables and n (%) for categorical variables.

^b Maternal age at urine sample collection, except in ALSPAC (age at last menstrual period).

^c Data were not imputed, due to no missing values for age (in ALSPAC, Generation R), pre-pregnancy BMI (INMA), parity (Generation R) and child's sex (ALSPAC). The rest of the data are shown after imputation of the missing values (see methods).

^d ALSPAC (Reference group=White); Generation R (Reference group=Dutch, Non-Dutch=Indonesian, Cape Verdian, Moroccan, Dutch Antilles, Surinamese, Turkish, Other non-western, Asian, or Other western, see Table 3); INMA (Reference group=Spanish, Non-Spanish=Latin American, European, or Others).

^e Non-Dutch group in Generation R presented in detail in Table 3.

^f ALSPAC (Other=widowed, divorced, or separated).

Abbreviations: BMI, body mass index; IQRs, interquartile ranges; N/A, data not available or not applicable; SD, standard deviation.

919 **Table 2** Urinary iodine status in early pregnancy (≤ 18 gestational weeks) expressed as UIC, UI/Creat and proportion of mothers with UI/Creat
 920 below 150 $\mu\text{g/g}$

	ALSPAC (n=2852)	Generation R (n=2254)	INMA (n=1460)
Gestational age at urine sampling, weeks ^a	11.0 (8.0 - 15.0)	13.1 (12.1 - 14.6)	13.0 (12.4 - 13.9)
Urinary iodine concentration (UIC), $\mu\text{g/L}$ ^a	95 (56 - 151)	165 (94 - 277)	130 (76 - 219)
Iodine-to-creatinine ratio (UI/Creat), $\mu\text{g/g}$ ^a	121 (81 - 193)	210 (140 - 303)	151 (96 - 255)
UI/Creat < 150 $\mu\text{g/g}$, n (%)	1792 (62.8)	650 (28.8)	723 (49.5)

921 ^a Data presented as median (25th - 75th percentiles).

922 **Abbreviations:** UI/Creat, urinary iodine-to-creatinine ratio; UIC, urinary iodine concentration.

23

Table 3 Determinants ^a of urinary iodine-to-creatinine ratio measured at ≤ 18 gestational weeks, statistically significant in at least one cohort

Determinants	ALSPAC (n=2852)						Generation R (n=2254)						INMA (n=1460)					
	Adjusted Model 1 ^b			Adjusted Model 2 ^c			Adjusted Model 1 ^b			Adjusted Model 2 ^c			Adjusted Model 1 ^b			Adjusted Model 2 ^c		
	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>
Gestational age at urine sampling, weeks	2852	0.051	<0.001	2710	0.052	<0.001	2254	0.007	0.300	1580	0.011	0.133	1460	0.004	0.755	1446	-0.009	0.513
Age ^d , years	2852	0.014	<0.001	2710	0.010	0.002	2254	0.018	<0.001	1580	0.015	<0.001	1460	0.020	<0.001	1446	0.014	0.008
Pre-pregnancy BMI, kg/m ²	2852	-0.013	<0.001	2710	-0.012	<0.001	2254	-0.011	<0.001	1580	-0.013	<0.001	1460	-0.013	0.005	1446	-0.010	0.033
Family adversity index																		
None 0	1395	Ref.		1334	Ref.		N/A	.	.	.	.	.	N/A	.	.	.	.	.
Mild 1-2	1124	-0.007	0.765	1069	-0.003	0.906	N/A	.	.	.	.	.	N/A	.	.	.	.	.
Severe > 3	333	-0.100	0.016	307	-0.086	0.046	N/A	.	.	.	.	.	N/A	.	.	.	.	.
Marital status																		
Never-married	355	Ref.		320	Ref.		1136	Ref.		825	Ref.		N/A	.	.	.	.	.
Married	2357	0.095	0.015	2269	0.119	0.003	1118	0.030	0.285	755	0.044	0.168	N/A	.	.	.	.	.
Other ^e	140	0.028	0.666	121	0.051	0.447	N/A	.	.	.	.	.	N/A	.	.	.	.	.
Ethnicity ^f																		
Reference group	2800	Ref.		2674	Ref.		1095	Ref.		930	Ref.		1340	Ref.		1328	Ref.	
Non-white	52	-0.018	0.851	36	0.043	0.676	N/A	.	.	.	.	.	N/A	.	.	.	.	.
Non-Dutch:	N/A	.	.	.	.	.							N/A	.	.	.	.	.
- Indonesian	N/A	.	.	.	.	.	72	-0.047	0.494	58	-0.081	0.286	N/A	.	.	.	.	.
- Cape Verdian	N/A	.	.	.	.	.	87	-0.013	0.848	41	0.065	0.502	N/A	.	.	.	.	.
- Moroccan	N/A	.	.	.	.	.	165	0.255	<0.001	72	0.182	0.028	N/A	.	.	.	.	.
- Dutch Antilles	N/A	.	.	.	.	.	65	-0.174	0.025	37	-0.225	0.031	N/A	.	.	.	.	.
- Surinamese	N/A	.	.	.	.	.	196	-0.115	0.017	111	-0.160	0.014	N/A	.	.	.	.	.
- Turkish	N/A	.	.	.	.	.	226	0.391	<0.001	103	0.360	<0.001	N/A	.	.	.	.	.
- Other, non-western	N/A	.	.	.	.	.	107	0.157	0.011	53	0.125	0.137	N/A	.	.	.	.	.
- Asian	N/A	.	.	.	.	.	32	0.091	0.383	19	0.028	0.835	N/A	.	.	.	.	.
- Other, western	N/A	.	.	.	.	.	209	0.052	0.228	156	0.062	0.213	N/A	.	.	.	.	.
Non-Spanish	N/A	.	.	.	.	.	N/A	.	.	.	.	.	120	-0.016	0.810	118	-0.003	0.961

Table 3 Determinants ^a of urinary iodine-to-creatinine ratio measured at ≤ 18 gestational weeks, statistically significant in at least one cohort (*continued*)

Determinants	ALSPAC (n=2852)						Generation R (n=2254)						INMA (n=1460)					
	Adjusted Model 1 ^b			Adjusted Model 2 ^c			Adjusted Model 1 ^b			Adjusted Model 2 ^c			Adjusted Model 1 ^b			Adjusted Model 2 ^c		
	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>	<i>n</i>	<i>B</i>	<i>P</i>
Smoking status																		
Never smoked	2169	Ref.		2077	Ref.		1671	Ref.		1202	Ref.		1020	Ref.		1011	Ref.	
Stopped smoking	333	-0.003	0.943	312	-0.010	0.797	211	-0.093	0.035	147	-0.121	0.020	187	-0.060	0.296	186	-0.060	0.287
Continued smoking	350	0.018	0.620	321	0.060	0.116	372	-0.008	0.831	231	0.049	0.296	253	-0.001	0.990	249	0.020	0.707

^a Effect estimates (*B*=unstandardised regression coefficient) and *P*-values from multiple linear regression models performed for each cohort with (natural) log-transformed iodine-to-creatinine ratio (UI/Creat) as the dependent variable and maternal characteristics and dietary intakes as independent variables (for full models, see footnotes b and c). Reported *B* coefficients represent the change in the mean (natural) log of UI/Creat per unit increase in the continuous independent variables and for each category compared to the reference for the categorical independent variables.

^b Adjusted Model 1 (adjusted for maternal and pregnancy characteristics); ALSPAC ($R^2=0.123$, $P < 0.0001$): gestational age (weeks), age (years), pre-pregnancy BMI (kg/m^2), ethnicity, parity, smoking status, alcohol consumption, education, home ownership, crowding index, family adversity index, life event score, marital status and child's sex; Generation R ($R^2=0.086$, $P < 0.0001$): gestational age (weeks), age (years), pre-pregnancy BMI (kg/m^2), ethnicity, parity, smoking status, alcohol consumption, education, net household income, marital status and child's sex; INMA ($R^2=0.021$, $P < 0.0001$): gestational age (weeks), age (years), pre-pregnancy BMI (kg/m^2), ethnicity, parity, smoking status, alcohol consumption, education, living with a partner and child's sex.

^c Adjusted Model 2 (adjusted for maternal and pregnancy characteristics + dietary intakes); ALSPAC ($R^2=0.147$, $P < 0.0001$): Model 1 + energy intake (kcal/day) + intake of vegetables (g/day), fruit (g/day), nuts and seeds (g/day), potatoes (g/day), legumes (g/day), cereals and cereal products (g/day), cakes, confectionery and added sugar (g/day), added fats (g/day), milk and dairy products (g/day), meat and meat products (g/day), eggs (g/day), fish and shellfish (g/day), processed and fried foods (g/day), non-alcoholic beverages (g/day), miscellaneous (g/day); Generation R ($R^2=0.091$, $P < 0.0001$): Model 1 + energy intake (kcal/day) + intake of vegetables (g/day), fruit (g/day), nuts and seeds (g/day), potatoes (g/day), legumes (g/day), cereals and cereal products (g/day), cakes, confectionery and added sugar (g/day), added fats (g/day), milk and dairy products (g/day), meat and meat products (g/day), eggs (g/day), fish and shellfish (g/day), condiments and seasoning (g/day), processed and fried foods (g/day), non-alcoholic beverages (g/day), alcoholic beverages (g/day), miscellaneous (g/day); INMA ($R^2=0.060$, $P < 0.0001$): Model 1 + energy intake (kcal/day) + intake of vegetables (g/day), fruit (g/day), nuts and seeds (g/day), potatoes (g/day), legumes (g/day), cereals and cereal products (g/day), cakes, confectionery and added sugar (g/day), added fats (g/day), milk and dairy products (g/day), meat and meat products (g/day), eggs (g/day), fish and shellfish (g/day), condiments and seasoning (e.g., salt) (g/day), processed and fried foods (g/day), non-alcoholic beverages (g/day), alcoholic beverages (g/day).

^d Maternal age at urine sample collection, except in ALSPAC (age at last menstrual period).

^e ALSPAC (Other=widowed, divorced, or separated).

^f ALSPAC (Reference group=White); Generation R (Reference group=Dutch, Non-Dutch=Indonesian, Cape Verdian, Moroccan, Dutch Antilles, Surinamese, Turkish, Other non-western, Asian, or Other western); INMA (Reference group=Spanish, Non-Spanish=Latin American, European, or Others).

47 **Abbreviations:** BMI, body mass index; N/A, data not available or not applicable; Ref, reference category; UI/Creat, urinary iodine-to-creatinine ratio.

948 **Table 4** Multivariable associations of food group intakes estimated from FFQ (per portion size, g) ^a with urinary iodine-to-creatinine ratio
 949 measured at ≤ 18 gestational weeks, statistically significant in at least one cohort

Food group	Standard portion (g) ^a	Description ^a	ALSPAC (n=2710)		Generation R (n=1580)		INMA (n=1446)	
			<i>B</i> (95% CI) ^b	<i>P</i> ^c	<i>B</i> (95% CI) ^b	<i>P</i> ^d	<i>B</i> (95% CI) ^b	<i>P</i> ^e
Fruit	80 g	<i>a portion '5-a-day'</i>	2.22 (0.47, 4.03)	0.012	0.90 (-0.90, 2.75)	0.328	-3.51 (-7.16, 0.20)	0.064
Nuts and seeds	30 g	<i>a handful</i>	5.08 (-0.63, 11.53)	0.083	12.92 (4.83, 21.73)	0.001	18.63 (-10.61, 52.17)	0.223
Cereals and cereal products	36 g	<i>a medium slice of bread</i>	0.51 (-0.33, 1.36)	0.238	4.25 (2.23, 6.31)	<0.001	0.07 (-5.12, 5.40)	0.978
Cakes, confectionery and added sugar	50 g	<i>a chocolate bar</i>	2.39 (0.25, 4.63)	0.028	2.64 (-1.13, 6.57)	0.172	-9.19 (-22.95, 5.59)	0.217
Added fats	10 g	<i>spread on a slice of bread</i>	1.12 (-0.14, 2.41)	0.082	2.70 (0.39, 5.08)	0.022	-4.19 (-11.03, 2.89)	0.243
Milk and dairy products	200 g	<i>a glass of milk</i>	7.77 (5.65, 9.99)	<0.001	4.74 (2.01, 7.56)	0.001	14.07 (5.07, 23.45)	0.002
Meat and meat products	130 g	<i>a medium portion chicken breast</i>	1.65 (-2.96, 6.78)	0.497	3.75 (-5.35, 13.87)	0.433	-41.48 (-63.44, -16.20)	0.002
Eggs	50 g	<i>an average egg</i>	2.64 (-0.60, 6.12)	0.112	28.10 (12.27, 46.47)	<0.001	39.69 (-6.67, 96.68)	0.099
Fish and shellfish	120 g	<i>a medium cod fillet</i>	6.18 (1.07, 11.87)	0.016	2.42 (-19.04, 30.69)	0.845	34.08 (3.28, 69.37)	0.029
Condiments and seasoning (e.g., salt) ^f	5 g	<i>a level teaspoon (WHO) ^g</i>	N/A	.	1.34 (-1.50, 4.29)	0.359	214.41 (56.88, 465.35)	0.003

950 ^a Portion sizes are based on Food Standard Agency (1998) Food Portion Sizes (3ed.) London: TSO.

951 ^b Effect estimates (*B*=unstandardised regression coefficient) represent the actual change in the geometric mean of UI/Creat ($\mu\text{g/g}$) associated with a portion size
 952 increase in intake of a food group. *B* coefficients and their 95% CIs are calculated by back-transformation from logarithmic scale (see methods). Values are
 953 adjusted for dietary intake of other food groups, energy intake and other potential confounders (for full models, see Table 3, footnote c). When calculating the
 954 *B* coefficients all categorical covariates were set to their reference group and the continuous covariates gestational age, maternal age, pre-pregnancy BMI and
 955 energy intake were centred to their means.

956 ^c *P*-value adjusted for energy (kcal/day) + gestational age (weeks), age (years), pre-pregnancy BMI (kg/m²), ethnicity, parity, smoking status, alcohol
957 consumption, education, home ownership, crowding index, family adversity index, life event score, marital status and child's sex ($R^2=0.147$, $P < 0.0001$).

958 ^d *P*-values adjusted for energy (kcal/day) + gestational age (weeks), age (years), pre-pregnancy BMI (kg/m²), ethnicity, parity, smoking status, alcohol
959 consumption, education, net household income, marital status and child's sex ($R^2=0.091$, $P < 0.0001$).

960 ^e *P*-values adjusted for energy (kcal/day) + gestational age (weeks), age (years), pre-pregnancy BMI (kg/m²), ethnicity, parity, smoking status, alcohol
961 consumption, education, living with a partner and child's sex ($R^2=0.060$, $P < 0.0001$).

962 ^f In INMA this food group also includes iodised salt. Separate information about the consumption of table salt and use of iodine-fortified table salt was collected
963 only in the INMA cohort.

964 ^g Maximum daily salt intake, as recommended by WHO (5 g/day salt, or 2 g/day Sodium).

965 **Abbreviations:** BMI, body mass index; 95% CI, confidence interval; FFQ, food frequency questionnaire; N/A, data not available; UI/Creat, urinary iodine-to-
966 creatinine; WHO, World Health Organisation.

Fig. 1 Flow chart of the study population selection

^a Urine samples with UIC > 500 µg/L or UI/Creat > 700 µg/g were excluded only from the ALSPAC cohort, as there was a concern about contamination by the use of iodine-containing test strips (see

971 methods). There was no such concern in Generation R and INMA, therefore the exclusion criteria was
972 not applicable to urine samples from these two cohorts.

973 **Abbreviations:** FFQ, food frequency questionnaire; IVF, in-vitro fertilisation; N/A, not applicable;
974 UI/Creat, urinary iodine-to-creatinine ratio; UIC, urinary iodine concentration.