ESTs and SNPs: What large scale sequencing projects can tell us about ADME

NICK PLANT

School of Biomedical and Molecular Sciences, University of Surrey, Guildford,
Surrey, GU2 7XH, UK

1. To date over 800 complete genomes have been sequenced, with many more partially complete. Coupled with the large amount of mRNA transcript sequence data being produced from expression studies there is now a daunting amount of information available to the research scientist. This review will aim to examine how this information may be best used, focussing on examples from sequences encoding ADME-related proteins in particular.
2. Through the use of phylogenetic, splice variant and SNP analysis I shall examine how insights into not only species-specific responses to drug exposure may be gained, but also how best we utilize this information to predict both individual human responses and the impact of population variance in response.

Keywords: Phylogenetic analysis, Expressed sequence tag, Single nucleotide polymorphism, species difference

Running Title: ESTs and SNPs

Correspondence: Nick Plant, School of Biomedical and Molecular Sciences,
University of Surrey, Guildford, Surrey, GU2 7XH, UK. Fax: +44 (0)1483 686401.
Email: N.Plant@Surrey.ac.uk

General Introduction

The ultimate aim of biological sciences has always been to understand how the body works, to actually explain what makes humans tick. An extension of this is required by the ADME scientist; not only to understand how the body works but, more importantly, to be able to *predict* how it will respond to chemical exposure. In the last decade a step-change has occurred in the capacity of man to generate information about the biological subsystems of the body, and how these alter in response to chemical exposure. This increase in technological ability has resulted in a huge amount of data available in silico to the research scientist; the challenge now is how to make the most use of this information.

This review will aim to examine the emergent technologies available for large scale information gathering, as well as the databases where such information is stored. Whereas public data storage is a central part of open-access research for the scientist, perhaps of more import today is the development of tools with which to analyse this freely available data, and a selection of these tools will also be covered in this review.

Public domain storage of sequence information and the development of computational tools to examine such a large amount of data are central to our further understanding of systems biology. However, six years after the sequencing of the human genome how far have we actually come? What insights (if any) have we learnt through the mining of these large databases? This review will attempt to address this question, or at least some parts of it, by examining how in silico analysis has aided our understanding of human biology, and in particular ADME: First, how the sequencing of non-human genomes may increase our knowledge of how the human body functions. Second, the discovery of novel genes, which may represent novel therapeutic targets, or whose protein products may impact upon ADME processes.

Finally, how single nucleotide polymorphism (SNP) analysis may help researchers to understand/predict inter-individual variation in response to chemical exposure.

Looking in to the past: What can phylogenetic analysis tell us?

There are presently fully sequenced genomes from just over 800 different species (www.ncbi.nlm.nih.gov/), the majority of which are available through public domain repositories such as NCBI, EBI and GenomeNet (based in the USA, Europe and Japan respectively). If one also includes all currently active genome sequencing projects then this number increases to over 4500 (Table 1), with approximately one third of these being animal species. However, a cursory glance at these will tell you that of these 800 only a relatively small number represent species directly applicable to the AMDET process (Table 2).

[TABLE 1+2 AROUND HERE]

The impact of the human genome sequence has been obvious and immediate; with it we are (theoretically) able to identify every human gene and extrapolate this information to functional biology. Indeed, some startling discoveries have been made such as the relatively low number of genes (now estimated at 20-25,000; Consortium 2004), and the overall high level of identity to supposedly very different organisms such as the mouse (Lander, et al. 2001, Waterston, et al. 2002). What has not been achieved (yet) however is a full understanding of the human biological system and all its intricate workings, and for that the other 800+ sequenced genomes have a significant role to play. From a homocentric point of view one can imagine that humans are at the (current) end of an evolutionary process that began some 3.5 billion

years ago (Kumar and Hedges 1998). This evolutionary process, culminating in *Homo sapiens* has revolved around the emergence of traits that allow humans to survive in our particular environment. Thus, we cannot say that we are the pinnacle of evolution, merely the end of this particular branch; other organisms have evolved to suit their particular environments and are hence the ‘top’ of their evolutionary tree. Traits that are beneficial for survival are kept, whereas those that are non-essential to life may be lost or altered through time. Following this argument in reverse it can be argued that traits (or genes) maintained in a large number of organisms are important for life in general, whereas those that exist in only a single species are important for that organism to exist in its particular environment. Through the study of the genomes of our evolutionary ancestors it may thus be possible to gain insights into the functioning of the human body. In addition, the less complex biological systems exhibited by many of these organisms means that gene functions are much better described, and hence it is possible to gain insights into human systems biology through the study of these ‘simpler’ organisms.

Genes whose protein products are involved in ADME lend themselves well to this type of analysis for a number of reasons. First, the need to respond to external chemicals has existed for a long time, thus meaning that these genes are present through millions of years of phylogenetic lineage. Second, the requirement to respond to a large number of different chemical entities has resulted in relatively large gene families: Such large numbers per species are useful in phylogenetic analysis for examining differences between evolutionarily close species (e.g. rats and humans). Third, the different environments that organisms are exposed to results in different pressures upon them, favouring the emergence of traits to cope with these specific pressures. By studying how different organisms have evolved to cope with their

particular environments we may gain important insights into how humans have evolved to cope with ours.

Perhaps the two best examples of where phylogenetic analysis has yielded insights into biological function for ADME-related genes are the cytochrome P450s and their controlling ligand-activated transcription factors, the nuclear receptors.

Cytochrome P450 phylogenetic analysis

Cytochrome P450 (CYP) enzymes are amongst the most ancient proteins known, with gene present in all species examined from archaeobacteria to humans (Nelson, et al. 1996). The reason for the prevalence is simple; CYPs are a fundamental enzyme family in the metabolism (both anabolic and catabolic) of the endogenous chemicals required for life (Nebert and Gonzalez 1987). Their ability to metabolise xenobiotics is to a great extent an acquired secondary characteristic, albeit one central to ADME. Indeed, the majority of human CYPs are still concerned with endogenous metabolism, with only CYP families 1-3 (representing 18 out of 57 human CYPs) primarily concerned with the metabolism of xenobiotics (Werck-Reichhart and Feyereisen 2000). The number of CYP genes underwent a rapid increase following the development of organisms with a gastro-intestinal tract, with increased chemical intake through the diet becoming a significant driver in the need to evolve new proteins capable of metabolising these chemicals (Nebert and Gonzalez 1987). An interesting example of this is how rodents have evolved over 100 individual CYP genes, far greater than the 57 found in humans. A cogent argument is that omnivorous rodents take in many more chemicals through their diet due to their scavenging lifestyle, and hence require more CYPs to protect them from potential toxins.

The Committee for Standardised Cytochrome P450 nomenclature has accepted the unenviable task of bringing all CYP sequence-related research under one roof, and

more importantly under one naming system. Sequences are named according to their identity, being assigned to families and sub-families at the 70 % and 40 % identity levels respectively. Such a nomenclature relies upon the accuracy of gene identification, and by inference the accuracy of gene prediction software (Wang, et al. 2004). Unfortunately, Cytochrome P450 gene sequences do not lend themselves well to automated identification; CYP genes sequences are often highly similar, and often exist as chromosome clusters of related genes. Such clusters are easily misinterpreted by automated gene prediction programs, producing 'fusion-genes' from two CYP-gene sequences within the gene cluster (Nelson 2002, Nelson 2005). For example, in human the four CYP3A sub-family members CYP3A4, 5, 7 and 43 are located on chromosome 7q21 within a single cluster, demonstrating an average 82 % identity (Gibson, et al. 2002). CYP gene prediction is particularly difficult in plant genomes, where gene clusters can extend to 10 or more highly related gene sequences (Nelson, et al. 2004). To further complicate the situation some CYP genes have evolved remarkably short exons, often comprised of only three or four amino acids (Doddapaneni, et al. 2005). Such short exons are often excluded by automated gene prediction programs, obviously impacting upon any naming system relying upon sequence identity, as well as potentially altering the implied biologically functioning of the protein products. Thus, it can be seen that whereas there have been huge improvements in the gene-hunting software available, for many gene families such as the CYPs there is no alternative but painstaking individual annotation by scientists. More than anything else this justifies the existence of websites such as the Committee for Standardised Cytochrome P450 Nomenclature, preventing duplication of this painstaking work, or invalid investigations based upon the erroneous gene sequences produced via automated data mining.

The many genome sequencing projects underway now make it possible to examine CYP evolution among a wide range of organisms, including humans and rodents (96 million years divergence), humans and pufferfish (420 million years divergence; Nelson 2003) and humans and *Arabidopsis* (670 million years divergence; Nelson, et al. 2004). Through the study of CYPs in a monocot (*Oryza sativa*, Japanese rice) and dicot (*Arabidopsis*) plant it is apparent that plants have many more CYPs than humans, with these species having ~350 and ~250 CYP genes respectively. The reason for this large number of CYP genes is not clear, but Nelson and colleagues speculate that it may be as a result of the pressures exerted on early plants during colonization of land (Nelson, et al. 2004). Of the rice and *Arabidopsis* CYPs approximately two-thirds are shared, having emerged prior to the monocot-dicot divergence 200 million years ago; however, despite this large overlap in shared genes only approximately one fifth of the CYP subfamilies are conserved, with these presumably encoding those CYPs involved in reaction fundamental to survival.

Comparing CYPs from humans and the pufferfish (*Takifugu rubripes*) produces far fewer differences than observed between plants and humans (Nelson 2003). Despite over 420 million years of divergence pufferfish and humans share 17 out of the 18 CYP families, and within these families gene structures (e.g. intron-exon boundaries) are well conserved. Such results suggest that the defining characteristics of vertebrate CYPs have been in place for at least 420 million years, with the only major divergences seen in those subfamilies, such as CYP2C, which are primarily targeted towards the metabolism of xenobiotics (Nelson 2003). The unique family in humans compared to pufferfish is CYP39, and this in itself is interesting as the emergence of this sub-family provides an alternate pathway for the biosynthesis of bile acids: This is a demonstration of the ability of CYPs to produce safety-nets,

whereby important metabolic functions exhibit redundancy in an attempt to prevent toxicity due to disruption of cellular homeostasis.

Nuclear receptor phylogenetic analysis

If the CYP-family members are the engines of Phase I metabolism then the nuclear receptors can be envisaged as the drivers; they control the when, where and how much for expression of many of the *CYP* genes as well as a large number of other gene families involved in ADME. It is thus logical to apply the same phylogenetic analysis to this family of ligand-activated transcription factors as it was to their target genes; understanding the evolution and biological functioning of nuclear receptors may help researchers further understand the regulation/functioning of their target genes.

Following the complete sequencing of the first three mammalian genomes (human, rat and mouse) Zhang and colleagues undertook an examination of the nuclear receptor family in these three species, employing a different approach to identifying novel related genes (Zhang, et al. 2004). Using the Pfam database (<http://www.sanger.ac.uk/Software/Pfam/>; Bateman, et al. 2004) these latter authors identified six structural and functional protein domains specific for the members of the nuclear receptor family, and then identified these in the 62 previously annotated human, rat or mouse nuclear receptors. Next, the complete genomes of the three species were examined for these Pfam domains using GENEWISEDB (Birney and Durbin 2000). Finally, alignment to the known nuclear receptor family members was then used to complete gene characterization. Using this methodology an additional 82 complete nuclear receptor genes were identified, along with two decayed aberrant genes sequences and ten pseudogenes. Of the 144 complete genes, 47 were present in rats, 49 in mice and 48 in humans; in itself an interesting finding when one considers

the disparity in murine and human CYP number, archetypal nuclear receptor target genes. The ‘missing genes’ in rats are NR1D2 and NR2E3, but their absence may be explained by gaps in the rat genome sequence at the places where they would be predicted to occur based upon their location in the mouse genome.

Examination of the rate of nucleotide substitutions within the nuclear receptor family, resulting in either synonymous (K_s) or non-synonymous (K_a) amino acid substitutions within the translated protein, revealed little species-specific differences, suggesting that for most of the nuclear receptors ligands were unlikely to be species-specific. However, for PXR and CAR K_a/K_s values were 4.0 and 5.6 times greater than average respectively, suggesting positive selection for ligands; this is perhaps unsurprising considering the role of these two nuclear receptors in regulating the body’s ability to respond to external chemicals, exposure to which is often species-specific due to the different diet compositions of these species. Such information must also raise concerns for ADME-testing and species extrapolation; this data confirms the observation first made in CYPs, that the very proteins whose responses we wish to study/predict are those that evolve the most rapidly and are therefore the most problematic to predict for. Increased K_a/K_s rates were also observed for members of the NR0B and NR1H sub-families, although the biological causes/consequences of this have not yet been elucidated.

The examination of the nuclear receptor family has resulted in a number of important discoveries. First, rapid evolution of members of this family, in particular PXR and CAR, demonstrates a strong positive selection, probably driven by the need to respond to external stimuli. Indeed, the Chicken Xenobiotic Receptor (CXR) has diverged into the Constitutive Androstane Receptor (CAR, NR1I3) and Pregnane X Receptor (PXR, NR1I2) in the space of only 310 million years (Handschin, et al.

2000). As a result of this divergence, higher vertebrates, including primates, have gained a second receptor targeted toward the capture, interpretation and response to external stimuli, potentially an important factor in the success of these species in surviving on an omnivorous diet. Second, this evolution has resulted in the same sort of redundancy seen in the CYP enzymes, where one chemical may be a ligand for several receptors (e.g. CAR and PXR) providing a 'metabolic safety net' (Xie, et al. 2000), or where internal control mechanisms (e.g. auto-regulation of *PXR* expression by PXR protein) are present to regulate the biological action of these interactions, thus preventing/reducing negative impacts on cellular homeostasis (Aouabdi, et al. 2006).

Perhaps the most surprising conclusion from the phylogenetic analysis was the ancestry of the SHP and DAX1 genes. SHP and DAX1 are the two members of the NR0 family of nuclear receptors and are unique in the fact that they do not contain a DNA-binding domain (Bavner, et al. 2005). The lack of functional DNA-binding capacity within these two proteins was suggestive of ancient nuclear receptors, from which the DNA-binding, fully functional family members had evolved (Guo, et al. 1996). However, this more complete phylogenetic analysis clearly demonstrates that these two latter nuclear receptors evolved from the NR2 branch of the phylogenetic tree, rather than acting as the root for the entire tree. During, or closely after the emergence of these NR0 family members the DNA-binding domain coding sequence appear to have been lost, with rapid sequence evolution following. The biological consequence of nuclear receptors without DNA-binding capacity is now emerging, with both SHP and DAX1 being implicated in the regulation of other nuclear receptors via protein:protein interactions (Bavner, et al. 2005, Niakan and McCabe 2005). The formation of non-DNA binding heterodimers of SHP/DAX1 with other

nuclear receptors effectively acts as a sink, removing nuclear receptors from the pool available to interact with ligand and/or DNA, regulating their biological effect.

New gene discovery

Expressed Sequence Tag Analysis

The advent of high throughput sequence analysis has had two major benefits. As discussed in the previous section, the first of these is the production of complete genome sequences for a large number of organisms, allowing phylogenetic analysis of gene families, providing further insights into their biological roles. The second major benefit is a large increase in the number of expressed sequences available in databases. These sequences are often multiple repeat sequencing reads of known genes/transcripts, which can be used to identify sequence variation, as will be discussed in the next section. However, a large number of these sequences are unassigned to any particular gene/transcript and hence may represent ‘novel’ sequences; herein I will discuss how these potentially novel sequences can be further examined.

Entire sections of sequence repositories are now devoted to expressed sequence tags (ESTs); those sequences are derived from mRNA and which should therefore result in a protein product, but for which no protein product or biological function has been assigned. It is from this latter pool that the exciting possibility of novel gene identification emerges. Before classifying any EST as a product of a ‘novel gene’ it is important to note that the majority of ESTs merely represent *unassigned* transcripts and not *novel* transcripts. Due to technical considerations in transcript isolation/processing and the use of high throughput single-read sequencing, much of the transcript sequence deposited in databases is not full length and biased towards the

3' terminus. Given that the majority of eukaryotic genes produce transcripts with untranslated regions (UTRs) at either extremity, and that the average length of a 3' UTR is 826 bp for humans (Wren, et al. 2000) and http://innovation.swmed.edu/research/informatics/res_inf_pom.html), it is perhaps not surprising that the majority of ESTs contain mainly 3' UTR sequences. As UTRs are in general not as well conserved as the coding sequence within a transcript (Wren, et al. 2000) identification of any transcript from this sequence may be problematic as identification via orthologues may be inaccurate, particularly for transcripts from large gene families, such as those associated with ADME. Hence, ESTs often merely represent 3' sequences of transcripts that have not been married to their known genes, and this can be achieved with relative ease using simple database searching. However, a number of ESTs do not map to known genes, nor translate to proteins of known function; these represent transcripts from completely novel genes, and are of particular interest both in understanding the potential biological effects of chemicals, but also for the identification of potential novel therapeutic targets for the treatment of disease.

Target Gene Identification

An important step in understanding/predicting how the body responds to chemical exposure is the characterisation of what molecular events occur immediately following exposure. Such events may be classed as direct (e.g. binding of a nuclear receptor to its target gene) or indirect (e.g. expression of other genes as a result of the direct actions of the chemical), and it is important to separate these two if we are to fully understand the molecular responses to chemical stimulation. To understand the chronology of response, and hence the mode of action of chemicals it is important to

identify the primary gene targets, with these later being associated with (secondary) biological responses.

An example of such an approach is provided by Horie-Inoue and colleagues, who mined the human genome sequence to identify potential binding sites for nuclear receptors involved in hormone homeostasis/response (Horie-Inoue, et al. 2006). The androgen receptor, glucocorticoid receptor, progesterone receptor and mineralocorticoid receptor all bind to an IR3 response element comprised of two inverted AGAACA half-sites, separated by three nucleotides. Genome wide scanning for this sequence identified 565 exact matches for this element, and of these 26 were within 10 Kb of the known transcription start sites for annotated human genes. Chromosome immunoprecipitation (ChIP) was then used to identify which nuclear receptors (if any) bound to these sites, revealing 14/26 to be active. This data shows the potential and limitations of in silico screening; whereas such data mining identified 565 potential binding sites, only 5% were close to known genes and of these only one-half were shown to be bound by a nuclear receptor. Thus, while such techniques can be seen to generate a lot of potential *clues*, they do not provide the *answers* – for that wet-lab experiments are a necessity.

It should also be noted that this type of analysis concentrates on perfect matches to the consensus binding site, and will almost certainly miss some functional binding sites as nuclear receptors have been clearly demonstrated to bind to sequences divergent from their consensus binding site (Song, et al. 2004, Xie, et al. 2000). This is highlighted by the recent work of Steketee and colleagues who, rather than search for one perfect ‘consensus’ element within a large amount of DNA, searched for many near-perfect elements within a restricted region of DNA (Steketee, et al. 2004). Using this approach these latter authors identified a non-perfect HRE within the first

intron of the specifically androgen-regulated gene (SARG), and went on to show that this site was functional via ChIP.

Hence, it can be seen that a combination of the two previous methods is recommended to provide the most complete coverage for identifying putative transcription factor binding sites – an arduous task given the complexity of the human genome. A potential solution for this problem is the use of phylogenetic fingerprinting. Through the use of programs such as rVISTA (Loots and Ovcharenko 2004, Loots, et al. 2002) and PipMaker (Schwartz, et al. 2000) it is possible to scan genome sequences for areas of conservation across species (Pennacchio and Rubin 2003). Those regions that show highest conservation usually correspond to genes, with the next most highly conserved regions often representing regulatory elements. Using such an approach Ozyildirim and colleagues recently examined lacrimal gland transcriptomes to identify ‘lacrimal-specific’ transcripts, plus the regulatory elements putatively involved in their expression (Ozyildirim, et al. 2005). They were able to identify a number of ‘lacrimal-specific’ transcripts but showed that conservation of the expression and regulation of these specific transcripts between mouse and human was poor; questioning the relevance of currently used murine models of human dry eye syndromes.

Biological function identification

Another benefit from the large sequence repositories is that there is now a large amount of data available for transcript expression levels in different tissues and under different conditions. With such information it should be possible to identify differences between, for example, normal and diseased tissue (Mandel, et al. 2003, Mariadason, et al. 2002) or before and after exposure to a chemical (Natsoulis, et al. 2005, Rogler, et al. 2004). This would allow the identification of the expression

changes elicited by a stimulus, and through modelling of the pathways affected by these changes insights into the molecular mechanisms of response can be gained.

Massively parallel signature sequencing (MPSS) is a technique for generating large numbers of very short sequences very rapidly (Brenner, et al. 2000). These can then be identified through mining of sequence databases to identify longer sequences for which genes/protein products have been assigned, and the number of tags present for each signal provides a measure of the expression level. The throughput of MPSS means that all transcripts within a given cell (>200,000) can be sampled very rapidly, meaning that the complete cataloguing of gene expression within cell types/conditions is now possible (Jongeneel, et al. 2005). Chen and colleagues used such an approach to first identify 1,056 transcripts predominantly expressed in human testis, and then compared the expression of these genes in normal tissue and testis cell lines. Using this approach they were able to identify 20 cancer/testis antigen genes, whose expression is linked to the hypomethylation that occurs during cell transformation (Feinberg, et al. 2002), including a previously unidentified gene family. Such genes offer exciting potential therapeutic targets for the development of antibody-based anti-cancer agents.

SNPs and Splicing: Producing alternate transcripts

As previously discussed, the rapid increase in the size of sequence databases is due to two main driving forces, the sequencing of novel genes/transcripts/genomes from disparate species and the repeated sequencing of these targets. The repeat sequencing of already 'known' sequences is important for two functions: First, it confirms the accuracy of a sequencing project, analogous to the requirement to carry out a wet-lab experiment at least twice to confirm the validity/robustness of conclusions, and

second, it reveals any variation seen within that sequence that is not due to experimental/technical error but a true variation seen within the general population. This variation may be due to alternative splicing of identical pre-mRNA transcripts, or due to variations within the genome itself; both of these scenarios will be discussed in the following sections, along with their potential implications for ADME.

Splice variants of ADME-related genes

Following the discovery that the human genome contains only approximately 20,000 genes came the realisation that added complexity must be factored in to achieve the total number of observed proteins in humans, estimated at an order of magnitude higher. A large amount of this variation can be accounted for by the presence of post-translational modifications such as phosphorylation, which occur to most proteins. However, it is becoming increasingly clear that at least some of this increase in complexity is achieved through the use of alternative splicing, where different exons are spliced together from the nascent pre-mRNA to form the mature mRNA, translation of which produces a protein product with the inclusion/removal of different amino acid cassettes (Lareau, et al. 2004). Indeed, it has been estimated that more than half of all human genes are capable of producing alternative splice variants (Hanke, et al. 1999). Such splice variants may have a number of roles; for example the tumour necrosis factor receptor gene (TNFR) gives rise to at least ten different splice variants, which may affect factors such as sub-cellular localisation of the receptor (Xing, et al. 2003) or cellular predisposition to apoptosis (Screaton, et al. 1997). A number of specialist websites have now been developed to provide central repositories for alternate splice variants of genes, some of which are given in Table 3.

[TABLE 3 AROUND HERE]

Alternative splicing of ADME-related genes is a common phenomenon, with examples existing at the level of drug transporters, nuclear receptors and enzymes. This review will concentrate on splice variants within ADME-related enzymes as examples of both the prediction and biological consequences of alternative splicing. N-acetyltransferases are widely expressed enzymes important in the conjugation of amino and hydroxyl amino groups with acetate, and are responsible for the metabolism of a number of xenobiotics, including the activation of arylamine and heterocyclic amine procarcinogens (Gooderham, et al. 2001). In humans, two N-acetyltransferase enzymes are present, NAT1 and NAT2, and recent work by Butcher et al described the genomic organization of the former of these genes, and identifies a number of splice variants (Butcher, et al. 2005). The NAT1 gene is comprised of nine exons, but, unusually, the protein coding region of NAT1 is contained within just a single exon, with the remaining eight coding for the 5' untranslated region (UTR). Examination of the NCBI EST database identified these non-coding exons and suggested that at least three alternative promoters were involved in the production of nine separate splice variants from this gene. As the splice variants alter the non-coding region of the transcript they do not produce proteins with different activities but rather have different translation efficiencies. Butcher et al suggested that the differential expression of these splice variants throughout the body may go some way to explaining the different activity of NAT1 in different organs of the body.

An interesting form of alternative splicing seen in a number of ADME-related genes is the use of alternative promoters. Here, transcription may begin at a number of alternate first exons within a gene, presumably driven via alternative biological/xenobiotic stimuli. The resultant pre-mRNA will contain the requisite

number of conserved exons plus at least one alternate first exon(s): The mature mRNA is then generated through the splicing out of any extra first exons. Examples of ADME-related genes for which alternative promoters/first exons play an important role in their biological functioning are the UDP-glucuronosyltransferase (UGT1A) gene (Tukey and Strassburg 2000) and the N-acetyltransferase 1 gene (Butcher, et al. 2005). These two examples demonstrate the different biological affects that alternative splicing may cause: In the UGT1A gene, splice variants result in proteins with significantly different catalytic activities, whereas NAT1 splice variants result in different levels of protein expression. The UGT1A gene is of particular interest since the protein products generated from four common exons and one, variable, first exon are sufficiently different to be termed separate proteins, forming the members of the UGT1A-family (Tukey and Strassburg 2000). This is thus an example of how alternative splicing can considerably alter the functionality of a protein product, to the point where it can be considered a separate entity rather than just a variant.

Alternate splicing shows the remarkable flexibility of genomic DNA, in that the derivation of protein products from static nucleotide sequences can be manipulated to increase the complexity of the biological system. Not only is this a very elegant method for increasing the ability of a system to change in response to external situations, be they developmental stage of chemical exposure, but it also allows the maximal variation in protein products to be formed with the minimum changes in genomic DNA: Thus, alternative splicing represents a very elegant method for biological systems to respond to the World in which it lives.

DNA variation in ADME-related genes

Alternative splicing may be seen as a consistent form of variation (if such a thing exists!) in that it allows all members of a species to produce alternate transcripts, and therefore protein products from a single gene, under different biological conditions. In contrast, genetic polymorphisms are defined as variation in genome sequences within a population, with not all members of that species having the same sequence. These features may be combined, with a polymorphism creating splice variants through the addition/removal of splice donor/acceptor sites. For example, expression of CYP3A5 in humans is varied, with only approximately 25% of American Caucasians expressing a functional protein (Gibson, et al. 2002). This variation is, at least in part, controlled by a polymorphism that allows the introduction of mini-exons into the mature mRNA; as these mini-exons contain stop codons the resultant splice variant produces a truncated protein that is non-functional (Kuehl, et al. 2001).

Polymorphisms are a likely source for the well known variation in predisposition to disease, and as such much study has been undertaken on examining associations between genotype status and predisposition to disease. However, due to a large number of factors (population size/source screened, environmental considerations etc) many of these studies have failed to show convincing, or reproducible, associations. Perhaps the most convincing association to date between a DME and disease predisposition is CYP19 and hyperandrogenism. In comparison to the relatively sparse evidence for associations between DME status and disease predisposition, much evidence exists for the role of DME status and variation in response to therapeutic agents (Table 4 for examples), and this is hence an important research target, particularly given the increasingly global market for drugs. The ability to predict the efficacy, or potential adverse effects, caused by drugs when given to individuals from different genetic backgrounds would be of great benefit, perhaps

even heading towards the holy grail of personalised medicine (Vizirianakis 2004). However, before such a goal can be reached not only must the sources of genetic variation be identified, but their effect on ADME established.

[TABLE 4 AROUND HERE]

As can be seen from Table 4 even a cursory search of PubMed will reveal a host of SNPs associated with altered response to therapeutic agents, resulting in loss/reduced efficacy or an increased risk of adverse effects. However, such information is not particularly accessible, with the search term largely determining the information retrieved. To allow more efficient identification of SNPs and their biological function a number of specialized websites have been developed such as HGVBase (<http://hgvbase.cgb.ki.se/>; (Fredman, et al. 2002), dbSNP (<http://www.ncbi.nlm.nih.gov/projects/SNP/>; (Sherry, et al. 1999), SNP Consortium (<http://snp.cshl.org/>; (Thorisson and Stein 2003) and HapMap (<http://www.hapmap.org/>; (Consortium 2003), which are fully searchable curated databases of SNPs. Perhaps the most ambitious of these is HapMap, which aims to produce a freely available resource of the frequency and associations between in excess of one million variants from population deriving from Africa, Asia and Europe; the first phase of this project was recently completed, with 269 DNA samples from diverse ethnic backgrounds analyzed for these SNPs (Consortium 2005). In addition to these more specialized databases, general databases such as Gene and OMIM (both at <http://www.ncbi.nlm.nih.gov/>), Ensembl (<http://www.ensembl.org/>; (Birney, et al. 2004)) and GeneCards (<http://bioinformatics.weizmann.ac.il/cards/>) contain much useful information on a large number of genes, including

common/biologically important polymorphisms, with links to relevant literature; as such they should be the first stop for any researcher wishing to assess the potential impact of polymorphisms on their pet gene. In relation to ADME-related genes it is also worth noting that for CYPs the CYP alleles nomenclature website (<http://www.imm.ki.se/CYPalleles/>) includes details of polymorphisms present within the CYP family of genes, along with biological significance, prevalence and related literature, although this information is limited to human alleles.

Summary

The emergence of high throughput sequencing has resulted in a vast increase in the information available to researchers in their quest to understand human biology. Genome sequencing projects provide information on where we have come *from*, with phylogenetic analysis helping scientists to examine the way humans have evolved to cope with their biological needs. Large scale sequencing projects have also revealed the large numbers of splice variants used by biological organisms to further increase the complexity of their protein pool, allowing humans to respond to variations in their *present* environments. Finally, SNP analysis may give some hints as to where we are *going*, showing genetic variations that are fixed within the population, and their effects on an individual's disease predisposition or response to therapeutic agents.

References:

- Ando M, Hasegawa Y and Ando Y. 2005. Pharmacogenetics of irinotecan: a promoter polymorphism of UGT1A1 gene and severe adverse reactions to irinotecan. *Investigations on New Drugs*. 23:539-545.
- Aouabdi S, Gibson GG and Plant N. 2006. Transcriptional regulation of PXR: Identification of a PPRE within the proximal promoter responsible for fibrate-mediated transcriptional activation of PXR. *Drug Metabolism and Disposition*. 34:138-144.
- Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, Khanna A, Marshall M, Moxon S, Sonnhammer EL, Studholme DJ, Yeats C and Eddy SR. 2004. The Pfam protein families database. *Nucleic Acids Research*. 32:D138-41.
- Bavner A, Sanyal S, Gustafsson JA and Treuter E. 2005. Transcriptional corepression by SHP: molecular mechanisms and physiological consequences. *Trends in Endocrinology and Metabolism*. 16:478-488.
- Bertilsson L, Dahl ML, Dalen P and Al-Shurbaji A. 2002. Molecular genetics of CYP2D6: clinical relevance with focus on psychotropic drugs. *British Journal of Clinical Pharmacology*. 53:111-122.
- Birney E, Andrews TD, Bevan P, Caccamo M, Chen Y, Clarke L, Coates G, Cuff J, Curwen V, Cutts T, Down T, Eyraas E, Fernandez-Suarez XM, Gane P, Gibbins B, Gilbert J, Hammond M, Hotz H-R, Iyer V, Jekosch K, Kahari A, Kasprzyk A, Keefe D, Keenan S, Lehvaslaiho H, McVicker G, Melsopp C, Meidl P, Mongin E, Pettett R, Potter S, Proctor G, Rae M, Searle S, Slater G, Smedley D, Smith J, Spooner W, Stabenau A, Stalker J, Storey R, Ureta-Vidal A, Woodwark KC, Cameron G, Durbin R, Cox A, Hubbard T and Clamp M. 2004. An overview of Ensembl. *Genome Research*. 14:925-926

Birney E and Durbin R. 2000. Using GeneWise in the *Drosophila* annotation experiment. *Genome Research*. 10:547-548.

Brenner S, Johnson M, Bridgham J, Golda G, Lloyd DH, Johnson D, Luo S, McCurdy S, Foy M, Ewan M, Roth R, George D, Eletr S, Albrecht G, Vermaas E, Williams SR, Moon K, Burcham T, Pallas M, DuBridge RB, Kirchner J, Fearon K, Mao J and Corcoran K. 2000. Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays. *Nature Biotechnology*. 18:630-634.

Burset M, Seledtsov IA and Solovyev VV. 2001. SpliceDB: database of canonical and non-canonical mammalian splice sites. *Nucleic Acids Research*. 29:255-259.

Butcher NJ, Arulpragasam A, Goh HL, Davey T and Minchin RF. 2005. Genomic organization of human arylamine N-acetyltransferase Type I reveals alternative promoters that generate different 5'-UTR splice variants with altered translational activities. *Biochemical Journal*. 387:119-27.

Chantry-Darmon C, Urien C, Hayes H, Bertaud M, Chadi-Taourit S, Chardon P, Vaiman D and Rogel-Gaillard C. 2005. Construction of a cytogenetically anchored microsatellite map in rabbit. *Mammalian Genome*. 16:442-459.

Collier DA, Stober G, Li T, Heils A, Catalano M, Di Bella D, Arranz MJ, Murray RM, Vallada HP, Bengel D, Muller CR, Roberts GW, Smeraldi E, Kirov G, Sham P and Lesch KP. 1996. A novel functional polymorphism within the promoter of the serotonin transporter gene: possible role in susceptibility to affective disorders. *Molecular Psychiatry*. 1:453-60.

Consortium HG. 2004. Finishing the euchromatic sequence of the human genome. *Nature*. 431:931-945.

Consortium IH. 2003. The International HapMap project. *Nature*. 426:789-796

Consortium IH. 2005. A haplotype map of the human genome. *Nature*. 437:1299-1320

Dai Y, Hebert MF, Isoherranen N, Davis CL, Marsh C, Shen DD and Thummel KE. 2006. Effect of cyp3a5 polymorphism on tacrolimus metabolic clearance in vitro. *Drug Metabolism and Disposition*. 34:836-847.

Doddapaneni H, Chakraborty R and Yadav JS. 2005. Genome-wide structural and evolutionary analysis of the P450 monooxygenase genes (P450ome) in the white rot fungus *Phanerochaete chrysosporium*: evidence for gene duplications and extensive gene clustering. *BMC Genomics*. 6:92.

Donald PR, Sirgel FA, Venter A, Parkin DP, Seifart HI, van de Wal BW, Werely C, van Helden PD and Maritz JS. 2004. The influence of human N-acetyltransferase genotype on the early bactericidal activity of isoniazid. *Clinical Infection and Disease*. 39:1425-1430.

Feinberg AP, Cui H and Ohlsson R. 2002. DNA methylation and genomic imprinting: insights from cancer into epigenetic mechanisms. *Seminars in Cancer Biology*. 12:389-398.

Fredman D, Siegfried M, Yuan YP, Bork P, Lehvaslaiho H and Brookes AJ. 2002. HGVBase: a human sequence variation database emphasizing data quality and a broad spectrum of data sources. *Nucleic Acids Research*. 30:387-391.

Frikke-Schmidt R, Nordestgaard BG, Jensen GB and Tybjaerg-Hansen A. 2004. Genetic variation in ABC transporter A1 contributes to HDL cholesterol in the general population. *Journal of Clinical Investigation*. 114:1343-1353.

Furuta T, Shirai N, Sugimoto M, Nakamura A, Hishida A and Ishizaki T. 2005. Influence of CYP2C19 pharmacogenetic polymorphism on proton pump inhibitor-based therapies. *Drug Metabolism and Pharmacokinetics*. 20:153-167.

Gawronska-Szkwarz B, Lubinski J, Kladny J, Kurzawski G, Bielicki D, Wojcicki M, Sych Z and Musial HD. 1999. Polymorphism of GSTM1 gene in patients with colorectal cancer and colonic polyps. *Experimental and Toxicologic Pharmacology*. 51:321-5.

Gibbs RA, Weinstock GM, Metzker ML, Muzny DM, Sodergren EJ, Scherer S, Scott G, Steffen D, Worley KC, Burch PE, Okwuonu G, Hines S, Lewis L, DeRamo C, Delgado O, Dugan-Rocha S, Miner G, Morgan M, Hawes A, Gill R, Celera, Holt RA, Adams MD, Amanatides PG, Baden-Tillson H, Barnstead M, Chin S, Evans CA, Ferriera S, Fosler C, Glodek A, Gu Z, Jennings D, Kraft CL, Nguyen T, Pfannkoch CM, Sitter C, Sutton GG, Venter JC, Woodage T, Smith D, Lee HM, Gustafson E, Cahill P, Kana A, Doucette-Stamm L, Weinstock K, Fechtel K, Weiss RB, Dunn DM, Green ED, Blakesley RW, Bouffard GG, De Jong PJ, Osoegawa K, Zhu B, Marra M, Schein J, Bosdet I, Fjell C, Jones S, Krzywinski M, Mathewson C, Siddiqui A, Wye N, McPherson J, Zhao S, Fraser CM, Shetty J, Shatsman S, Geer K, Chen Y, Abramzon S, Nierman WC, Havlak PH, Chen R, Durbin KJ, Egan A, Ren Y, Song XZ, Li B, Liu Y, Qin X, Cawley S, Cooney AJ, D'Souza LM, Martin K, Wu JQ, Gonzalez-Garay ML, Jackson AR, Kalafus KJ, McLeod MP, Milosavljevic A, Virk D, Volkov A, Wheeler DA, Zhang Z, Bailey JA, Eichler EE, Tuzun E, Birney E, Mongin E, Ureta-Vidal A, Woodwark C, Zdobnov E, Bork P, Suyama M, Torrents D, Alexandersson M, Trask BJ, Young JM, Huang H, Wang H, Xing H, Daniels S, Gietzen D, Schmidt J, Stevens K, Vitt U, Wingrove J, Camara F, Mar Alba M, Abril JF, Guigo R, Smit A, Dubchak I, Rubin EM, Couronne O, Poliakov A, Hubner N, Ganten D, Goesele C, Hummel O, Kreitler T, Lee YA, Monti J, Schulz H, Zimdahl H, Himmelbauer H, Lehrach H, Jacob HJ, Bromberg S, Gullings-Handley J, Jensen-Seaman MI, Kwitek AE, Lazar J, Pasko D, Tonellato PJ, Twigger S, Ponting CP,

Duarte JM, Rice S, Goodstadt L, Beatson SA, Emes RD, Winter EE, Webber C, Brandt P, Nyakatura G, Adetobi M, Chiaromonte F, Elnitski L, Eswara P, Hardison RC, Hou M, Kolbe D, Makova K, Miller W, Nekrutenko A, Riemer C, Schwartz S, Taylor J, Yang S, Zhang Y, Lindpaintner K, Andrews TD, Caccamo M, Clamp M, Clarke L, Curwen V, Durbin R, Eyraes E, Searle SM, Cooper GM, Batzoglou S, Brudno M, Sidow A, Stone EA, Payseur BA, Bourque G, Lopez-Otin C, Puente XS, Chakrabarti K, Chatterji S, Dewey C, Pachter L, Bray N, Yap VB, Caspi A, Tesler G, Pevzner PA, Haussler D, Roskin KM, Baertsch R, Clawson H, Furey TS, Hinrichs AS, Karolchik D, Kent WJ, Rosenbloom KR, Trumbower H, Weirauch M, Cooper DN, Stenson PD, Ma B, Brent M, Arumugam M, Shteynberg D, Copley RR, Taylor MS, Riethman H, Mudunuri U, Peterson J, Guyer M, Felsenfeld A, Old S, Mockrin S and Collins F. 2004. Genome sequence of the Brown Norway rat yields insights into mammalian evolution. *Nature*. 428:493-521.

Gibson G, Plant N, Swales K, Ayrton A and El-Sankary W. 2002. Receptor-dependent transcriptional activation of cytochrome P450 3A genes: Induction mechanisms, species differences and inter-individual variation in man. *Xenobiotica*. 32:165-206

Gooderham NJ, Murray S, Lynch AM, Yadollahi-Farsani M, Zhao K, Boobis AR and Davies DS. 2001. Food-derived heterocyclic amine mutagens: Variable metabolism and significance to humans. *Drug Metabolism and Disposition*. 29:529-534

Guo W, Burris TP, Zhang YH, Huang BL, Mason J, Copeland KC, Kupfer SR, Pagon RA and McCabe ER. 1996. Genomic sequence of the DAX1 gene: an orphan nuclear receptor responsible for X-linked adrenal hypoplasia congenita and hypogonadotropic hypogonadism. *Journal of Clinical Endocrinology and Metabolism*. 81:2481-2486.

Handschin C, Podvinec M and Meyer U. 2000. CXR, a chicken xenobiotic-sensing orphan nuclear receptor is related to both mammalian pregnane X receptor (PXR) and constitutive androstane receptor (CAR). *Proceedings of the National Academy of Sciences USA*. 97:10769-10774

Hanke J, Brett D, Zastrow I, Aydin A, Delbruck S, Lehmann G, Luft F, Reich J and Bork P. 1999. Alternative Splicing of human genes: more the rule than the exception? *Trends in Genetics*. 15:389-390

Horie-Inoue K, Takayama K, Bono HU, Ouchi Y, Okazaki Y and Inoue S. 2006. Identification of novel steroid target genes through the combination of bioinformatics and functional analysis of hormone response elements. *Biochemical and Biophysical Research Communications*. 339:99-106.

Huang HD, Horng JT, Lee CC and Liu BJ. 2003. ProSplicer: a database of putative alternative splicing information derived from protein, mRNA and expressed sequence tag sequence data. *Genome Biology*. 4:R29

Hughes DA, Vilar FJ, Ward CC, Alfirovic A, Park BK and Pirmohamed M. 2004. Cost-effectiveness analysis of HLA B*5701 genotyping in preventing abacavir hypersensitivity. *Pharmacogenetics*. 14:335-42.

Jongeneel CV, Delorenzi M, Iseli C, Zhou D, Haudenschield CD, Khrebtukova I, Kuznetsov D, Stevenson BJ, Strausberg RL, Simpson AJ and Vasicek TJ. 2005. An atlas of human gene expression from massively parallel signature sequencing (MPSS). *Genome Research*. 15:1007-1014.

Jun JB, Cho DY, Kang C and Bae SC. 2005. Thiopurine S-methyltransferase polymorphisms and the relationship between the mutant alleles and the adverse effects in systemic lupus erythematosus patients taking azathioprine. *Clinical and Experimental Rheumatology*. 23:873-976.

Kirchheiner J and Brockmoller J. 2005. Clinical consequences of cytochrome P450 2C9 polymorphisms. *Clinical Pharmacology and Therapeutics*. 77:1-16.

Kuehl P, Zhang J, Lin Y, Lanba J, Assem M, Scheutz J, Watkins P, Daly A, Wrighton S, Hall S, Maurel P, Relling M, Brimer C, Yasuda K, Venkataramanan R, Strom S, Thummel K, Boguski M and Schuetz E. 2001. Sequence diversity in CYP3A promoters and characterisation of the genetic basis of polymorphic CYP3A5 expression. *Nature Genetics*. 27:383-391

Kumar S and Hedges SB. 1998. A molecular timescale for vertebrate evolution. *Nature*. 392:917-20.

Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, Funke R, Gage D, Harris K, Heaford A, Howland J, Kann L, Lehoczky J, LeVine R, McEwan P, McKernan K, Meldrim J, Mesirov JP, Miranda C, Morris W, Naylor J, Raymond C, Rosetti M, Santos R, Sheridan A, Sougnez C, Stange-Thomann N, Stojanovic N, Subramanian A, Wyman D, Rogers J, Sulston J, Ainscough R, Beck S, Bentley D, Burton J, Clee C, Carter N, Coulson A, Deadman R, Deloukas P, Dunham A, Dunham I, Durbin R, French L, Grafham D, Gregory S, Hubbard T, Humphray S, Hunt A, Jones M, Lloyd C, McMurray A, Matthews L, Mercer S, Milne S, Mullikin JC, Mungall A, Plumb R, Ross M, Shownkeen R, Sims S, Waterston RH, Wilson RK, Hillier LW, McPherson JD, Marra MA, Mardis ER, Fulton LA, Chinwalla AT, Pepin KH, Gish WR, Chissole SL, Wendl MC, Delehaunty KD, Miner TL, Delehaunty A, Kramer JB, Cook LL, Fulton RS, Johnson DL, Minx PJ, Clifton SW, Hawkins T, Branscomb E, Predki P, Richardson P, Wenning S, Slezak T, Doggett N, Cheng JF, Olsen A, Lucas S, Elkin C, Uberbacher E, Frazier M, Gibbs RA, Muzny DM, Scherer SE, Bouck JB, Sodergren EJ, Worley KC, Rives CM, Gorrell JH, Metzker ML, Naylor SL, Kucherlapati RS,

Nelson DL, Weinstock GM, Sakaki Y, Fujiyama A, Hattori M, Yada T, Toyoda A, Itoh T, Kawagoe C, Watanabe H, Totoki Y, Taylor T, Weissenbach J, Heilig R, Saurin W, Artiguenave F, Brottier P, Bruls T, Pelletier E, Robert C, Wincker P, Rosenthal A, Platzer M, Nyakatura G, Taudien S, Rump A, Yang HM, Yu J, Wang J, Huang GY, Gu J, Hood L, Rowen L, Madan A, Qin SZ, Davis RW, Federspiel NA, Abola AP, Proctor MJ, Myers RM, Schmutz J, Dickson M, Grimwood J, Cox DR, Olson MV, Kaul R, Shimizu N, Kawasaki K, Minoshima S, Evans GA, Athanasiou M, Schultz R, Roe BA, Chen F, Pan HQ, Ramser J, Lehrach H, Reinhardt R, McCombie WR, de la Bastide M, Dedhia N, Blocker H, Hornischer K, Nordsiek G, Agarwala R, Aravind L, Bailey JA, Bateman A, Batzoglou S, Birney E, Bork P, Brown DG, Burge CB, Cerutti L, Chen HC, Church D, Clamp M, Copley RR, Doerks T, Eddy SR, Eichler EE, Furey TS, Galagan J, Gilbert JGR, Harmon C, Hayashizaki Y, Haussler D, Hermjakob H, Hokamp K, Jang WH, Johnson LS, Jones TA, Kasif S, Kasprzyk A, Kennedy S, Kent WJ, Kitts P, Koonin EV, Korf I, Kulp D, Lancet D, Lowe TM, McLysaght A, Mikkelsen T, Moran JV, Mulder N, Pollara VJ, Ponting CP, Schuler G, Schultz JR, Slater G, Smit AFA, Stupka E, Szustakowski J, Thierry-Mieg D, Thierry-Mieg J, Wagner L, Wallis J, Wheeler R, Williams A, Wolf YI, Wolfe KH, Yang SP, Yeh RF, Collins F, Guyer MS, Peterson J, Felsenfeld A, Wetterstrand KA, Patrinos A and Morgan MJ. 2001. Initial sequencing and analysis of the human genome. *Nature*. 409:860-921

Lareau LF, Green RE, Bhatnagar RS and Brenner SE. 2004. The evolving roles of alternative splicing. *Current Opinion in Structural Biology*. 14:273-282.

Lee C, Atanelov L, Modrek B and Xing Y. 2003. ASAP: the Alternative Splicing Annotation Project. *Nucleic Acids Research*. 31:101-105.

Lim YP, Liu CH, Shyu LJ and Huang JD. 2005. Functional characterization of a novel polymorphism of pregnane X receptor, Q158K, in Chinese subjects. *Pharmacogenetics and Genomics*. 15:337-341.

Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, Clamp M, Chang JL, Kulbokas EJ, 3rd, Zody MC, Mauceli E, Xie X, Breen M, Wayne RK, Ostrander EA, Ponting CP, Galibert F, Smith DR, DeJong PJ, Kirkness E, Alvarez P, Biagi T, Brockman W, Butler J, Chin CW, Cook A, Cuff J, Daly MJ, DeCaprio D, Gnerre S, Grabherr M, Kellis M, Kleber M, Bardeleben C, Goodstadt L, Heger A, Hitte C, Kim L, Koepfli KP, Parker HG, Pollinger JP, Searle SM, Sutter NB, Thomas R, Webber C, Baldwin J, Abebe A, Abouelleil A, Aftuck L, Ait-Zahra M, Aldredge T, Allen N, An P, Anderson S, Antoine C, Arachchi H, Aslam A, Ayotte L, Bachantsang P, Barry A, Bayul T, Benamara M, Berlin A, Bessette D, Blitshteyn B, Bloom T, Blye J, Boguslavskiy L, Bonnet C, Boukhgalter B, Brown A, Cahill P, Calixte N, Camarata J, Cheshatsang Y, Chu J, Citroen M, Collymore A, Cooke P, Dawoe T, Daza R, Decktor K, DeGray S, Dhargay N, Dooley K, Dooley K, Dorje P, Dorjee K, Dorris L, Duffey N, Dupes A, Egbiremolen O, Elong R, Falk J, Farina A, Faro S, Ferguson D, Ferreira P, Fisher S, FitzGerald M, Foley K, Foley C, Franke A, Friedrich D, Gage D, Garber M, Gearin G, Giannoukos G, Goode T, Goyette A, Graham J, Grandbois E, Gyaltzen K, Hafez N, Hagopian D, Hagos B, Hall J, Healy C, Hegarty R, Honan T, Horn A, Houde N, Hughes L, Hunnicutt L, Husby M, Jester B, Jones C, Kamat A, Kanga B, Kells C, Khazanovich D, Kieu AC, Kisner P, Kumar M, Lance K, Landers T, Lara M, Lee W, Leger JP, Lennon N, Leuper L, LeVine S, Liu J, Liu X, Lokyitsang Y, Lokyitsang T, Lui A, Macdonald J, Major J, Marabella R, Maru K, Matthews C, McDonough S, Mehta T, Meldrim J, Melnikov A, Meneus L, Mihalev A, Mihova T, Miller K, Mittelman R, Mlenga V, Mulrain L, Munson G, Navidi A,

Naylor J, Nguyen T, Nguyen N, Nguyen C, Nguyen T, Nicol R, Norbu N, Norbu C, Novod N, Nyima T, Olandt P, O'Neill B, O'Neill K, Osman S, Oyono L, Patti C, Perrin D, Phunkhang P, Pierre F, Priest M, Rachupka A, Raghuraman S, Rameau R, Ray V, Raymond C, Rege F, Rise C, Rogers J, Rogov P, Sahalie J, Settipalli S, Sharpe T, Shea T, Sheehan M, Sherpa N, Shi J, Shih D, Sloan J, Smith C, Sparrow T, Stalker J, Stange-Thomann N, Stavropoulos S, Stone C, Stone S, Sykes S, Tchuinga P, Tenzing P, Tesfaye S, Thoulutsang D, Thoulutsang Y, Topham K, Topping I, Tsamla T, Vassiliev H, Venkataraman V, Vo A, Wangchuk T, Wangdi T, Weiland M, Wilkinson J, Wilson A, Yadav S, Yang S, Yang X, Young G, Yu Q, Zainoun J, Zembek L, Zimmer A and Lander ES. 2005. Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature*. 438:803-819.

Loots GG and Ovcharenko I. 2004. rVISTA 2.0: evolutionary analysis of transcription factor binding sites. *Nucleic Acids Research*. 32:W217-221.

Loots GG, Ovcharenko I, Pachter L, Dubchak I and Rubin EM. 2002. rVista for comparative sequence-based discovery of functional transcription factor binding sites. *Genome Research*. 12:832-839.

Mandel S, Weinreb O and Youdim MB. 2003. Using cDNA microarray to assess Parkinson's disease models and the effects of neuroprotective drugs. *Trends in Pharmacological Sciences*. 24:184-91.

Mariadason JM, Arango D, Corner GA, Aranes MJ, Hotchkiss KA, Yang W and Augenlicht LH. 2002. A gene expression profile that defines colon cell maturation in vitro. *Cancer Research*. 62:4791-804.

Murata M, Watanabe M, Yamanaka M, Kubota Y, Ito H, Nagao M, Katoh T, Kamataki T, Kawamura J, Yatani R and Shiraishi T. 2001. Genetic polymorphisms in cytochrome P450 (CYP) 1A1, CYP1A2, CYP2E1, glutathione S-transferase (GST)

M1 and GSTT1 and susceptibility to prostate cancer in the Japanese population. *Cancer Letters*. 165:171-177

Natsoulis G, El Ghaoui L, Lanckriet GR, Tolley AM, Leroy F, Dunlea S, Eynon BP, Pearson CI, Tugendreich S and Jarnagin K. 2005. Classification of a large microarray data set: algorithm comparison and analysis of drug signatures. *Genome Research*. 15:724-736.

Nebert DW and Gonzalez FJ. 1987. P450 genes: structure, evolution, and regulation. *Annual Review of Biochemistry*. 56:945-993.

Nelson DR. 2002. Mining databases for cytochrome P450 genes. *Methods in Enzymology*. 357:3-15.

Nelson DR. 2003. Comparison of P450s from human and fugu: 420 million years of vertebrate P450 evolution. *Archives of Biochemistry and Biophysics*. 409:18-24.

Nelson DR. 2005. Gene nomenclature by default, or BLASTing to Babel. *Human Genomics*. 2:196-201.

Nelson DR, Koymans L, Kamataki T, Stegeman JJ, Feyereisen R, Waxman DJ, Waterman MR, Gotoh O, Coon MJ, Estabrook RW, Gunsalus IC and Nebert DW. 1996. P450 superfamily: Update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics*. 6:1-42

Nelson DR, Schuler MA, Paquette SM, Werck-Reichhart D and Bak S. 2004. Comparative genomics of rice and Arabidopsis. Analysis of 727 cytochrome P450 genes and pseudogenes from a monocot and a dicot. *Plant Physiology*. 135:756-772.

Neuhaus T, Ko YD, Lorenzen K, Fronhoffs S, Harth V, Brode P, Vetter H, Bolt HM, Pesch B and Bruning T. 2004. Association of cytochrome P450 2E1 polymorphisms and head and neck squamous cell cancer. *Toxicology Letters*. 151:273-282.

Niakan KK and McCabe ER. 2005. DAX1 origin, function, and novel role. *Molecular Genetics and Metabolism*. 86:70-83.

Niemi M, Backman JT, Kajosaari LI, Leathart JB, Neuvonen M, Daly AK, Eichelbaum M, Kivisto KT and Neuvonen PJ. 2005. Polymorphic organic anion transporting polypeptide 1B1 is a major determinant of repaglinide pharmacokinetics. *Clinical Pharmacology and Therapeutics*. 77:468-478.

Niemi M, Schaeffeler E, Lang T, Fromm MF, Neuvonen M, Kyrklund C, Backman JT, Kerb R, Schwab M, Neuvonen PJ, Eichelbaum M and Kivisto KT. 2004. High plasma pravastatin concentrations are associated with single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide-C (OATP-C, SLCO1B1). *Pharmacogenetics*. 14:429-440.

Ozyildirim AM, Wistow GJ, Gao J, Wang J, Dickinson DP, Frierson HF, Jr. and Laurie GW. 2005. The lacrimal gland transcriptome is an unusually rich source of rare and poorly characterized gene transcripts. *Investigations in Ophthalmology and Visual Science*. 46:1572-1580.

Pennacchio LA and Rubin EM. 2003. Comparative genomic tools and database: Providing insights into the human genome. *Journal of Clinical Investigation*. 111:1099-1106

Petry CJ, Ong KK, Michelson KF, Artigas S, Wingate DL, Balen AH, de Zegher F, Ibanez L and Dunger DB. 2005. Association of aromatase (CYP 19) gene variation with features of hyperandrogenism in two populations of young women. *Human Reproduction*. 20:1837-1843.

Rebbeck. 1999. Modification of clinical presentation of prostate tumors by a novel genetic variant in CYP3A4 (vol 90, pg 1225 1998). *Journal of the National Cancer Institute*. 91:1082

Rogler CE, Tchaikovskaya T, Norel R, Massimi A, Plescia C, Rubashevsky E, Siebert P and Rogler LE. 2004. RNA expression microarrays (REMs), a high-throughput method to measure differences in gene expression in diverse biological samples. *Nucleic Acids Research*. 32:e120.

Sachse C, Brochmoller J, Bauer S and Roots I. 1999. Functional significance of a C - > A polymorphism in intron I of the cytochrome P450 CYP1A2 gene tested with caffeine. *British Journal of Clinical Pharmacology*. 47:445-449

Saitoh A, Singh KK, Powell CA, Fenton T, Fletcher CV, Brundage R, Starr S and Spector SA. 2005. An MDR1-3435 variant is associated with higher plasma nelfinavir levels and more rapid virologic response in HIV-1 infected children. *AIDS*. 19:371-80.

Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R and Miller W. 2000. PipMaker--a web server for aligning two genomic DNA sequences. *Genome Research*. 10:577-586.

Screaton GR, Xu XN, Olsen AL, Cowper AE, Tan R, McMichael AJ and Bell JL. 1997. LARD: a new lymphoid-specific death domain containing receptor regulated by alternative pre-mRNA splicing. *Proceedings of the National Academy of Science USA*. 94:4615-4619.

Sherry ST, Ward M and Sirotkin K. 1999. dbSNP-database for single nucleotide polymorphisms and other classes of minor genetic variation. *Genome Research*. 9:677-679.

Song X, Xie M, Zhang H, Li Y, Sachdeva K and Yan B. 2004. The pregnane X receptor binds to response elements in a genomic context-dependent manner, and PXR activator rifampicin selectively alters the binding among target genes. *Drug Metabolism and Disposition*. 32:35-42.

Spiecker M, Darius H, Hankeln T, Soufi M, Sattler AM, Schaefer JR, Node K, Borgel J, Mugge A, Lindpaintner K, Huesing A, Maisch B, Zeldin DC and Liao JK. 2004. Risk of coronary artery disease associated with polymorphism of the cytochrome P450 epoxigenase CYP2J2. *Circulation*. 110:2132-6. Epub 2004 Oct 4.

Steketee K, Ziel-van der Made AC, van der Korput HA, Houtsmuller AB and Trapman J. 2004. A bioinformatics-based functional analysis shows that the specifically androgen-regulated gene SARG contains an active direct repeat androgen response element in the first intron. *Journal of Molecular Endocrinology*. 33:477-491.

Thanaraj TA, Stamm S, Clark F, Riethoven JJ, Le Texier V and Muilu J. 2004. ASD: the Alternative Splicing Database. *Nucleic Acids Research*. 32:D64-69.

Thorisson GA and Stein LD. 2003. The SNP Consortium website: past, present and future. *Nucleic Acids Research*. 31:124-127.

Tukey RH and Strassburg CP. 2000. Human UDP-glucuronosyltransferases: metabolism, expression, and disease. *Annu Rev Pharmacol Toxicol*. 40:581-616.

Vizirianakis IS. 2004. Challenges in current drug delivery from the potential application of pharmacogenomics and personalized medicine in clinical practice. *Current Drug Delivery*. 1:73-80.

Wang Y, O'Brate A, Zhou W and Giannakakou P. 2005. Resistance to microtubule-stabilizing drugs involves two events: beta-tubulin mutation in one allele followed by loss of the second allele. *Cell Cycle*. 4:1847-1853.

Wang Z, Chen Y and Li Y. 2004. A brief review of computational gene prediction methods. *Genomics Proteomics Bioinformatics*. 2:216-221.

Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P, Antonarakis SE, Attwood J, Baertsch R, Bailey J, Barlow K, Beck S, Berry E, Birren B, Bloom T, Bork P, Botcherby M, Bray

N, Brent MR, Brown DG, Brown SD, Bult C, Burton J, Butler J, Campbell RD,
 Carninci P, Cawley S, Chiaromonte F, Chinwalla AT, Church DM, Clamp M, Clee C,
 Collins FS, Cook LL, Copley RR, Coulson A, Couronne O, Cuff J, Curwen V, Cutts
 T, Daly M, David R, Davies J, Delehaunty KD, Deri J, Dermitzakis ET, Dewey C,
 Dickens NJ, Diekhans M, Dodge S, Dubchak I, Dunn DM, Eddy SR, Elnitski L, Emes
 RD, Eswara P, Eyraas E, Felsenfeld A, Fewell GA, Flicek P, Foley K, Frankel WN,
 Fulton LA, Fulton RS, Furey TS, Gage D, Gibbs RA, Glusman G, Gnerre S, Goldman
 N, Goodstadt L, Grafham D, Graves TA, Green ED, Gregory S, Guigo R, Guyer M,
 Hardison RC, Haussler D, Hayashizaki Y, Hillier LW, Hinrichs A, Hlavina W, Holzer
 T, Hsu F, Hua A, Hubbard T, Hunt A, Jackson I, Jaffe DB, Johnson LS, Jones M,
 Jones TA, Joy A, Kamal M, Karlsson EK, Karolchik D, Kasprzyk A, Kawai J,
 Keibler E, Kells C, Kent WJ, Kirby A, Kolbe DL, Korf I, Kucherlapati RS, Kulbokas
 EJ, Kulp D, Landers T, Leger JP, Leonard S, Letunic I, Levine R, Li J, Li M, Lloyd
 C, Lucas S, Ma B, Maglott DR, Mardis ER, Matthews L, Mauceli E, Mayer JH,
 McCarthy M, McCombie WR, McLaren S, McLay K, McPherson JD, Meldrim J,
 Meredith B, Mesirov JP, Miller W, Miner TL, Mongin E, Montgomery KT, Morgan
 M, Mott R, Mullikin JC, Muzny DM, Nash WE, Nelson JO, Nhan MN, Nicol R, Ning
 Z, Nusbaum C, O'Connor MJ, Okazaki Y, Oliver K, Larty EO, Pachter L, Parra G,
 Pepin KH, Peterson J, Pevzner P, Plumb R, Pohl CS, Poliakov A, Ponce TC, Ponting
 CP, Potter S, Quail M, Reymond A, Roe BA, Roskin KM, Rubin EM, Rust AG,
 Santos R, Sapojnikov V, Schultz B, Schultz J, Schwartz MS, Schwartz S, Scott C,
 Seaman S, Searle S, Sharpe T, Sheridan A, Shownkeen R, Sims S, Singer JB, Slater
 G, Smit A, Smith DR, Spencer B, Stabenau A, Strange-Thomann NS, Sugnet C,
 Suyama M, Tesler G, Thompson J, Torrents D, Trevaskis E, Tromp J, Ucla C, Vidal
 AU, Vinson JP, von Niederhausern AC, Wade CM, Wall M, Weber RJ, Weiss RB,

Wendl MC, West AP, Wetterstrand K, Wheeler R, Whelan S, Wierzbowski J, Willey D, Williams S, Wilson RK, Winter E, Worley KC, Wyman D, Yang S, Yang SP, Zdobnov EM, Zody MC and Lander ES. 2002. Initial sequencing and comparative analysis of the mouse genome. *Nature*. 420:520-562

Werck-Reichhart D and Feyereisen R. 2000. Cytochromes P450: a success story. *Genome Biology*. 1:REVIEWS3003. Epub 2000 Dec 8.

Wren JD, Forgacs E, Fondon JW, Pertsemlidis A, Cheng S, Gallardo T, Williams RS, Shohet RV, Minna JD and Garner HR. 2000. Repeat polymorphisms within gene regions: Phenotypic and evolutionary implications. *American Journal of Human Genetics*. 67:345-356.

Xie W, Barwick JL, Simmon CM, Pierce AM, Safe S, Blumberg B, Guzelian PS and Evans RM. 2000. Reciprocal activation of xenobiotic response genes by nuclear receptors SXR/PXR and CAR. *Genes and Development*. 14:3014-3023

Xing Y, Xu Q and Lee C. 2003. Widespread production of novel soluble protein isoforms by alternative splicing removal of transmembrane anchoring domains. *FEBS Letters*. 555:572-578.

Zhang H, Ahmadi A, Arbman G, Zdolsek J, Carstensen J, Nordenskjold B, Soderkvist P and Sun XF. 1999. Glutathione S-transferase T1 and M1 genotypes in normal mucosa, transitional mucosa and colorectal adenocarcinoma. *International Journal of Cancer*. 84:135-8.

Zhang Z, Burch PE, Cooney AJ, Lanz RB, Pereira FA, Wu J, Gibbs RA, Weinstock G and Wheeler DA. 2004. Genomic analysis of the nuclear receptor family: new insights into structure, regulation, and evolution from the rat genome. *Genome Research*. 14:580-90.

Group	Number of Sequencing Projects
Bacteria	472
Fungi	508
Plants	1740
Lower Eukaryotes	203
Animals	1581
TOTAL	4504

Table 1: Current genome sequencing projects

(Data sourced from NCBI March 2006)

Order	Species	Genome Status	Reference
Primates	Human	Complete. Build 35.1	(Lander, et al. 2001)
Primates	Marmoset	In Progress	N/A
Carnivora	Dog	Complete Build 2.1	(Lindblad-Toh, et al. 2005)
Lagomorpha	Rabbit	Microsatellite map	(Chantry-Darmon, et al. 2005)
Rodentia	Rat	Complete. Build 3.1	(Gibbs, et al. 2004)
Rodentia	Mouse	Complete. Build 35.1	(Waterston, et al. 2002)
Rodentia	Hamster	Not Started	N/A

Table 2: Status of whole genome sequencing projects in ADME-relevant mammalian species.

Database	URL	Description	Reference
Alternative Splicing Database (ASDB)	http://www.ebi.ac.uk/asd/	Splice variants from annotated genes for human	(Thanaraj, et al. 2004)
Alternative Splicing Annotation Project	http://bioinfo.mbi.ucla.edu/ASAP/	EST based splice variant prediction with tissue information	(Lee, et al. 2003)
ProSplicer	http://prosplicer.mbc.nctu.edu.tw/	Putative alternative splice variant database for human	(Huang, et al. 2003)
SpliceDB	http://www.softberry.com/berry.phtml	Canonical and non-canonical splice sites in mammals	(Burset, et al. 2001)

Table 3: Alternative Splice Variant Databases

Gene with SNPs	Disease State	Reference
CYP19	Hyperandrogenism	(Petry, et al. 2005)
CYP2E1	Head/neck squamous cell cancer	(Neuhaus, et al. 2004)
CYP2J2	Coronary Heart Disease	(Spiecker, et al. 2004)
CYP1A1	Prostate Cancer	(Murata, et al. 2001)
CYP3A4	Prostate Cancer	(Rebbeck 1999)
GSTM1	Colorectal cancer	(Gawronska-Szklarz, et al. 1999)
GSTT1	Colorectal cancer	(Zhang, et al. 1999)
Serotonin Transporter	Affective Disorder	(Collier, et al. 1996)
ABCA1	Coronary heart disease	(Frikke-Schmidt, et al. 2004)

Table 4: Human SNPs associated with an increased predisposition to disease states

Gene with SNP	Drug Effected	Reference
CYP1A2	Caffeine	(Sachse, et al. 1999)
CYP2C9	Warfarin	(Kirchheiner and Brockmoller 2005)
CYP2C19	Omeprazole	(Furuta, et al. 2005)
CYP2D6	Psychotropic Drugs	(Bertilsson, et al. 2002)
CYP3A5	Tacrolimus	(Dai, et al. 2006)
NAT2	Isoniazid	(Donald, et al. 2004)
TMPT	Azothioprine	(Jun, et al. 2005)
UGT1A1	Irinotecan	(Ando, et al. 2005)
MDR1	Nelfinavir	(Saitoh, et al. 2005)
OATP2	Pravastatin, Repaglinide	(Niemi, et al. 2005, Niemi, et al. 2004)
PXR	Rifampicin and Paclitaxel	(Lim, et al. 2005)
HLA B	Abacavir	(Hughes, et al. 2004)
β -tubulin	Taxol/Epothilones	(Wang, et al. 2005)

Table 4: Human SNPs associated with altered response to therapeutic agents