Intruder dominance in the 0_2^+ state of ^{32}Mg studied with a novel technique for in-flight decays

R. Elder,^{1,2} H. Iwasaki,^{1,2} J. Ash,^{1,2} D. Bazin,^{1,2} P. C. Bender,^{1,3} T. Braunroth,⁴ B. A. Brown,^{1,2}
C. M. Campbell,⁵ H. L. Crawford,⁵ B. Elman,^{1,2} A. Gade,^{1,2} M. Grindler,^{1,2} N. Kobayashi,⁶ B. Longfellow,^{1,2}
A. O. Macchiavelli,⁵ T. Mijatović,^{1,7} J. Pereira,¹ A. Revel,¹ D. Rhodes,^{1,2} J. A. Tostevin,⁸ D. Weisshaar,¹
¹*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824, USA*
²*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA*
³*Department of Physics, University of Massachusetts Lowell, Lowell, Massachusetts 01854, USA*
⁴*Institut für Kernphysik, Universität zu Köln, 50937 Köln, Germany*
⁵*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
⁶*Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan*
⁷*Ruder Bošković Institute, HR-10 002 Zagreb, Croatia*
⁸*Department of Physics, University of Surrey, Guildford, Surrey GU2 7XH, United Kingdom*
(Dated: July 25, 2019)

The neutron-rich isotope ^{32}Mg is near the center of the $N = 20$ island of inversion where intruder configurations of cross-shell particle-hole excitations are energetically favored relative to the normal-order configurations. The classic picture of the inversion between the normal $0p0h$ and intruder $2p2h$ configurations may not be sufficient to describe the measured properties for the 0_2^+ state in ^{32}Mg such as the excitation energy and the (t, p) reaction cross section. More recently, a significant admixture of the intruder $4p4h$ configurations has been indicated in ^{32}Mg .

The development of advanced gamma-ray tracking arrays allows for a sensitive new technique to investigate elusive states of exotic nuclei with fast rare-isotope beams. In this study, we present a novel technique developed to identify in-flight isomeric decays of ^{32}Mg by taking advantage of the excellent energy and position resolution of the detector array GRETINA. We confirm the $0_2^+ \rightarrow 2_1^+$ gamma-ray transition of ^{32}Mg and constrain the 0_2^+ decay lifetime. The results are discussed in the context of admixtures of the normal and intruder configurations.

The existence of an island of inversion was first postulated to explain the unexpected excess binding for certain neutron-rich isotopes near the canonical $N = 20$ magic number [1–3]. Since then, similar islands of inversion have been proposed at $N = 8, 20, 28, 40$, [4–11] and more recently at 50 [12], where it has been suggested that the nuclear shell structure is dramatically modified, leading to the appearance of deformed nuclei near these magic numbers. The physics mechanisms underlying the structural changes in neutron-rich nuclei have been explored in recent decades, highlighting important aspects of nuclear interactions, such as the tensor and the three-body forces [13–16]. Many low-energy properties in the $N = 20$ island of inversion can be explained by the intruder $2p2h$ configurations dominating over the normal $0p0h$ configurations [2, 17, 18]. However, recent studies indicated that the intruder $4p4h$ configurations are also important [14, 15], calling into question the simple picture of the structural changes and configuration mixings in the islands of inversion.

In this work, we report the lifetime and reaction population of the 0_2^+ state of ^{32}Mg to examine the configuration mixing near the center of the $N = 20$ island of inversion. The $B(E2; 2_1^+ \rightarrow 0_2^+)$ value from the measured lifetime allows for a comparison of the quadrupole collectivity between the $2_1^+ \rightarrow 0_2^+$ and $2_1^+ \rightarrow 0_1^+$ transitions. Meanwhile, the two-proton removal cross section from the $^9\text{Be}(^{34}\text{Si}, ^{32}\text{Mg})X$ reaction is sensitive to the overlap between the ^{32}Mg 0_2^+ state and the ^{34}Si 0_1^+ state, the latter being dominated by $0p0h$ configurations [17, 19].

Many other ^{32}Mg properties such as the excess binding [1], the reduced 2_1^+ energy [7], and the enhanced $B(E2; 2_1^+ \rightarrow 0_1^+)$ value [5, 20–24] are consistent with $2p2h$ configurations dominating the low-energy states. However, the $4p4h$ configurations appear essential to reproduce the known properties of the long-lived ($\tau > 10$ ns) 0_2^+ state reported at 1058 keV [6]. In agreement with the importance of the $4p4h$ configurations suggested in recent shell-model studies [14, 15], a schematic model consisting of only three configurations, $0p0h$, $2p2h$, and $4p4h$, successfully characterized the structure of the 0_2^+ state of ^{32}Mg [25, 26]. This elusive 0_2^+ state has not been observed since the original discovery in 2010 [6], therefore an independent confirmation of this state and additional lifetime and reaction information are needed to validate the importance of the $4p4h$ configurations.

To study the 0_2^+ state of ^{32}Mg , we developed a novel in-beam spectroscopy technique that is now made possible by taking advantage of advanced gamma-ray tracking arrays. At relativistic beam velocities, the 0_2^+ state decays over an average range on the order of 1 m past the target. Therefore, in conventional in-beam experiments, the decay location of the 0_2^+ state remains unknown and the gamma-ray energies cannot be corrected for Doppler-shift. In our technique, we overcome this difficulty by tracking the isomeric decay position, as detailed later in the text. This technique can be applied more generally to in-beam experiments involving isomers with lifetimes on the order of nanoseconds depending on gamma-ray yield.

This experiment was performed at the NSCL Coupled Cyclotron Facility [27] using a ^{48}Ca primary beam at 140 MeV/nucleon on a ^9Be production target. A ^{34}Si secondary beam at 86 MeV/nucleon was selected by the A1900 fragment separator [28] with a purity of 82% and an intensity of 9×10^5 pps. The 0_2^+ state of ^{32}Mg was populated in the $^9\text{Be}(^{34}\text{Si}, ^{32}\text{Mg})\text{X}$ reaction on a 0.57 g/cm^2 -thick ^9Be target using the setup shown in Fig. 1. To validate our method, we examined the ^{31}Mg products created simultaneously from the same setup which populates a $(7/2^-)$ isomer at 461 keV with a lifetime of $\tau = 15.1(12) \text{ ns}$ [29]. Reaction products were identified by time-of-flight and energy-loss measurements from the S800 spectrograph [30].

FIG. 1. The present experimental setup. The $^9\text{Be}(^{34}\text{Si}, ^{32}\text{Mg})\text{X}$ reaction populates the 0_2^+ isomer which emits a cascade of gamma rays γ_1 and γ_2 at angles θ_1 and θ_2 relative to the ion trajectory.

In general, the in-flight detection of isomers with a lifetime from 1 ns to 100 ns is challenging because decays occur along a flight path on the order of meters. When the isomer produces a cascade of gamma rays emitted nearly simultaneously, one solution is to use the timing information of the cascade to locate the common decay position [31]. Our approach is similar except we use the energy information of the gamma-ray cascade, which is now feasible by using the excellent position and energy resolution of GRETINA [32]. In practice, if the energy of one observed gamma ray γ_1 is consistent with the Doppler-shifted energy of the known transition, then we assume it is associated with the gamma-ray cascade emitted from the isomer. In the ^{32}Mg case, the 885-keV transition has laboratory-frame energy in the range from about 610 to 1290 keV. Then γ_1 is used to find the emission angle θ_1 , which, when coupled with the gamma-ray hit position information from GRETINA, allows us to locate the decay position. We can then determine the emission angle θ_2 for the other gamma ray γ_2 observed in coincidence. At our ion velocity of $0.353c$, if the lifetime of the 0_2^+ state in ^{32}Mg is $\tau \approx 10 \text{ ns}$, the average flight path is approximately 1 m. Therefore, to cover the wide range of possible decay positions, the ^9Be target was placed 72 cm upstream of the center of GRETINA (Fig. 1). To improve the signal-to-background ratio for detecting isomeric decays, a cylindrical lead shield was installed downstream

of the target to attenuate the prompt gamma rays from short-lived states such as the 2_1^+ and 4_1^+ states of ^{32}Mg . GRETINA was arranged to have 4 detector modules at 58 degrees, 2 at 90 degrees, and 4 at 122 degrees relative to the beam axis measured from the center of GRETINA. In this arrangement GRETINA was most efficient for decays occurring 52 cm to 92 cm downstream of the target, corresponding to $\pm 20 \text{ cm}$ from the center of GRETINA, and we selected events within this range to reduce the background.

The decays we analyzed emit two gamma rays in cascade that both may interact with GRETINA multiple times. The highly segmented geometry of GRETINA allows one to distinguish the multiple interaction points but some criteria must be applied to determine which interaction points belong to each incoming gamma ray, and which of those points is the first interaction point within GRETINA. Following the technique used in Ref. [32], the interaction point with the largest energy deposit was chosen as the first interaction point of one gamma ray. Then an addback routine analogous to that used in Ref. [33] was implemented to sum energies within an $r = 80 \text{ mm}$ sphere centered on the first interaction point. Using the remaining interaction points, a second gamma ray was reconstructed with the first interaction point and addback energy found in the same manner. In general, if interaction points still remain, the same addback routine can be repeated to define additional gamma rays. However, to improve the signal-to-noise ratio, we selected events with exactly two interaction spheres (gamma-ray multiplicity two). As mentioned earlier, if one gamma ray γ_1 is assigned to the known transition, then the Doppler-corrected energy of the other gamma ray γ_2 is determined event by event based on the common decay position.

First, we demonstrate the new technique by applying it to the ^{31}Mg products as shown in Fig. 2a together with a partial level scheme. In this case, the 171-keV transition from the 221-keV excited state was used as a reference to determine the decay location of the isomer at 461 keV. The peak in Fig. 2a was fit with a Gaussian with a centroid energy of 244(5) keV which is consistent with the literature value of 239.9(5) keV. The low energy of the 240-keV gamma ray suggests that all interactions are likely to occur within a 20-mm sphere. Therefore, to improve the sensitivity to events of interest, an additional gate requiring the 240-keV gamma ray to deposit all energy within $r = 20 \text{ mm}$ of the first interaction point was implemented. Including this gate results in the lower, filled spectrum of Fig. 2a which shows a reduced background while retaining 66% of the peak counts. The scaled background spectrum in black in Fig. 2 was obtained by analyzing all ^{34}Si reaction products excluding the Mg isotopes, which mostly consists of ^{34}Si and ^{33}Al .

The energy spectrum of ^{31}Mg was studied further with a GEANT4 simulation [33, 34] that included the isomeric 461-keV state, the lower-lying 221- and 50-keV states, and the gamma-ray transitions at 240, 221, 171, and 50 keV, as shown in the level scheme of Fig. 2a. The

FIG. 2. Doppler-corrected gamma-ray energy spectra are shown for data (blue circles), background from the scaled response of beam products excluding the Mg isotopes (black lines), and the sums of the simulated and background responses (red lines). ^{31}Mg is shown (a) in the upper spectra and the constraint that the 244-keV transition only interacts within $r = 20$ mm of its first interaction is included in the lower, filled spectrum (scaled by 0.5). ^{32}Mg is shown (b) with the $r = 20$ mm gate applied to the 170-keV gamma ray in the upper spectrum (scaled by 0.6) and with additional gates requiring proper identification of the interaction points of the 885-keV gamma ray in the lower, filled spectrum. Insets show the number of decays near (250 to 525 mm), at mid-distance (525 to 700 mm) and far from the target (700 to 1000 mm). The data (open circles) is compared to simulations assuming lifetimes of 5 ns (red solid line), 15 ns (blue dashed line), and 45 ns (black dotted line).

simulated spectrum was added to the background and scaled to fit the measured peak at 244 keV. The simulated energy of the transition from the isomer that best reproduced the data was 239(1) keV, as is shown with the red line in Fig. 2a. The difference between the best-fit energy of 239 keV and the peak centroid energy of 244 keV was attributed to the $\tau = 192$ ps lifetime of the 221-keV state, corresponding to an average distance of 2 cm between the emission points of the 171-keV and the 240-keV gamma rays.

The lifetime of the 461-keV state was studied using the decay distribution along the beam line shown in the inset of Fig. 2a. Although GRETTINA is most efficient for decays from 52 cm to 92 cm downstream of the target, the decay trend was analyzed in a larger region from 25 to 100 cm past the target to improve the sensitivity

to the lifetime. The decay distribution does not change significantly for lifetimes greater than 10 ns so we cannot place an upper limit on the lifetime of the 461-keV state. However, this data places a 1σ lower limit of 9 ns which is consistent with the known lifetime of $\tau = 15.1(12)$ ns [35].

The ^{32}Mg result is shown in Fig. 2b where the same analysis approach is used except now the 885-keV transition from the 2_1^+ state of ^{32}Mg is used as a reference to find the decay location. Since the expected energy of the $0_2^+ \rightarrow 2_1^+$ transition (172 keV) is low, similar to the 240-keV transition of ^{31}Mg , we used the same condition that the gamma ray deposits its energy within $r = 20$ mm of the first interaction point. The upper spectrum of Fig. 2b shows a peak-like structure close to 170 keV corresponding to the $0_2^+ \rightarrow 2_1^+$ transition [6]. The background distribution is reproduced by analyzing to other products of the ^{34}Si beam and scaling the result as shown in black. To understand the significance of the peak-like structure at 170 keV the measured spectrum was compared to a simulation including the 0_2^+ isomer at 1058 keV and the cascade of gamma rays with energies 172 and 885 keV. The 2_1^+ state was also included in the simulation using the lifetime value $\tau = 16(3)$ ps determined from $B(E2)$ results [5, 20–24, 36]. The simulated distribution was added to the background distribution and scaled to fit the peak at 170 keV as shown by the red line.

The 170-keV peak was unambiguously confirmed by applying additional gates to the 885-keV candidate. The result is shown in Fig. 2b as the lower, filled spectrum presenting a clear signal with reduced background. In analogy with the $r = 20$ mm gate applied to the 170-keV transition in ^{32}Mg , a gate was applied for the 885-keV gamma ray that requires all interaction points to lie within 60 mm of the first interaction point. Additionally, we used the interaction point information of the detected 885-keV gamma ray to test if it is consistent with Compton scattering. The energies of the first interaction and remaining interactions were used in the Compton scattering formula to obtain the scattering angle (see eq. (21) of Ref. [37]). If this angle agreed within 0.7 rad with the scattering angle deduced from the decay position and interaction position information, the event passed the gate. In this work the $0_2^+ \rightarrow 2_1^+$ energy is $165 \pm 4(\text{stat}) \pm 2(\text{syst})$ keV which is included in the simulated response in Fig. 2b. The 7 keV difference between this measurement and the previous measurement of 172(2) keV [6] is larger than the systematic uncertainty in this measurement due to both the lifetime of the 2_1^+ state (3 ps uncertainty, corresponding to 0.6 keV uncertainty in energy) and the decay location calculation (5 mm uncertainty, corresponding to 2 keV uncertainty in energy). The apparent discrepancy between the observations may be due to the limited statistics of the elusive 0_2^+ gamma-ray decay in both studies. We adopt the weighted average of 170(2) keV for the energy of the $0_2^+ \rightarrow 2_1^+$ transition. The decay distribution of the 0_2^+ state places an independent 1σ lower limit on the lifetime of 8 ns, confirming the isomeric nature of this state [6]. The upper

limit could not be constrained, however, as shown in the inset of Fig. 2b.

The correlation between the lifetime and partial cross section populating the 0_2^+ state (including feeding from unobserved higher-lying states) can be studied from the yield of the 170-keV peak, shown as a gray band in Fig. 3. Since the gamma-ray efficiency in this measurement strongly depends on the lifetime, the possible cross section can be constrained for a given lifetime or vice versa. For example, if the assumed lifetime is 5 to 10 ns, the gamma-ray efficiency in this setup is maximized so the assumed cross section must be minimized. The total error in this result includes 20% statistical uncertainty in the yield of the 170-keV peak. Another important source of uncertainty is the efficiency of the gates. In order to keep this systematic uncertainty small, the spectrum including fewer gates (Fig. 2b, upper spectrum) was used in this portion of the analysis and contributes 13% relative uncertainty, estimated by applying the same gates to calibration data taken with a ^{152}Eu source. The statistical and systematic uncertainties were combined in quadrature to a total relative uncertainty of 25% in the cross section at any given lifetime.

FIG. 3. Possible values for partial cross section and lifetime of the 0_2^+ state of ^{32}Mg within 1σ are plotted (gray band). From the work of Ref. [38], the cross section upper limit of 0.10 mb is included (red line), and the overlap of that work and this work is highlighted (red hatched region).

The lifetime can be constrained by including the upper limit on the partial cross section of the 0_2^+ state from a previous two-proton removal reaction experiment [38], shown with a red horizontal line in Fig. 3. The previous and present removal reaction experiments populated states in ^{32}Mg using similar mid-target energies of 67 and 75 MeV/nucleon respectively. Eikonal model calculations based on the USDB two-nucleon amplitudes predict 5.11 and 5.23 mb for the inclusive cross section at 67 and 75 MeV/nucleon respectively, suggesting that the results of the previous removal experiment with a slightly different energy can be safely applied here. In the previous measurement, the 0_2^+ state was not observed but the 2_1^+ and 4_1^+ states were observed from their gamma-ray decays. The measured exclusive cross section populating the 2_1^+ and 4_1^+ states accounted for 100(12)% of the ^{32}Mg inclusive cross section, $\sigma_{inc} = 0.76(10)$ mb. This suggests that at most only 12% of the reactions populate the

unobserved 0_2^+ state, corresponding to an upper limit of 0.10 mb for the partial cross section of the 0_2^+ state. With this upper limit in conjunction with the present result, the partial 0_2^+ cross section and lifetime are constrained to $0.03 \text{ mb} < \sigma < 0.10 \text{ mb}$ and $1.5 \text{ ns} < \tau < 38 \text{ ns}$, respectively, as shown by the red hatched region in Fig. 3. The lifetime can be further constrained to $10 \text{ ns} < \tau < 38 \text{ ns}$ from Ref. [6].

Using the 0_2^+ lifetime result of $10 \text{ ns} < \tau < 38 \text{ ns}$ and the weighted average energy of the $0_2^+ \rightarrow 2_1^+$ transition of 170(2) keV, the reduced $E2$ transition probability is $28 \text{ e}^2\text{fm}^4 < B(E2; 2_1^+ \rightarrow 0_2^+) < 122 \text{ e}^2\text{fm}^4$. For physically reasonable values of $\rho^2(E0)$ the $E0$ branch is expected to be less than 1% [6, 39] therefore we assumed the $0_2^+ \rightarrow 0_1^+$ transition to be negligible. Table I summarizes the reduced $E2$ transition probabilities to 0^+ states in ^{32}Mg and neighboring even-even nuclei which can characterize the quadrupole collectivity in these transitions. For ^{30}Mg , a strong $B(E2)$ is observed for the $2_1^+ \rightarrow 0_1^+$ transition [17, 42], whereas in ^{34}Si a strong $B(E2)$ appears for the $2_1^+ \rightarrow 0_2^+$ transition since both the 0_2^+ and 2_1^+ states are considered to be dominated by the collective 2p2h configurations [19]. The present data for ^{32}Mg indicate that the 0_2^+ state is as collective as the 0_1^+ state in contrast with the sizeable difference in transition probabilities between 0^+ states in both ^{30}Mg and ^{34}Si . Notably, the $B(E2; 2_1^+ \rightarrow 0_2^+)$ value in ^{32}Mg is comparable to the strong transitions in ^{30}Mg and ^{34}Si . Indeed, the present result shows $E2$ strength that far exceeds the values calculated by the 3-level mixing model [26] and the SDPF-U-MIX model [19] which both predict $B(E2; 2_1^+ \rightarrow 0_2^+) = 15 \text{ e}^2\text{fm}^4$. The large collectivity in the 0_2^+ state of ^{32}Mg indicates prominent intruder contributions to the state. However, given the large experimental uncertainties, the current $B(E2)$ result does not allow for a stringent conclusion.

Additional direct information about the intruder contributions can be obtained from the partial cross section to populate the 0_2^+ state in comparison with theoretical calculations assuming a pure 0p0h configuration. The ^{34}Si 0_1^+ state is predominantly of 0p0h configuration [14, 17] and the two-proton removal reaction cross section is sensitive to the wave-function overlap between the incoming projectile and outgoing residual nucleus final state, allowing the 0p0h occupancy in the 0_2^+ state of ^{32}Mg to be quantified. Reaction calculations were performed following the method of Ref. [43] which applies

TABLE I. $B(E2)$ values of ^{32}Mg and neighboring even-even isotopes.

$B(E2)$ (e^2fm^4)	^{30}Mg	^{34}Si	^{32}Mg
$2_1^+ \rightarrow 0_1^+$	53(7) [40]	17(7) [41]	94(16) [36]
$2_1^+ \rightarrow 0_2^+$	10.9(12) [40]	61(40) [19]	48^{+74}_{-20} ^a

^a central value of $B(E2) = 48 \text{ e}^2\text{fm}^4$ corresponds to the central lifetime value of 24 ns

the two-neutron amplitudes (TNA) from shell-model calculations combined with eikonal, direct reaction theory. The suppression factor R_{2n} , the ratio of experimental to calculated inclusive two-nucleon removal cross sections, is not well-known for the ($^{34}\text{Si}, ^{32}\text{Mg}$) reaction. We use the value $R_{2n} = 0.5$ seen for a number of less-exotic *sd*-shell nuclei [43]. Assuming the essentially pure 0p0h ^{32}Mg ground state calculated with the USDB interaction [44] is the ^{32}Mg 0_2^+ state, the 0_2^+ cross section is $\sigma_{2n} = 0.42$ mb. This value is significantly larger than the experimental upper limit of 0.10 mb indicating the physical 0_2^+ state has a reduced 0p0h occupancy. The 3-level mixing model predicts a smaller 0p0h occupancy with a probability $\alpha^2 = 0.15$ for the 0_2^+ state [25] due to the sizeable 2p2h and 4p4h contributions to this state which reduces the overlap with the ^{34}Si 0_1^+ state. By scaling the cross section obtained from the USDB pure 0p0h calculation with the 0p0h probability α^2 of the 0_2^+ state, the 3-level mixing model results in a cross section of 0.06 mb. This result is consistent with the measured partial cross section, strongly suggesting that the 0_2^+ state contains strong admixtures of the 2p2h and 4p4h intruder configurations.

The present result raises the question, “Where does the 0p0h-dominant 0^+ state exist in ^{32}Mg , if anywhere?” The 3-level mixing model predicts the 0_3^+ state at 2.22 MeV with the 0p0h probability $\alpha^2 = 0.81$. This gives the partial cross section $\sigma_{2n} = 0.34$ mb for the 0_3^+ state, although associated events have not been experimentally observed. Using the USDB calculations the partial cross sections for the individual states are $\sigma(0^+) = 0.42$ mb, $\sigma(2^+) = 0.94$ mb and $\sigma(4^+) = 1.26$ mb for $R_{2n} = 0.50$. The resulting inclusive cross section of 2.62 mb is much larger than the experimental value of 0.76(10) mb [38]

as is the case for more exotic nuclei in this mass region [45, 46]. This discrepancy indicates either that the R_{2n} is strongly quenched in the ($^{34}\text{Si}, ^{32}\text{Mg}$) reaction where the structure is thought to change drastically between the two nuclei or that the 0p0h components in ^{32}Mg are widely spread or even fragmented above the neutron separation energy ($S_n = 5.778$ MeV), calling for future investigation.

In conclusion, a new method to study isomeric states decaying in-flight was used to observe the $0_2^+ \rightarrow 2_1^+$ transition at 170(2) keV in ^{32}Mg . The $B(E2; 2_1^+ \rightarrow 0_{1,2}^+)$ values of ^{32}Mg reveal that the 0_2^+ state is as collective as the 0_1^+ state which is an unexpected feature not yet observed in neighboring isotopes or available theoretical calculations. From the constrained reaction cross section it is implied that the 0p0h amplitude in the 0_2^+ state is much reduced by the presence of intruder configurations. The novel technique introduced here proved indispensable to observe the 0_2^+ state and, as rare-isotope beams with high velocities continue to be powerful tools, this method will prove vital to extend the sensitive lifetime range of in-beam experiments.

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation (NSF) under Grants No. PHY-1565546, and No. PHY-1811855, by the Department of Energy (DOE) National Nuclear Security Administration through the Nuclear Science and Security Consortium under Award No. DE-NA0003180, and by the Science and Technology Facilities (U.K.) Grant No. ST/L005743/1. GRETINA was funded by the DOE, Office of Science. Operation of the array at NSCL was supported by DOE under Grant No. DE-SC0014537 (NSCL) and No. DE-AC02-05CH11231 (LBNL).

-
- [1] C. Thibault, R. Klapisch, C. Rigaud, A. M. Poskanzer, R. Prieels, L. Lessard, and W. Reisdorf, *Phys. Rev. C* **12**, 2 (1975).
 - [2] E. K. Warburton, J. A. Becker, and B. A. Brown, *Phys. Rev. C* **41**, 3 (1990).
 - [3] B. H. Wildenthal and W. Chung, *Phys. Rev. C* **22**, 5 (1980).
 - [4] A. Navin, D. W. Anthony, T. Aumann, T. Baumann, D. Bazin, Y. Blumenfeld, B. A. Brown, T. Glasmacher, P. G. Hansen, R. W. Ibbotson, P. A. Lofy, V. Maddalena, K. Miller, T. Nakamura, B. V. Pritychenko, B. M. Sherrill, E. Spears, M. Steiner, J. A. Tostevin, J. Yurkon, and A. Wagner, *Phys. Rev. Lett.* **86**, 266 (2000).
 - [5] T. Motobayashi, Y. Ikeda, Y. Ando, K. Ieki, M. Inoue, N. Iwasa, T. Kikuchi, M. Kurokawa, S. Moriya, S. Ogawa, H. Murakami, S. Shimoura, Y. Tanagisawa, T. Nakamura, Y. Watanabe, M. Ishihara, T. Teranishi, H. Okuno, and R. F. Casten, *Phys. Lett. B* **346**, 9 (1995).
 - [6] K. Wimmer, T. Kröll, R. Krücken, V. Bildstein, R. Gernhäuser, B. Bastin, N. Bree, J. Diriken, P. V. Duppen, M. Huuysse, N. Patronis, P. Vermaelen, D. Voulot, J. V. de Walle, F. Wenander, L. M. Fraile, R. Chapman, B. Hadinia, R. Orlandi, J. F. Smith, R. Lutter, P. G. Thirolf, M. Labiche, A. Blazhev, M. Kalkühler, P. Reiter, M. Seidlitz, N. Warr, A. O. Macchiavelli, H. B. Jeppesen, E. Fiori, G. Gerigiev, G. Schrieder, S. D. Gupta, G. L. Bianco, S. Nardelli, J. Butterworth, J. Johansen, and K. Riisager, *Phys. Rev. Lett.* **105**, 252501 (2010).
 - [7] C. Détraz, D. Guillemaud, G. Huber, R. Klapisch, M. Langevin, F. Naulin, C. Thibault, and L. Carraz, *Phys. Rev. C* **19**, 1 (1979).
 - [8] B. Bastin, S. Grévy, D. Sohler, O. Sorlin, Z. Dombrádi, N. L. Achouri, J. C. Angélique, F. Azaiez, D. Baborodin, R. Borcea, C. Bourgeois, A. Buta, A. Bürger, R. Chapman, J. C. Dalouzy, Z. Dlouhy, A. Drouard, Z. Elekes, S. Franchoo, S. Iacob, B. Laurent, M. Lazar, X. Liang, E. Liénard, J. Mrazek, L. Nalpas, F. Negoita, N. A. Orr, Y. Penionzhkevich, Z. Podolyák, F. Pougheon, P. Roussel-Chomaz, M. G. Saint-Laurent, M. Stanoiu, I. Stefan, F. Nowacki, and A. Poves, *Phys. Rev. Lett.* **99**, 022503 (2007).
 - [9] A. Gade, R. V. F. Janssens, T. Baugher, D. Bazin, B. A.

- Brown, M. P. Carpenter, C. J. Chiara, A. N. Deacon, S. J. Freeman, G. F. Grinyer, C. R. Hoffman, B. P. Kay, F. G. Kondev, T. Lauritsen, S. McDaniel, K. Meierbachtol, A. Ratkiewicz, S. R. Stroberg, K. A. Walsh, D. Weisshaar, R. Winkler, and S. Zhu, *Phys. Rev. C* **81**, 051304(R) (2010).
- [10] B. A. Brown, *Physics* **3**, 104 (2010).
- [11] S. M. Lenzi, F. Nowacki, A. Poves, and K. Sieja, *Phys. Rev. C* **82**, 054301 (2010).
- [12] F. Nowacki, A. Poves, E. Caurier, and B. Bounthong, *Phys. Rev. Lett.* **117**, 272501 (2016).
- [13] O. Sorlin and M.-G. Porquet, *Prog. Part. Nucl. Phys.* **61**, 602 (2008).
- [14] E. Caurier, F. Nowacki, and A. Poves, *Phys. Rev. C* **90**, 014302 (2014).
- [15] N. Tsunoda, T. Otsuka, N. Shimizu, M. Hjorth-Jensen, K. Takayanagi, and T. Suzuki, *Phys. Rev. C* **95**, 021304(R) (2017).
- [16] T. Otsuka, T. Suzuki, J. D. Holt, A. Schwenk, and Y. Akaishi, *Phys. Rev. Lett.* **105**, 032501 (2010).
- [17] Y. Utsuno, T. Otsuka, T. Mizusaki, and M. Honma, *Phys. Rev. C* **60**, 054315 (1999).
- [18] B. A. Brown and B. H. Wildenthal, *Ann. Rev. Nucl. Part. Sci.* **38**, 29 (1988).
- [19] F. Rotaru, F. Negoita, S. Grévy, J. Mrazek, S. Lukyanov, F. Nowacki, A. Poves, O. Sorlin, C. Borcea, R. Borcea, A. Buta, L. Cáceres, S. Calinescu, R. Chevrier, Z. Dombrádi, J. M. Daugas, D. Lebhertz, Y. Penionzhovich, C. Petrone, D. Sohler, M. Stanoiu, and J. C. Thomas, *Phys. Rev. Lett.* **109**, 092503 (2012).
- [20] B. V. Pritychenko, T. Glasmacher, P. D. Cottle, M. Fauerback, R. W. Ibbotson, K. W. Kemper, V. Madalena, A. Navin, R. Ronningen, A. Sakharuk, H. Scheit, and V. G. Zelevinsky, *Phys. Lett. B* **461**, 322 (1999).
- [21] V. Chisté, A. Gillibert, A. Leépine-Szily, N. Alamanos, F. Auger, J. Barrette, F. Braga, M. D. Cortina-Gil, Z. Dlouhy, V. Lapoux, M. Lewitowicz, R. Lichtenthäler, R. L. Neto, S. M. Lukyanov, M. MacCormick, F. Marie, W. Mittig, F. de Oliveira Santos, N. A. Orr, A. N. Ostrowski, S. Ottini, A. Pakou, Y. E. Penionzhkevich, P. Roussel-Chomaz, and J. L. Sida, *Phys. Lett. B* **514**, 233 (2001).
- [22] H. Iwasaki, T. Motobayashi, H. Sakurai, K. Yoneda, T. Gomi, N. Aoi, N. Fukuda, Z. Fülöp, U. Futakami, Z. Gacsi, Y. Higurashi, N. Imai, N. Iwasa, T. Kubo, M. Kunibu, M. Kurokawa, Z. Liu, T. Minemura, A. Saito, M. Serata, S. Shimoura, S. Takeuchi, Y. X. Watanabe, K. Yamada, Y. Yanagisawa, and M. Ishihara, *Phys. Lett. B* **522**, 227 (2001).
- [23] J. A. Church, C. M. Campbell, D.-C. Dinca, J. Enders, A. Gade, T. Glasmacher, Z. Hu, R. V. F. Janssens, W. F. Mueller, H. Olliver, B. C. Perry, L. A. Riley, and K. L. Yurkewicz, *Phys. Rev. C* **72**, 054320 (2005).
- [24] K. Li, Y. Ye, T. Motobayashi, H. Scheit, P. Doornenbal, S. Takeuchi, N. Aoi, M. Matsushita, E. Takeshita, D. Pang, and H. Sakurai, *Phys. Rev. C* **92**, 014608 (2015).
- [25] A. O. Macchiavelli, H. L. Crawford, C. M. Campbell, R. M. Clark, M. Cromaz, P. Fallon, M. D. Jones, I. Y. Lee, M. Salathe, B. A. Brown, and A. Poves, *Phys. Rev. C* **94**, 051303(R) (2016).
- [26] A. O. Macchiavelli and H. L. Crawford, *Phys. Scr.* **92**, 064001 (2017).
- [27] B.-M. Sherrill, *Progress of Theoretical Physics Supplement* **146**, 60 (2002).
- [28] D. J. Morrissey, B. M. Sherrill, M. Steiner, A. Stolz, and I. Wiedenhoever, *Nucl. Instrum. Methods Phys. Res., Sect. B* **204**, 90 (2003).
- [29] H. Nishibata, T. Shimoda, A. Odahara, S. Morimoto, S. Kanaya, A. Yagi, H. Kanaoka, M. R. Pearson, C. D. P. Levy, and M. Kimura, *Phys. Lett. B* **767**, 81 (2017).
- [30] D. Bazin, J. A. Caggiano, B. M. Sherrill, J. Yurkon, and A. Zeller, *Nucl. Instrum. Methods Phys. Res., Sect. B* **204**, 629 (2003).
- [31] S. Shimoura, A. Saito, T. Minemura, Y. U. Matsuyama, H. Baba, H. Akiyoshi, N. Aoi, T. Gomi, Y. Higurashi, K. Ieki, N. Imai, N. Iwasa, H. Iwasaki, S. Kanno, S. Kubono, M. Kunibu, S. Michimasa, T. Motobayashi, T. Nakamura, H. Sakurai, M. Serata, E. Takeshita, S. Takeuchi, T. Teranishi, K. Ue, K. Yamada, Y. Yanagisawa, M. Ishihara, and N. Itagaki, *Phys. Lett. B* **560**, 31 (2003).
- [32] D. Weisshaar, D. Bazin, P. C. Bender, C. M. Campbell, F. Recchia, V. Bader, T. Baugher, J. Belarge, M. P. Carpenter, H. L. Crawford, M. Cromaz, B. Elman, P. Fallon, A. Forney, A. Gade, J. Harker, N. Kobayashi, C. Langer, T. Lauritsen, I. Y. Lee, A. Lemasson, B. Longfellow, E. Lunderberg, A. O. Macchiavelli, K. Miki, S. Momiyama, S. Noji, D. C. Radofrd, M. Scott, J. Sethi, S. R. Stroberg, Z. Sullivan, R. Titus, A. Wiens, S. Williams, K. Wimmer, and S. Zhu, *Nucl. Instrum. Methods Phys. Res., Sect. A* **847**, 187 (2017).
- [33] C. Loelius, H. Iwasaki, B. A. Brown, M. Honma, V. M. Bader, T. Baugher, D. Bazin, J. S. Berryman, T. Braunroth, C. M. Campbell, A. Dewald, A. Gade, N. Kobayashi, C. Langer, I. Y. Lee, A. Lemasson, E. Lunderberg, C. Morse, F. Recchia, D. Smalley, S. R. Stroberg, R. Wadsworth, C. Walz, D. Weisshaar, A. Westerberg, K. Whitmore, and K. Wimmer, *Phys. Rev. C* **94**, 024340 (2016).
- [34] S. Agostinelli, J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. Arce, M. Asai, D. Axen, S. Banerjee, G. Barrand, F. Behner, L. Bellagamba, J. Boudreau, L. Broglia, A. Brunengo, H. Burkhardt, S. Chauvie, J. Chuma, R. Chytracek, G. Cooperman, G. Cosmo, P. Degtyarenko, A. Dell'Acqua, G. Depaola, D. Dietrich, R. Enami, A. Feliciello, C. Ferguson, H. Fesefeldt, G. Folger, F. Foppiano, A. Forti, S. Garelli, S. Giani, R. Giannitrapani, D. Gibin, J. J. G. Cadenas, I. Gonzalez, G. G. Abril, G. Greeniaus, W. Greiner, V. Grichine, A. Grossheim, S. Guatelli, P. Gumplinger, R. Hamatsu, K. Hashimoto, H. Hasui, A. Heikkinen, A. Howard, V. Ivanchenko, A. Johnson, F. W. Jones, J. Kallenbach, N. Kanaya, M. Kawabata, Y. Kawabata, M. Kawaguti, S. Kelner, P. Kent, A. Kimura, T. Kodama, R. Kokoulin, M. Kossov, H. Kurashige, E. Lamanna, T. Lampn, V. Lara, V. Lefebure, F. Lei, M. Liendl, W. Lockman, F. Longo, S. Magni, M. Maire, E. Medernach, K. Minamimoto, P. M. de Freitas, Y. Morita, K. Murakami, M. Nagamatu, R. Nartallo, P. Nieminen, T. Nishimura, K. Ohtsubo, M. Okamura, S. O'Neale, Y. Oohata, K. Paech, J. Perl, A. Pfeiffer, M. G. Pia, F. Ranjard, A. Rybin, S. Sadilov, E. D. Salvo, G. Santin, T. Sasaki, N. Savvas, Y. Sawada, S. Scherer, S. Sei, V. Sirotenko, D. Smith, N. Starkov, H. Stoecker, J. Sulkimo, M. Takahata, S. Tanaka, E. Tcherniaev, E. S. Tehrani, M. Tropeano, P. Truscott, H. Uno, L. Urban, P. Urban, M. Verderi, A. Walkden, W. Wander, H. Weber, J. P.

- Wellisch, T. Wenaus, D. C. Williams, D. Wright, T. Yamada, H. Yoshida, and D. Zschesche, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [35] H. Mach, L. M. Fraile, O. Tengblad, R. Boutami, C. Jollet, W. A. Plóciennik, D. T. Yordanov, M. Stanoiu, M. J. G. Borge, P. A. Butler, J. Cedarkäll, P. Fogelberg, H. Fynbo, P. Hoff, A. Jokinen, A. Korgul, U. Köster, W. Kurcewicz, F. Marechal, T. Motobayashi, J. Mrazek, G. Neyens, T. Nilsson, S. Pedersen, A. Poves, B. Rubio, E. Ruchowska, and the ISOLDE Collaboration, Eur. Phys. J. A **s01**, 105 (2005).
- [36] C. Ouellet and B. Singh, Nucl. Data Sheets **112**, 2199 (Data extracted from the ENSDF online database June, 2019) (2011).
- [37] T. Lauritsen, A. Korichi, S. Zhu, A. N. Wilson, D. Weisshaar, J. Dudouet, A. D. Ayangeakaa, M. P. Carpenter, C. M. Campbell, E. Clément, H. L. Crawford, M. Cromaz, P. Fallon, J. P. Greene, R. V. F. Janssens, T. L. Khoo, N. Lalović, I. Y. Lee, A. O. Macchiavelli, R. M. Perez-Vidal, S. Pietri, D. C. Radford, D. Ralet, L. A. Riley, D. Seweryniak, and O. Stezowski, Nucl. Instrum. Methods Phys. Res., Sect. A **836**, 46 (2016).
- [38] D. Bazin, B. A. Brown, C. M. Campbell, J. A. Church, D. C. Dinca, J. Enders, A. Gade, T. Glasmacher, P. G. Hansen, W. F. Mueller, H. Olliver, B. C. Perry, B. M. Sherrill, J. R. Terry, and J. A. Tostevin, Phys. Rev. Lett. **91**, 1 (2003).
- [39] T. Kibédi, T. W. Burrows, M. B. Trzhaskovskaya, P. M. Davidson, and C. W. N. Jr., Nucl. Instrum. Methods Phys. Res., Sect. A **589**, 202 (2008).
- [40] M. S. Basunia, Nucl. Data Sheets **111**, 2331 (Data extracted from the ENSDF online database July, 2019 (2010)).
- [41] R. W. Ibbotson, T. Glasmacher, B. A. Brown, L. Chen, M. J. Chromik, P. D. Cottle, M. Fauerbach, K. W. Kemper, D. J. Morrissey, H. Scheit, and M. Thoennessen, Phys. Rev. Lett. **80**, 10 (1998).
- [42] O. Neidermaier, H. Scheit, V. Bildstein, H. Boie, J. Figging, R. von Hahn, F. Köck, M. Lauer, U. K. Pal, H. Podlech, R. Repnow, D. Schwalm, C. Alvarez, F. Ames, G. Bollen, S. Emhofer, D. Habs, O. Kester, R. Lutter, K. Rudolph, M. Pasini, P. G. Thirolf, B. H. Wolf, J. Eberth, G. Gersch, H. Hess, P. Reiter, O. Thelen, N. Warr, D. Weisshaar, F. Aksouh, P. V. den Bergh, P. V. Duppen, M. Huyse, O. Ivanov, P. Mayet, J. V. de Walle, J. Åystö, P. A. Butler, J. Cederkäll, P. Delahaye, H. O. U. Fynbo, L. M. Fraile, O. Forstner, S. Franchoo, U. Köster, T. Nilsson, M. Oinonen, T. Sieber, F. Wenander, M. Pantea, A. Richter, G. Schreider, H. Simon, T. Behrens, R. Gernhäuser, T. Kröll, Krücken, M. Münch, T. Davison, J. Gerl, G. Huber, A. Hurst, J. Iwanicki, B. Jonson, P. Lieb, L. Liljeby, A. S. nad A. Scherillo, P. Schmidt, and G. Walter, Phys. Rev. Lett. **94**, 172501 (2005).
- [43] J. A. Tostevin and B. A. Brown, Phys. Rev. C **74**, 064604 (2006).
- [44] B. A. Brown and W. A. Richter, Phys. Rev. C **74**, 034315 (2006).
- [45] P. Fallon, E. Rodriguez-Vieitez, A. O. Macchiavelli, A. Gade, J. A. Tostevin, P. Adrich, D. Bazin, M. Bowen, C. M. Campbell, R. M. Clark, J. M. Cook, M. Cromaz, D. C. Dinca, T. Glasmacher, I. Y. Lee, S. McDaniel, W. F. Mueller, S. G. Prussin, A. Ratkiewicz, K. Siwek, J. R. Terry, D. Weisshaar, M. Wiedeking, K. Yoneda, B. A. Brown, T. Otsuka, and Y. Utsuno, Phys. Rev. C **81**, 041302(R) (2010).
- [46] I. Murray, M. MacCormick, D. Bazin, P. Doornenbal, N. Aoi, H. Baba, H. Crawford, P. Fallon, K. Li, J. Lee, M. Matsushita, T. Motobayashi, T. Otsuka, H. Sakurai, H. Scheit, D. Steppenbeck, S. Takeuchi, J. A. Tostevin, N. Tsunoda, Y. Utsuno, H. Wang, and K. Yoneda, Phys. Rev. C **99**, 011302(R) (2019).