Title: Aberrant Ca^{2+} oscillations in smooth muscle cells from overactive human bladders

Authors: Sui G^{1,3}, Fry CH^{1,3}, Malone-Lee J² and Wu C^{2,3}

Institutions where the work was undertaken:

Institute of Urology¹ and Department of Medicine², University College London; Postgraduate Medical School, Faculty of Health and Medical Sciences³, University of Surrey, U.K.

Running title: Smooth muscle Ca^{2+} oscillations in overactive bladders

Key words: human detrusor, spontaneous activity, intracellular Ca^{2+} , overactive bladders, L-type Ca^{2+} channel, T-type Ca^{2+} channel

Correspondence:

Dr C Wu

Postgraduate Medical School

Faculty of Health and Medical Sciences

University of Surrey

Guildford

Surrey GU2 7XH, U.K.

Tel: +44(0)1483682510

e-mail: c.wu@surrey.ac.uk

Abstract

Overactive bladder syndrome (OAB) is highly prevalent and costly, but its pathogenesis remains unclear; in particular, the origin of involuntary detrusor muscle activity. To identify the functional substrate for detrusor muscle overactivity, we examined intracellular Ca^{2+} oscillations in smooth muscle cells from pathologically overactive human bladders. Basal cytoplasmic Ca^{2+} concentration was elevated in smooth muscle cells from overactive bladders. Unprovoked, spontaneous rises of Ca^{2+} were also identified. These spontaneous Ca^{2+} oscillations were Ca^{2+} -dependent, sensitive to L-type Ca^{2+} channel antagonist verapamil and also attenuated by blocking SR Ca^{2+} reuptake. The fraction of spontaneously active cells was higher in cells from overactive bladders and the magnitude of spontaneous Ca^{2+} oscillations also greater. Spontaneous action potentials or depolarising oscillations were also observed, associated with Ca^{2+} rise; with a higher percentage of cells from idiopathic OAB, but not in neurogenic OAB. Low concentrations of NiCl_2 attenuated both spontaneous electrical and Ca^{2+} activation. This study provides first evidence that spontaneous, autonomous cellular activity - Ca^{2+} and membrane potential oscillations, originates from detrusor smooth muscle in human bladders, mediated by extracellular Ca^{2+} influx and intracellular release. Such cellular activity underlies spontaneous muscle contraction and defective Ca^{2+} activation contributes to upregulated contractile activity in overactive bladders.

1. Introduction

Overactive bladder (OAB) is a symptom complex characterised by an increase of the frequency of micturition and feelings of urgency, accompanied sometimes by incontinence[1]. It is a highly prevalent disorder, especially in the elderly and women[2;3], has a profound impact on the quality of life [4;5], and is economically very burdensome to health providers [6]. The symptoms of overactive bladder can be attributed to up-regulated activity of detrusor smooth muscle, and detrusor overactivity is associated with spontaneous, involuntary contractions. The exact pathophysiology remains unclear, although neurological disorders may be causative in a proportion of patients [7;8]. Moreover, functional changes are heterogeneous in isolated smooth muscle from OAB patients [9]. However, a myogenic mechanism, whereby there are fundamental changes to the electromechanical properties of detrusor muscle has been proposed as a cause of detrusor overactivity [10]. Unprovoked, spontaneous muscle activity could be a significant pathophysiological process as it would explain enhanced bladder activity and is more prevalent in tissue from patients with overactive bladders [11], as well as in animal models of detrusor overactivity [12].

However, the cellular basis for enhanced spontaneous activity in detrusor smooth muscle from patients with overactive bladders has not been explored and in particular whether spontaneous activity originates from smooth muscle itself or other cell types [13]. The role of intracellular Ca^{2+} is of particular importance, as it is the main determinant of detrusor contractile activity. The aims of this study were: to test the hypothesis that spontaneous intracellular Ca^{2+} oscillations are increased in human detrusor smooth muscle from overactive bladders; and to elucidate the cellular mechanisms underlying such activity.

2. Materials and methods

2.1. Tissue samples, cell isolation and solutions

Detrusor samples were obtained after approval of the Joint UCL/UCLH Committees on the Ethics of Human Research and with informed patient consent in accordance with the Declaration of Helsinki. Samples were either from control patients with no symptoms of detrusor overactivity - stable bladders, but undergoing open operations for bladder cancer or other disorders; or from patients undergoing clam cystoplasty who had detrusor overactivity, as assessed by urodynamics. For the stable group 48 samples were used (31 male, 17 female, age 58.3 ± 16.4 years, $\pm$ SD), and for the overactive group 66 samples were obtained (21 male, 45 female, age 42.2 ± 16.7 years). Of the overactive group, 38 were idiopathic – bladder overactivity with no clinically detectable neurological disorders (8 male, 30 female, age 49.3 ± 15.1 years), and 28 neuropathic – bladder overactivity with a neurological disorder (13 male, 15 female, age 32.6 ± 13.8 years). Data from the idiopathic and neurogenic OAB groups are compared to control either separately or collectively. Neither gender- nor age-dependent differences were observed for any experimental variables within or between groups.

All samples were treated similarly: they were placed immediately in a Ca-free HEPES-buffered solution after excision and transported to the laboratory within 60 minutes. The mucosa was removed and the underlying muscle tissue was disrupted into isolated cells with a collagenase-containing medium [14] and stored in a Ca-containing HEPES-buffered medium. The yield of cells from samples of both experimental groups was similar, so that the procedure was equally effective in all samples. Viability of the cells was confirmed by trypan blue extrusion and their Ca^{2+} response to stimuli such as carbachol and high-K solutions. The HEPES-buffered solution contained (mM): NaCl, 105.4; NaHCO_3 , 22.3; KCl, 3.6; $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 0.9; $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$, 0.4; HEPES, 19.5; Na pyruvate, 4.5; adjusted to pH 7.1 with NaOH, and with 1.8 CaCl_2 as required. For experiments cells were perfused with Tyrode's solution (mM): NaCl, 118; NaHCO_3 , 24; KCl, 4.0; $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 1.0; $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$, 0.4; CaCl_2 , 1.8; Na pyruvate, 5.0; glucose, 6.1, gassed with 95% O_2 :5% CO_2 to pH 7.35. Ca-free Tyrode's solution (0-Ca solution) had no added CaCl_2 and 0.1 mM EGTA to buffer Ca^{2+} activity at $\text{pCa} > 8$. All chemicals were from Sigma UK.

2.2. Measurement of the intracellular free Ca^{2+} concentration $[\text{Ca}^{2+}]_i$.

[Ca²⁺]_i was measured by epifluorescence microscopy as described previously [15]. Cells were loaded with the Ca²⁺-sensitive fluorochrome Fura-2 AM (Calbiochem, Germany) at 25°C for 30 min and, following Fura-2 AM washout, for 20 minutes, to de-esterify the intracellular fluorochrome to Fura 2 acid, used for experiments. Myocytes were then continuously superfused with normal Tyrode's solution at 37°C, at 2 ml min⁻¹. The cell of interest was illuminated alternately (50 Hz) at 340 or 380 nm, fluorescent light collected between 410 to 510 nm. The ratio, *R*, of fluorescent light when illuminated at 340 and 380 nm was used as an index of the intracellular Ca²⁺ concentration and calibrated as described previously [15].

2.3. Electrophysiological recordings

Electrophysiological recordings used patch-type pipettes (3-5 MΩ) made from borosilicate glass and filled with a high-K based intracellular medium (mM): KCl 20; aspartic acid, 110; MgCl₂, 5.45; Na₂ATP, 5.0; Na₄GTP, 0.2; EGTA, 0.1; HEPES, 5.0; pH adjusted to 7.1 with KOH. Membrane potentials were recorded in current-clamp mode; resting potentials recorded with no current passed into the cells (*I_h* = 0). An Axopatch-1D system (Axon Instruments) was used and data recorded via an A/D converter (Digidata 1200, Axon Instruments) at 4 kHz, and filtered with a low-pass filter, corner frequency 2 kHz.

2.4. Simultaneous recording of [Ca²⁺]_i and membrane potential

To record simultaneously electrical activity and intracellular Ca²⁺ Fura-2 was used in its non-ester form as a pentapotassium salt (Fura-2 K₅, Calbiochem). 100μM Fura-2 K₅ was included in the intracellular medium as above, but with the EGTA concentration reduced to 0.05mM, and dialysed into the cell via the patch pipette[15]. These concentrations of Ca²⁺ buffers contribute about 10% to the total Ca²⁺-buffering power of detrusor smooth muscle cells[16].

2.5. Data analysis and statistical methods

Results are expressed as means±SEM, except where otherwise stated; *n* is the number of preparations. Student's t-test and ANOVA (Bonferroni post hoc test) were used to test the difference between two and multiple mean values respectively, a chi-square test used for two incidences. A Kolmogorov-Smirnov test was used to test the normal distribution of a data set,

137 where $|d|$ is the statistic denoting the maximum difference between two cumulative fraction
138 curves, one for a normal distribution and the other for the experimental data curve with the same
139 mean. The null hypothesis was rejected for $p < 0.05$. A Clampfit program (Axon Instruments)
140 was used for data processing.
141

3. Results

3.1. Resting intracellular $[Ca^{2+}]_i$ in human detrusor cells

With cells from stable bladders, the resting $[Ca^{2+}]_i$ was 66 ± 36 nM (mean $\pm$ SD, $n=196$ cells from 48 patients). The value was not significantly different from a separate study using guinea-pig cells [15]. The variation of data values could be described by a normal distribution ($|d|=0.086$, $p>0.05$). However, the resting $[Ca^{2+}]_i$ in myocytes from overactive bladders was significantly higher (75 ± 47 nM, $p<0.05$, $n=256$ cells from 66 patients). Figure 1 presents data on per patient basis and shows no significant difference in cells from idiopathic and neurogenic overactivity; both data sets were significantly greater than control set.

FIGURE 1 NEAR HERE

3.2. Spontaneous rises of $[Ca^{2+}]_i$

Unprovoked, spontaneous rises of $[Ca^{2+}]_i$ were observed in a proportion of cells from human detrusor samples. This spontaneous activity occurred as: discrete, isolated spike transients; frequent fast, transients; or fused tetanic rises of $[Ca^{2+}]_i$ (Figure 2 A-C). The frequency of these events varied and was on average 3.5 ± 0.3 transients per minute during phases of active firing. The frequency calculated for a longer period of observation (10-15 minutes), to include quiescent periods and thus the overall incidence of the activity, was 0.60 ± 0.10 per minute in cells from stable bladders ($n=50$). The spontaneous Ca-transients were large; the majority were between one-third and two-thirds of those evoked by the muscarinic agonist carbachol and on average were not significantly different in cells from the three groups of biopsy samples. Figure 2D shows a cell in which a maximum carbachol concentration evoked a transient initially, followed by several spontaneous transients of comparable magnitude. In general, the carbachol-transient exhibited an “undershoot” following relaxation [17], whilst the spontaneous transient did not, suggesting a different origin for the Ca^{2+} -transient. The spontaneous transients were associated with cell shortening and indicate their important functional relevance to spontaneous cell contractions (Figure 2E).

FIGURE 2 NEAR HERE

The fraction of spontaneously active cells was greater in OAB compared to control samples (73.7 ± 3.9 and $40.2 \pm 5.2\%$; $n=61, 44$, respectively, $p<0.01$). This difference was maintained if the

two OAB subgroups were compared separately to control (Figure 3A). Furthermore, the frequency of spontaneous activity was greater in cells from OAB bladders; autonomous events, over the long observation period, occurred at a higher frequency of 0.93 ± 0.12 per minute ($n=95$ cells, $p<0.05$) compared to control cells.

The morphology of spontaneous Ca^{2+} -oscillations was variable, due to varying duration and fusion of individual events. Therefore, a standardised index was devised to quantify the overall increase of the $[\text{Ca}^{2+}]$ over base-line – the mean magnitude of $[\text{Ca}^{2+}]_i$ rise, $M\Delta[\text{Ca}^{2+}]_i$, obtained by integrating the area under the $[\text{Ca}^{2+}]_i$ trace over 10-15 minutes, divided by time. This value thus reflects the magnitude, frequency and duration of Ca^{2+} -transients. The value was significantly greater in cells from OAB bladders (13.2 ± 2.0 nM vs 5.7 ± 1.2 nM; $n=40$, 24 respectively, $p<0.01$). The difference was again maintained if the two OAB groups were compared separately to control (Figure 3B). The value of $M\Delta[\text{Ca}^{2+}]_i$ to the maximum carbachol-evoked Ca-transient from the same cell was also significantly greater in the OAB group compared to control ($10.2 \pm 1.6\%$ and $3.6 \pm 0.7\%$; $p<0.01$), again the difference was preserved when each OAB sub-group was compared to control (Figure 3C).

FIGURE 3 NEAR HERE

3.3. The cellular basis of spontaneous Ca^{2+} transients

The magnitude of extracellular entry and intracellular release of Ca^{2+} are sufficient in detrusor muscle to cause significant increases of intracellular Ca^{2+} [15]. The sources of Ca^{2+} that generate spontaneous activity were thus investigated. Ca^{2+} influx through L-type Ca^{2+} channels is the main source of extracellular entry: it can produce large Ca^{2+} transients and is also essential to replenish intracellular Ca^{2+} stores [18]. Figure 4A shows that spontaneous intracellular Ca^{2+} transients were rapidly and completely abolished in a 0-Ca solution, furthermore there was an equally prompt fall of the resting $[\text{Ca}^{2+}]$. Similar observations were made in 12 cells from control and OAB bladders. Similarly, the L-type Ca^{2+} channel blocker verapamil abolished intracellular Ca-transients (Figure 4B, $n=11$). These data suggest the importance of transmembrane Ca^{2+} fluxes through L-type Ca^{2+} channels in generating such activity. Figure 4C implicates also the role of intracellular Ca-stores through the use of cyclopiazonic acid (CPA), which attenuates Ca^{2+} uptake into this compartment. CPA caused a maintained rise of the intracellular $[\text{Ca}^{2+}]$, which returned to control only after its removal: spontaneous activity was

209 suppressed and subsequently returned (n=8). Similar results were observed in OAB and control
210 cells.

211 FIGURE 4 NEAR HERE

213 3.4. Electrophysiological properties of isolated cells

214
215 After membrane break-through, the resting membrane potential showed low-level fluctuations,
216 superimposed sometimes were spontaneous depolarisations in a variety of patterns (Figure 5). In
217 the quiescent phases the averaged potential over several seconds was recorded as the resting
218 membrane potential. The value was similar in cells from the control and overactive groups (-
219 54.0 ± 2.6 mV, $n=82$ cells vs -51.6 ± 2.1 mV, $n=105$ cells, $p>0.05$); there was also no difference
220 between subgroups of OAB (idiopath vs neuropath: -52.3 ± 2.4 mV, $n=81$ cells vs -49 ± 4.5 mV,
221 $n=24$ cells, $p>0.05$). Figure 5A shows examples of spontaneous action potentials, as seen by
222 them overshooting the 0-mV level and 5B is a fast time-base showing upstroke and repolarising
223 phases of an action potential. In other cells smaller, more prolonged depolarisations were
224 recorded (part C), whilst in others slow depolarisations, with occasional superimposed action
225 potentials (part D). It is of note that the spontaneous action potentials and depolarisations were
226 initiated from relatively negative membrane potentials, well below the threshold for the opening
227 of the L-type Ca^{2+} channels at -40 to -35 mV [19] suggesting that other ionic currents contribute
228 to their initiation.

229 FIGURE 5 NEAR HERE

230
231 A higher proportion of cells from overactive bladders displayed spontaneous electrical activity
232 (54.9%, $n=91$ cells vs 29.5%, $n=78$ cells from stable patients, $p<0.01$). This excess of activity
233 was attributable to cells from idiopathic OAB bladder samples (59.2%, $n=71$ cells, $p<0.01$ vs
234 control). With cells from the neuropathic OAB group the incidence of activity was not
235 statistically distinguished from control (40.0%, $n=20$ cells, $p>0.05$ vs control). Thus, although
236 cells from the neuropathic and idiopathic OAB groups showed a similar increased incidence of
237 intracellular Ca-transients, the difference of electrical spontaneous activity suggests that different
238 mechanisms may be responsible in the two pathologies.

240 3.5. The relationship between spontaneous electrical activity and intracellular Ca-transients

The previous measurements suggest some underlying inter-relationship between electrical and Ca events, at least in some pathologies. A causal relationship between these events was thus investigated by simultaneous recording of both variables. Figure 6A shows a single spontaneous action potential associated with a corresponding Ca-transient. Figure 6B exemplifies discrete, more sustained but smaller oscillating depolarisations mirrored by similar changes to the intracellular $[Ca^{2+}]$. In part C a more random pattern of slow depolarisations was again generally accompanied by similar changes to the intracellular $[Ca^{2+}]$. The rise of Ca^{2+} was generally associated with membrane depolarisations greater than 20mV and also related to the duration of the depolarisations.

FIGURE 6 NEAR HERE

To determine the causal relationship, Figure 7A shows an example of a spontaneous depolarisation and rise of intracellular $[Ca^{2+}]$ on a faster time-base. The change of membrane potential proceeded and peaked before the $[Ca^{2+}]$. Such temporal relationship was more clearly demonstrated by the phase plot in Figure 7B in which membrane depolarisation occurred first, followed by the increase of intracellular $[Ca^{2+}]$. The plot also shows that the threshold membrane potential to trigger a detectable Ca^{2+} rise was between -30 to -40 mV, consistent with the L-type Ca^{2+} channel being pivotal in providing Ca^{2+} entry for spontaneous Ca^{2+} oscillations.

FIGURE 7 NEAR HERE

3.6. Involvement of T-type Ca^{2+} channels in spontaneous activity

The above observations suggested that spontaneous electrical activity could be initiated at relatively negative potentials by an inward current separate from the L-type Ca^{2+} channels. T-type Ca^{2+} current which operates at such negative membrane potentials is present in human detrusor muscle and sensitive to low concentrations of Ni^{+} ions [19]. Figure 8A shows a spontaneous action potential in a cell, which fired from a relatively negative potential, and the suppression of such activity by $200\mu M$ $NiCl_2$. Consistently, Figure 8B demonstrates that spontaneous Ca^{2+} oscillations were also suppressed by the same intervention. Similar results were observed in 6 cells. Figure 8C and D suggests that Ni^{+} -sensitive T-type Ca^{2+} current triggered Ca^{2+} -activated BK current at relatively negative membrane potentials. These observations suggest that the T-type Ca^{2+} current is an important contributor in the initiation of spontaneous electrical and hence Ca^{2+} -transients in human detrusor smooth muscle.

FIGURE 8 NEAR HERE

4. Discussion

4.1. The significance of spontaneous activity

A myogenic component to spontaneous activity has been proposed as an important contributor to the overactive bladder as evidenced by: it is increased in OAB both from human and animal models[10;20;21]; present in isolated muscle preparations and whole bladders especially from overactive bladders[22]; modulated by inotropic interventions and insensitive to neurotoxins[23]. Several histological and immunocytochemical studies have examined differences between human detrusor preparations from stable and overactive bladders to identify the underlying pathological process [24]. However, no functional investigations at the cellular level have been carried out to help understand the enhanced spontaneous activity in OAB.

This is the first study to demonstrate that spontaneous Ca^{2+} -transients exist in isolated human detrusor myocytes, and more importantly, occur at a higher frequency (unit or integrated) in cells from overactive bladder biopsies. Furthermore, the resting intracellular $[\text{Ca}^{2+}]$ was greater in overactive bladder myocytes that may facilitate the development of such transients. These differences are unlikely to be due to tissue handling or cell isolation because all tissue samples were handled similarly and the percentage of viable cells similar with all biopsy samples. Individual Ca^{2+} -transients were sufficiently large to elicit contraction. The functional relevance of this spontaneous cellular activity to in situ smooth muscle properties is indicated by the observation of spontaneous Ca^{2+} transients and associated contractile activity in multicellular preparations from both guinea-pig and human tissue [22;25;26] and the fact that such activity is enhanced in specimens from overactive human bladders [11;22]. Thus, increased spontaneous Ca^{2+} -transients may serve as a functional substrate for OAB. The deranged Ca^{2+} regulation may be a common cellular pathway for both OAB modalities and is partly attributable to hyperactive electrical activity. Whether membrane-independent mechanisms are involved in neuropathic bladder overactivity needs further investigation.

The population behaviour of individual smooth muscle cells in the bladder is likely to be more complicated than that of individual cells, as intercellular communications and mediators in extracellular milieu would exert additional influence on their activities. However, the study has revealed fundamental mechanisms underlying such activity that would be difficult to characterize otherwise. Whilst un-coordinated contractile activity of individual cells may

increase tone in the intact muscle bundle, synchronised activation of many cells will produce a bladder contraction. Synchrony may be achieved by intercellular electrical or chemical coupling, and/or stretch-activated channels. The attenuation of OAB contractions by gap junction blockers suggests intercellular co-ordination has a role in such pathological overactivity[27].

Such enhanced muscle tone and contractile activity driven by more spontaneous Ca^{2+} -transients will not just increase contractile function *per se*, but also stimulate sensory fibres in the detrusor layer by mechanical stretch, triggering an enhanced micturition reflex. Studies in cats, for example, have shown that A- δ afferents respond in a graded manner to passive distension and active contraction of the bladder[28].

The origin of the spontaneous activity in detrusor muscle bundles has been debated and non-muscle “pace making cells” have been speculated [13;29]. Recently interstitial cells within the detrusor muscle bundle have been suggested to be a potential candidate[30]. Although these interstitial cells have been extensively characterized by immunocytochemistry [31], functional evidence for pace-making activity and connectivity to smooth muscle cells is still lacking. Contrary to common believe, the present observation has confirmed that smooth muscle cells are capable of generating spontaneous activity. Furthermore, upregulation of spontaneous activity in smooth muscle accounts for the increased contraction in OABs. This not only provides an answer to pathophysiology of detrusor overactivity, but more importantly, a rationale for smooth muscle as the therapeutic target for muscle overactivity.

4.2. The origin of spontaneous Ca^{2+} -transients

Spontaneous Ca^{2+} -transients in guinea-pig detrusor muscle are importantly determined by L-type Ca^{2+} channels[25;26]. A similar involvement was suggested in human myocytes by their sensitivity to removal of extracellular Ca or addition of Ca-antagonists. In addition, Ca^{2+} -transients were associated with action potentials or smaller depolarisations. The time-course analysis indicated that membrane depolarisation preceded the rise of Ca^{2+} , opposite to the case for interstitial cells [32], and again is consistent with membrane events being ultimately responsible for spontaneous Ca^{2+} -transients. A significant observation was that small, but relatively long-lasting depolarisations, were mirrored by changes to intracellular Ca^{2+} . The prolonged depolarisations may be due to a non-inactivating or long-open state of the L-type Ca^{2+} current preparations[33], and is consistent with the efficacy of hyperpolarising agents on

overactive bladders[34] . In addition, Ca^{2+} influx through $\text{Na}^+/\text{Ca}^{2+}$ exchange has been shown in detrusor muscle [16] so that a depolarisation, from whatever source, would also favour Ca^{2+} entry.

An up-regulated T-type Ca^{2+} channel activity, as demonstrated in human detrusor from idiopathic overactive bladders [35], could explain the enhanced electrical activity. T-type Ca^{2+} current is activated at more negative membrane potentials, i.e., near the resting potential recorded in these experiments. Thus, T-type channel activity would depolarize the membrane to a level where L-type channels would open and generate an action potential and enable Ca^{2+} influx. The depressant effect on spontaneous electrical activity of submillimolar NiCl_2 concentrations, fairly selective for T-type Ca^{2+} channels [19], would indicate their importance in initiating spontaneous activity. The ability of the T-type Ca^{2+} channel to trigger Ca^{2+} -activated current suggests its additional role in modulating BK channels of the excitation-relaxation cycle of spontaneous Ca^{2+} oscillations[36].

However, intracellular Ca-stores will also influence spontaneous activity as cyclopiazonic acid (CPA) increased the resting intracellular $[\text{Ca}^{2+}]$ and abolished spontaneous activity. CPA prevents Ca^{2+} uptake into Ca-stores, so that membrane-evoked release is not possible. However, it was not investigated in this study if the size of intracellular Ca-stores is altered in different bladder pathologies.

4.3. Pathophysiological implications and conclusion

Ca^{2+} regulation has been widely recognised to be fundamental in a variety of biological functions[37] and thus deranged Ca^{2+} metabolism would have pathological implications. To this end, Ca^{2+} dysregulation as a pathological basis for muscular dysfunction has been recognised in skeletal muscle and cardiac muscle [38;39] and as such similar mechanisms may mediate smooth muscle disorders. Whilst data from animal models and cell culture strongly suggest a role of intracellular Ca^{2+} in vascular smooth muscle disorders[40;41], results from the present study present the first evidence that Ca^{2+} dysregulation can cause such smooth muscle dysfunction in human. Elucidation of similar mechanisms in other smooth muscle hyperactivity and identification of their modulators may reveal specific drug targets for treatment of the associated human conditions[42].

In summary, the present observations have confirmed that smooth muscle cells are capable of generating spontaneous activity and may thus serve as one origin for spontaneous activity in intact muscle, among other candidates[13;43]. Furthermore, upregulation of spontaneous activity in smooth muscle could account for the increased contraction in OABs. Distinct cellular pathways participate in such spontaneous activation. This not only provides insights to pathophysiology of detrusor overactivity, but also a rationale for smooth muscle as the therapeutic target for muscle overactivity.

386   **Acknowledgements**

387   We thank the Wellcome Trust, St Peter's Trust and The Whittington Hospital NHS Trust for
388   financial support.

389

References

1. Abrams P. Describing bladder storage function: overactive bladder syndrome and detrusor overactivity. *Urology* 2003; **62**: 28-37.
2. Hampel C, Gillitzer R, Pahernik S, Hohenfellner M, Thuroff JW. [Epidemiology and etiology of overactive bladder]. *Urologe A* 2003; **42**: 776-786.
3. Payne CK. Epidemiology, pathophysiology, and evaluation of urinary incontinence and overactive bladder. *Urology* 1998; **51**: 3-10.
4. Abrams P, Kelleher CJ, Kerr LA, Rogers RG. Overactive bladder significantly affects quality of life. *Am J Manag Care* 2000; **6**: S580-S590.
5. Bulmer P, Abrams P. The unstable detrusor. *Urol Int* 2004; **72**: 1-12.
6. Reeves P, Irwin D, Kelleher C et al. The current and future burden and cost of overactive bladder in five European countries. *Eur Urol* 2006; **50**: 1050-1057.
7. de Groat WC. A neurologic basis for the overactive bladder. *Urology* 1997; **50**: 36-52.
8. Fingerman JS, Finkelstein LH. The overactive bladder in multiple sclerosis. *J Am Osteopath Assoc* 2000; **100**: S9-12.
9. Turner WH, Brading AF. Smooth muscle of the bladder in the normal and the diseased state: pathophysiology, diagnosis and treatment. *Pharmacol Ther* 1997; **75**: 77-110.
10. Brading AF. A myogenic basis for the overactive bladder. *Urology* 1997; **50**: 57-67.
11. Mills IW, Greenland JE, McMurray G et al. Studies of the pathophysiology of idiopathic detrusor instability: the physiological properties of the detrusor smooth muscle and its pattern of innervation. *J Urol* 2000; **163**: 646-651.
12. Milicic I, Buckner SA, Daza A et al. Pharmacological characterization of urinary bladder smooth muscle contractility following partial bladder outlet obstruction in pigs. *Eur J Pharmacol* 2006; **532**: 107-114.
13. Brading AF, McCloskey KD. Mechanisms of Disease: specialized interstitial cells of the urinary tract--an assessment of current knowledge. *Nat Clin Pract Urol* 2005; **2**: 546-554.
14. Montgomery BS, Fry CH. The action potential and net membrane currents in isolated human detrusor smooth muscle cells. *J Urol* 1992; **147**: 176-184.
15. Wu C, Fry CH. The effects of extracellular and intracellular pH on intracellular Ca^{2+} regulation in guinea-pig detrusor smooth muscle. *J Physiol* 1998; **508 (Pt 1)**: 131-143.
16. Wu C, Fry CH. $\text{Na}^+/\text{Ca}^{2+}$ exchange and its role in intracellular Ca^{2+} regulation in guinea pig detrusor smooth muscle. *Am J Physiol Cell Physiol* 2001; **280**: C1090-C1096.

17. Baro I, O'Neill SC, Eisner DA. Changes of intracellular $[Ca^{2+}]$ during refilling of sarcoplasmic reticulum in rat ventricular and vascular smooth muscle. *J Physiol* 1993; **465**: 21-41.
18. Wu C, Sui G, Fry CH. The role of the L-type Ca^{2+} channel in refilling functional intracellular Ca^{2+} stores in guinea-pig detrusor smooth muscle. *J Physiol* 2002; **538**: 357-369.
19. Sui GP, Wu C, Fry CH. A description of Ca^{2+} channels in human detrusor smooth muscle. *BJU Int* 2003; **92**: 476-482.
20. Bradshaw HD, Radley SC, Rosario DJ, Chapple CR. Towards a better understanding of involuntary detrusor activity. *BJU Int* 2005; **95**: 799-803.
21. Kato K, Wein AJ, Radzinski C et al. Short term functional effects of bladder outlet obstruction in the cat. *J Urol* 1990; **143**: 1020-1025.
22. Kinder RB, Mundy AR. Pathophysiology of idiopathic detrusor instability and detrusor hyper-reflexia. An in vitro study of human detrusor muscle. *Br J Urol* 1987; **60**: 509-515.
23. Levin RM, Ruggieri MR, Velagapudi S, Gordon D, Altman B, Wein AJ. Relevance of spontaneous activity to urinary bladder function: an in vitro and in vivo study. *J Urol* 1986; **136**: 517-521.
24. Andersson KE. Storage and voiding symptoms: pathophysiologic aspects. *Urology* 2003; **62**: 3-10.
25. Hashitani H, Fukuta H, Takano H, Klemm MF, Suzuki H. Origin and propagation of spontaneous excitation in smooth muscle of the guinea-pig urinary bladder. *J Physiol* 2001; **530**: 273-286.
26. Hashitani H, Brading AF, Suzuki H. Correlation between spontaneous electrical, calcium and mechanical activity in detrusor smooth muscle of the guinea-pig bladder. *Br J Pharmacol* 2004; **141**: 183-193.
27. Ikeda Y, Fry C, Hayashi F, Stolz D, Griffiths D, Kanai A. Role of gap junctions in spontaneous activity of the rat bladder. *Am J Physiol Renal Physiol* 2007; **293**: F1018-F1025.
28. Habler HJ, Janig W, Koltzenburg M. Myelinated primary afferents of the sacral spinal cord responding to slow filling and distension of the cat urinary bladder. *J Physiol* 1993; **463**: 449-460.
29. Brading AF. Spontaneous activity of lower urinary tract smooth muscles: correlation between ion channels and tissue function. *J Physiol* 2006; **570**: 13-22.

30. McCloskey KD, Gurney AM. Kit positive cells in the guinea pig bladder. *J Urol* 2002; **168**: 832-836.
31. Davidson RA, McCloskey KD. Morphology and localization of interstitial cells in the guinea pig bladder: structural relationships with smooth muscle and neurons. *J Urol* 2005; **173**: 1385-1390.
32. Wu C, Sui GP, Fry CH. Purinergic regulation of guinea pig suburothelial myofibroblasts. *J Physiol* 2004; **559**: 231-243.
33. Nakayama S, Brading AF. Possible contribution of long open state to noninactivating Ca^{2+} current in detrusor cells. *Am J Physiol* 1995; **269**: C48-C54.
34. Foster CD, Fujii K, Kingdon J, Brading AF. The effect of cromakalim on the smooth muscle of the guinea-pig urinary bladder. *Br J Pharmacol* 1989; **97**: 281-291.
35. Sui GP, Wu C, Severs N, Newgreen D, Fry CH. The association between T-type Ca^{2+} current and outward current in isolated human detrusor cells from stable and overactive bladders. *BJU Int* 2007; **99**: 436-441.
36. Herrera GM, Heppner TJ, Nelson MT. Regulation of urinary bladder smooth muscle contractions by ryanodine receptors and BK and SK channels. *Am J Physiol Regul Integr Comp Physiol* 2000; **279**: R60-R68.
37. Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* 2003; **4**: 517-529.
38. Avila G. Intracellular Ca^{2+} dynamics in malignant hyperthermia and central core disease: established concepts, new cellular mechanisms involved. *Cell Calcium* 2005; **37**: 121-127.
39. Yano M, Ikeda Y, Matsuzaki M. Altered intracellular Ca^{2+} handling in heart failure. *J Clin Invest* 2005; **115**: 556-564.
40. Ward JP, Snetkov VA, Aaronson PI. Calcium, mitochondria and oxygen sensing in the pulmonary circulation. *Cell Calcium* 2004; **36**: 209-220.
41. Tostes RC, Wilde DW, Bendhack LM, Webb RC. Calcium handling by vascular myocytes in hypertension. *Braz J Med Biol Res* 1997; **30**: 315-323.
42. Lehnart SE, Wehrens XH, Marks AR. Defective ryanodine receptor interdomain interactions may contribute to intracellular Ca^{2+} leak: a novel therapeutic target in heart failure. *Circulation* 2005; **111**: 3342-3346.
43. Brading AF. Spontaneous activity of lower urinary tract smooth muscles: correlation between ion channels and tissue function. *J Physiol* 2006; **570**: 13-22.

Figure legends

Figure 1. Resting intracellular $[Ca^{2+}]$ in human detrusor smooth muscle cells. Mean $\pm$ SEM, $n=48, 38, 28$ patients, for stable, idiopathic and neurogenic groups; * $p<0.05$, vs stable, ANOVA (Bonferroni post hoc test)..

Figure 2. Spontaneous intracellular $[Ca^{2+}]$ transients in human detrusor myocytes. A, discrete transients. B, more rapid transients. C, fused, tetanic responses. D, comparison to maximal carbachol-evoked Ca-transient. E, transmitted-light image showing shortening of detrusor myocyte during a spontaneous rise of intracellular Ca; top: images taken at resting state (a) and during the peak (b) of the spontaneous Ca^{2+} rise, bottom: the time-course of such activity.

Figure 3. Characteristics of spontaneous $[Ca^{2+}]$ transients in smooth muscle cells from stable and overactive bladders. A, the proportion of spontaneously active cells: B, the time-averaged increase of intracellular $[Ca^{2+}]$, $M\Delta[Ca^{2+}]_i$ (see text for details): C, the value of $M\Delta[Ca^{2+}]_i$ normalized to the maximal carbachol-induced intracellular $[Ca^{2+}]$ rise. Data are mean $\pm$ SEM, ANOVA (Bonferroni post hoc test); * $p<0.05$, ** $p<0.01$ vs stable.

Figure 4. Effect of interventions on spontaneous intracellular Ca^{2+} oscillations. A, exposure to zero-Ca solution. B, effect of 20 μ M verapamil. C, effect of 40 μ M cyclopiazonic acid (CPA).

Figure 5. Spontaneous electrical activity in human detrusor myocytes. A, action potentials, either in bursts, as in the start of this trace, or as individual events. B, an action potential from part A on fast time-base. C, low amplitude, more sustained depolarisations. D, slow waves of activity.

Figure 6. The association between spontaneous membrane potential and intracellular $[Ca^{2+}]$ changes. A, single events. B, sustained changes. C, more random changes.

Figure 7. The relationship between spontaneous membrane potential and intracellular Ca^{2+} changes. A, fast time-base of changes to membrane potential and intracellular $[Ca^{2+}]$ during the initial phase of a spontaneous activity. B, phase-plot of the paired event: the arrows indicate the arrow of time.

Figure 8. Effect of 200 μ M $NiCl_2$ on spontaneous activity. A, effect on electrical activity before and after exposure to 200 μ M $NiCl_2$, 2-min interval between traces. B, effect on spontaneous Ca^{2+} rises, the myocyte was exposed to $NiCl_2$ during the time denoted by the horizontal bar. C, Ni^{2+} -sensitive outward current by clamp step to selectively activate T-type current (from -90mV to -50mV). D, Abolition of outward current by iberiotoxin.

Figure 1

Figure 1

Figure 2

Figure 2
A

$[Ca^{2+}]_i$, nM

B

$[Ca^{2+}]_i$, nM

C

$[\text{Ca}^{2+}]_i$, nM

D

E

Figure 3

Figure 3

A

Spontaneous activity
% cells

B

$\Delta M[Ca^{2+}]_i$, nM

C

Figure 4

Figure 4
A

B

c

40 μ M CPA

$[\text{Ca}^{2+}]_i$, nM

200

50

2 min

Figure 5

Figure 5
A

B

C

D

Figure 6

Figure 6

A

B

c

$[\text{Ca}^{2+}]_i$, nM

300
100

----- 0 mV

20
mV

2 min

Figure 7

Figure 7

A

B

Figure 8

Figure 8

A

B

C

D