8074474

UNIVERSITY OF SURREY LIBRARY

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code
Microform Edition © ProQuest LLC.

ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 – 1346

***The regulation of porcine classical and
non-classical MHC class I expression***

by

Laura Tennant

A thesis submitted in partial fulfilment of the requirements
of the University of Surrey for the degree of
Doctor of Philosophy

School of Biomedical and Molecular Science
University of Surrey

The programme of research was carried out at the
Institute for Animal Health, Pirbright

August 2005

Abstract

Cellular responses to viral infection and non-self tissues depend on presentation of antigenic peptides by polymorphic MHC class I molecules. The regulatory mechanisms controlling MHC class I expression are fundamental to effective immune responses, and in the pig are not fully understood. In this study the cellular responses of SLA class I genes to cytokines were studied by measuring SLA class I expression at the cell surface and also by fusing the promoters of the classical genes SLA-1, -2 and -3 and non-classical genes SLA 6 and -7 independently to a luciferase reporter gene. Cell surface expression of SLA class I was measured on Max cells, Shimosuma cells and porcine aortic endothelial cells isolated from inbred pigs of the d/d haplotype and outbred pigs of an unknown haplotype. IFN- α and γ treatment increased SLA class I expression on d/d PAECs and Max cells but not on outbred PAECs or Shimosuma cells. Analysis of SLA class I promoter activity in Max cells showed constitutive activity of SLA-1, -2, -3, -6, -7 and MIC-2 promoters. In summary, classical SLA promoters were responsive to IFNs and co-expression of the transcription factors IRF-1, NF- κ B p65 and CIITA. In contrast to their human counterparts, combined treatment with TNF- α and IFN- α/γ had no synergistic effect on classical SLA class I promoter activity. SLA-1 responded to TNF- α and co-expression of the transcription factors IRF-3, -7 and -9. Non-classical promoters were not induced by IFN- α or γ or CIITA. SLA-7 was responsive to TNF- α and co-expression of IRF-1 but not co-expression of NF- κ B. Interestingly, basal expression of MIC-2 remained unaffected by cytokines or co-expression of transcription factors. These results suggest locus-specific responses of SLA class I genes to cytokines and transcription factors, which can be explained in part by sequence variation in three key SLA class I promoter motifs: ISRE, Enhancer A and SXY. Furthermore, this study has demonstrated that the porcine virus CSFV decreases SLA class I surface expression early during infection and that this effect correlates to decreased constitutive activity of SLA-1, -2 and -7 promoters.

Contents

Abstract	2
Contents	3
List of tables and figures	10
Abbreviations	14
1: Introduction	18
1.1 The major histocompatibility complex	18
1.2 The MHC of swine	18
1.2.1 The SLA class I region	21
1.2.2 Classical (class Ia) genes	21
1.2.2.1 Classical SLA class I gene promoter regions	24
1.2.3 Non-classical (class Ib) genes	25
1.2.3.1 Non-classical SLA class I gene promoter regions	27
1.2.4 MHC class I chain-related (class Ic) genes	27
1.3 Structure and functions of MHC class I proteins	29
1.3.1 Overview of the structure of MHC class I proteins	29
1.3.2 Presentation of peptides to T cells	31
1.3.3 Requirements of peptides for MHC class I presentation	32
1.3.4 The antigen presentation pathway	33
1.3.4.1 Role of the proteasome in cytosolic proteolysis and peptide generation.	33
1.3.4.2 Peptide trimming	36
1.3.4.3 Translocation of peptides into the ER	37

1.3.4.4 Folding of MHC class I heavy and light chains – the peptide loading complex.-----	38
1.3.5 MHC class I molecules are ligands for NK cell receptors -----	39
1.3.6 Regulation of NK cell activity by inhibitory and activating receptors -----	40
1.3.7 Killer Immunoglobulin like receptors (KIR) and MHC class I -----	42
1.3.8 Leucocyte immunoglobulin-like receptors (LIR or ILT)-----	44
1.3.9 Murine Ly49 and MHC class I -----	45
1.3.10 NKG2 family -----	45
1.4 <i>Regulation of MHC class I expression by cytokines and transcription factors.</i> -----	46
1.4.1 Interferon -----	47
1.4.1.1 Induction of type I interferon during viral infections -----	49
1.4.1.2 Induction of IFN- γ -----	53
1.4.1.3 IFNs induce the transcription of ISGs - the 'delayed' response -----	54
1.4.2 The Interferon regulatory factor (IRF) family of transcription factors -----	54
1.4.3 Nuclear Factor kappa B -----	57
1.4.4 The SXY region and the MHC class I enhanceosome-----	59
1.4.5 Summary of Human MHC class I promoter regulation-----	60
1.4.6 Regulation of expression of MHC class I genes of the d/d pig ----	63
1.4.7 Regulation of expression of MHC class I genes of the H01 pig ---	66
1.5 <i>Viruses and MHC class I</i> -----	66
1.6 <i>Classical Swine Fever</i> -----	70

1.6.1 Clinical perspective and causative agent-----	71
Objectives -----	74
Chapter 2. Materials and Methods -----	75
2.1 Reagent suppliers-----	75
2.2 Media, buffers and solutions -----	75
2.2.1 Cell culture media-----	75
2.2.2 Bacterial culture media -----	76
2.2.3 Immunofluorescence microscopy buffers -----	76
2.2.4 FACS Buffers -----	77
2.2.5 DNA manipulation buffers -----	77
Agarose gel DNA electrophoresis buffers -----	77
2.2.6 Plasmid and Bacterial Artificial Chromosome buffers -----	77
DNA preparation buffers-----	77
2.3 Cell lines and viruses-----	78
2.3.1 Cell lines -----	78
2.3.2 Virus and infection protocols-----	79
2.3.2.1 CSFV stock and preparation -----	79
2.3.2.2 Virus infection -----	79
2.4 Antibodies -----	80
2.4.1 Antibodies recognising MHC class I -----	80
2.4.2 Conjugated antibodies and dyes -----	80
2.5 Plasmid DNA tools -----	81
2.5.1 Transfection control plasmids -----	81

2.5.2 Transcription factor activity control plasmids -----	81
2.5.3 Transcription factor expression plasmids-----	82
2.6 <i>DNA manipulation techniques</i> -----	84
2.6.1 Small scale preparation of plasmid DNA (miniprep)-----	84
2.6.2 Large scale preparation of plasmid DNA (maxiprep)-----	84
2.6.3 Purification of DNA fragments from agarose -----	85
2.6.4 Restriction enzymes digestion-----	86
2.6.5 Ligation -----	86
2.6.6 Transformation of competent bacteria -----	86
2.6.7 Generation and small-scale preparation of Bacterial Artificial Chromosomes-----	87
2.7 <i>Production of SLA class I promoter reporter constructs</i> -----	88
2.7.1 Primer design -----	88
2.7.2 Polymerase chain reaction -----	90
2.7.3 Sub-cloning of SLA class I promoters -----	92
2.7.4 Sequencing-----	93
2.7.4.1 Sequencing primers -----	93
2.7.4.2 Sequencing reaction-----	93
2.7.5 Computational analysis of DNA sequence data -----	94
2.8 <i>Cell culture techniques</i> -----	95
2.8.1 Standard transfection protocol -----	95
2.8.2 Co-transfections with transcription factor expression plasmids ---	96
2.8.3 Luciferase and β -galactosidase assays-----	98
2.8.4 Cytokine treatments-----	98
2.9 <i>Indirect Immunohistochemistry</i> -----	99

2.9.1 Indirect immunofluorescence -----	99
2.9.2 Indirect immunofluorescence analysis of MHC class I-----	100
2.9.3 Image analysis-----	101
2.9.4 Analysis of MHC class I surface expression by FACS-----	101
2.10 Statistical analyses-----	102

Chapter 3: Cellular characterisation of SLA class I

expression in porcine cells. -----103

3.1 Introduction-----	103
3.2 Expression of SLA class I on primary cells-----	104
3.2.1 Porcine aortic endothelial cells and responses to interferon -----	104
3.2.2 Primary pig lymphoma cells (Shimozuma cells) and responses to cytokine induction -----	109
3.2.3. Primary Max pig kidney cells and responses to interferon-----	110
3.3 Effect of CSFV infection on MHC class I expression in porcine cells	114
3.3.1 Max cells can be infected with field isolates of Classical Swine Fever Virus. -----	114
3.3.2 CSFV down-regulates constitutive SLA class I surface expression. -----	116
3.4 Discussion-----	122
3.4.1 Summary of results -----	122
3.4.2 General considerations for studying MHC class I -----	123
3.4.2.1 Antibody specificity -----	123
3.4.2.2 Antibody affinity-----	125

3.4.2.3 Tissue specific expression -----	125
3.4.2.4 Non-genetic factors -----	127
3.4.2.5 Summary of the limitations of SLA class I antibody use ----	128
3.4.3 SLA class I expression in primary cells-----	128
3.4.4 Interferon induction of SLA class I expression -----	129
3.4.5 CSFV down-regulates MHC class I expression on porcine cells- -----	131
3.4.6 Summary -----	133

Chapter 4: Physical and Structural Characterisation of the

SLA class I Locus-----134

4.1 Introduction-----	134
4.2 Description of the porcine MHC class I locus and regulatory promoter elements. -----	135
4.3 Cloning of SLA class I promoter regions-----	141
4.4.1 Summary of results-----	144
4.4.2 The sequences of the SLA-2 and SLA-3 promoter differ to the published sequences. -----	146
4.4.3 Functional analysis of SLA class I promoters from the H01 pig -	146

Chapter 5: Transcriptional regulation of classical and non-

classical SLA class I genes.-----148

5.1 Introduction-----	148
5.2 Constitutive activity of classical and non-classical SLA class I promoters in porcine cells-----	148

5.3.1 Regulation by interferons -----	151
5.4 Regulation of SLA promoters by transcription factors -----	159
5.4.1 The interferon-stimulated response element (ISRE) -----	159
5.4.2 Enhancer A region (NF- κ B consensus site) -----	167
5.4.3 S, X and Y elements within SLA class I promoters -----	170
5.5 Viral disruption of SLA class I promoters -----	173
5.5.1 CSFV down-regulates the constitutive activity of SLA class I promoters. -----	173
5.6 Discussion -----	176
5.6.1 Summary of results -----	176
5.6.2 General considerations -----	177
5.6.3 Constitutive activity of SLA promoters in porcine cells -----	178
5.6.4 Interferon induces classical SLA class I promoter activity -----	180
5.6.5 Non-classical promoter activity was not induced by IFNs -----	183
5.6.6 Regulation of SLA class I promoters by Interferon Regulatory Factors -----	184
5.6.7 Transcriptional regulation by TNF- α and NF- κ B -----	187
5.6.8 CIITA induces classical SLA class I promoter activity -----	188
5.6.9 Regulation of MIC-2 expression -----	189
5.6.10 CSFV down-regulates basal SLA class I promoter activity ----	190
5.6.11 Summary and future perspectives -----	191
Chapter 6. Conclusion -----	195
Chapter 7. References -----	201

List of tables and figures

Tables

Table 1. Summary of restriction sites within SLA promoter	89
Table 2. Primers used for PCR amplification of SLA promoters	90
Table 3. Summary of SLA promoter reporter constructs produced.....	92
Table 4. Antibodies and dilutions used in FACS experiments	102
Table 5. CSFV down regulates MHC class I surface expression on Max cells: summary of raw MFI data	117
Table 6. Putative STAT binding motifs in SLA class I promoters	140
Table 7 Anticipated sizes of SLA class I gene promoter regions to be cloned into pGL3-basic	143
Table 8. Summary of the expected results.....	176
Table 9. Summary of the SLA promoter response results... ..	177

Figures

Figure 1.1 Comparative map of the major histocompatibility complex in humans and swine... ..	20
Figure 1.2 A: Schematic of SLA class Ia genes	23
Figure 1.2 B: Schematic of SLA class Ib and Ic genes	23
Figure 1.3: Structure of MHC class I molecules and the peptide binding groove	30
Figure 1.4: Summary of the MHC class I presentation pathway	34
Figure 1.5: Activating and inhibitory signals mediated by ITAM and ITIM motifs upon NK receptor-MHC engagement	41
Figure 1.6: HLA class I alleles are inhibitory ligands for NK killer immunoglobulin receptors (KIRs).....	43
Figure 1.7: Induction of type I IFNs during a viral infection	50
Figure 1.8: Transcriptional regulation of HLA class I genes	61
Figure 1.9: Viruses target the MHC class I antigen presentation pathway in order to avoid immune recognition	67
Figure 1.10: Schematic organisation of the CSFV genome and functions of polyprotein cleavage products	72
Figure 2.1: Protocol for co-transfection of Max cells with multiple plasmids.....	97
Figure 3.1: Comparison of SLA class I expression and IFN responses on PAECs derived from inbred and outbred swine	106
Figure 3.2: Cell surface and intracellular MHC class I expression in PAECs	108
Figure 3.3: MHC class I expression is up regulated by IFN- α and IFN- γ on Max cells.....	111
Figure 3.4: IFN- γ induces cell surface expression of MHC class I	113

Figure 3.5: Max cells can be infected with field strains of CSFV	115
Figure 3.6: CSFV down-regulates MHC class I surface expression on Max cells	117
Figure 3.7: CSFV down regulates MHC class I expression on Max cells	120
Figure 4.1: Physical structure of SLA promoter regions on chromosome 7 and identification of SLA Class I genes from Bacterial Artificial Chromosomes (BACs)	136
Figure 4.2: Comparison of porcine SLA-1 promoter sequence with human HLA-A1, -B7 and -Cw7 alleles	142
Figure 4.3: Analysis of DNA amplified from BACs and cloned into pGL3 basic	143
Figure 4.4: Sequence analysis of SLA promoters	145
Figure 5.1: Constitutive activity of SLA class I promoters in porcine cells	150
Figure 5.2: Effect of interferon stimulation on classical SLA class I promoters	153
Figure 5.3: Effect of interferon stimulation on non-classical SLA class I promoters	156
Figure 5.4: Stimulation of SLA-1 and SLA-7 promoters with TNF- α and co-stimulation with IFN- α /- γ and TNF- α	157
Figure 5.5: Analysis of ISRE sequences from MHC class I promoters	161
Figure 5.6: Effect of IRF-1 and IRF-2 on SLA class I promoters	161
Figure 5.7: Effect of IRF-3 expression on SLA class I promoters	164
Figure 5.8: Effect of IRF-7 expression on SLA class I promoters	164
Figure 5.9: Effect of IRF-9 expression on SLA class I promoters	166
Figure 5.10: Analysis of Enhancer A sequences from MHC class I promoters	168
Figure 5.11: Effect of NF- κ B p65 and p50 on SLA class I promoters	168
Figure 5.12: Analysis of S, X and Y domain sequences from MHC class I promoters	172
Figure 5.13: Effect of CIITA on SLA class I promoters	172
Figure 5.14: CSFV down regulates constitutive classical and non-classical SLA class I promoter activity	175
Figure 5.15: Summary of the transcriptional regulation of SLA class I genes	193

Acknowledgements

First and foremost I would like to thank my supervisor, Dr. Penny Powell her help, guidance, endless enthusiasm and support during this project and of course, for allowing me into her lab in the first place! Special thanks also to Dr. Thomas Wileman for his support and advice. At Surrey University, I would also like to thank my collaborative supervisor Dr. Kikki Bodman-Smith for all of her support and help in preparing the final manuscript (especially with the dreaded stats!).

Across the channel, very special thanks to the members of the LREG lab at INRA in Jouy-en-Josas who I spent three very enjoyable months with. Thank you for giving me the opportunity to collaborate with you, use your resources and for having the patience to listen to me attempt to speak French!! Warmest thanks to Christine Renard and Patric Chardon for their support, critique and for looking after me whilst in a foreign country.

At Pirbright, thank you to everyone who helped me with this project, special thanks to Miriam Windsor and to Haru Takamatsu and Mick Denyer for help with all things related to pigs and FACS !!! it goes without saying that this work would not have been possible without Pirbright's support task force: Sheila and Janet in the library and Wendy and Morgan from computing who have resolved countless computing nightmares for me over the years! I must also say thank you to the laundry staff for providing me with a particularly glamorous wardrobe for the duration of my studies (and the largest pants I have ever seen!). Thank all of my friends at Pirbright for making the last few years such a fun and special time, I have some fantastic memories and miss

you all. In particular I would like to thank my old housemate/drinking buddy/confidante/sounding board Sandra, Volim te! good luck in the USA! Also 'ma poulette' Claire, you've been a great friend and thanks for keeping me motivated for the last few months of writing. On a more serious note, the last year has been a difficult time at IAH and as I final point I would like to wish my friends and colleagues who are moving onto pastures new the very best of luck with whatever, and wherever they have chosen to continue their careers.

In Manchester, thanks to my colleagues at UMIP for being patient and supportive during the final months of preparation of this manuscript.

Special thanks to Anthony for being so understanding, for putting a roof over my head in times of financial crisis and generally for putting up with me (are you insane?), especially during the preparation of this thesis!

Finally, this thesis is dedicated to my parents who gave me the love, support and self-belief that I needed to complete it. Thank you.

Abbreviations

ATF-2	activating transcription factor-2
BAC	Bacterial artificial chromosome
BVDV	Bovine viral diarrhoea virus
cAMP	cyclic adenosine monophosphate
cDNA	complimentary DNA
CIITA	MHC class II transactivator
CREB	cyclic AMP response element binding protein
CSF	Classical swine fever
CSFV	classical swine fever virus
CTL	cytotoxic T lymphocyte
DNA	deoxyribonucleic acid
dsRNA	double stranded ribonucleic acid
DC	Dendritic cells
E.coli	Escherichia coli
eIF2	Eukaryotic initiation factor 2
ER	endoplasmic reticulum
ERAAP	endoplasmic reticulum-associated aminopeptidase P
ERAP1	endoplasmic reticulum-associated aminopeptidase 1
Erp57	endoplasmic reticulum protein 57
FACS	fluorescence-activated cell sorter
FCS	foetal calf serum
GAS	gamma activated sequence
HCMV	human cytomegalovirus
HLA	human leukocyte antigen

HSP	heat shock protein
IFN	interferon
IFN- α	interferon alpha
IFNAR-1	Interferon-alpha receptor 1
IFNAR-2	Interferon-alpha receptor 2
IFN- γ	interferon gamma
IFNGR-1	Interferon-gamma receptor 1
IFNGR-2	Interferon-gamma receptor 2
Ig	Immunoglobulin
IgG	Immunoglobulin G
IgM	Immunoglobulin M
I κ B	Inhibitor of kappa B
IL	Interleukin
ILT2	inhibitory receptor Ig-like transcript 2
ILT4	inhibitory receptor Ig-like transcript 4
IMDM	Iscoves modified Dulbecos medium
IRAK	Interleukin-1 receptor-associated kinase
IRF	Interferon regulatory factor
ISG	Interferon sensitive gene
ISRE	interferon sensitive response element
ITAM	immunoreceptor tyrosine-based activation motif
ITIM	immunoreceptor tyrosine-based inhibitory motif
Jak	Janus activated kinase
κ B	kappa b
Kb	kilobases

kD	kilodaltons
KIR	Killer immunoglobulin-like receptor
KIR2DL	killer immunoglobulin like receptor 2D long
KIR2DS	killer immunoglobulin like receptor 2D short
LB	Luria broth
LIR	Leucocyte immunoglobulin-like receptor
mAB	monoclonal antibody
MFI	Mean fluorescence intensity
MHC	major histocompatibility complex
MIC	MHC-class I chain related
mRNA	messenger ribonucleic acid
NF- κ B	nuclear factor b
NK	natural killer
NF-Y	nuclear factor Y
ORF	open reading frame
PAEC	porcine aortic endothelial cell
PBMC	peripheral blood mononuclear cells
PCR	Polymerase chain reaction
PKR	protein kinase R
PLC	peptide loading complex
RFX	regulatory factor X
RLU	relative light units
RNA	ribonucleic acid
RT	Room temperature
RT-PCR	reverse transcriptase polymerise chain reaction

SLA	swine leukocyte antigen
STAT	signal transducer and activator of transcription
TAP	transporter associated with antigen presentation
TcR	T cell receptor
TGN	trans-golgi network
Th cell	T helper cell
TLR	toll-like receptor
TNF- α	tumour necrosis factor alpha
ULBP	UL16 binding protein

1: Introduction

1.1 The major histocompatibility complex

The major histocompatibility complex (MHC) is a genomic region found in all higher vertebrates studied to date (Trowsdale, 1995). The complex was first described in 1937 following transplantation studies in mice as a genetic region that plays an important role in the acceptance or rejection of grafts between allogeneic individuals (Gorer, 1937; Gorer, 1948). The complex contains an array of genes, many of which are expressed at the cell surface and have immunological functions including a set of highly polymorphic genes that code for two classes of MHC molecules; MHC class I and MHC class II (Horton *et al.*, 2004). In humans these genes are also known as Human Leukocyte antigen (HLA) I and II. MHC class I and II gene products are vital in the immunological dialogue that occurs between T lymphocytes and other cells of the body and have two principal functions. Firstly they present antigenic peptides derived from intracellular pathogens such as viruses, bacteria and parasites to cytotoxic T lymphocytes and secondly they are ligands for natural killer cells.

1.2 The MHC of swine

The porcine equivalent of MHC is known as the Swine Leucocyte Antigen (SLA). It was first described in the seventies, when a correlation was found between acute graft rejection and a group of surface antigens present on peripheral lymphocytes in pigs (Vaiman *et al.*, 1970; Viza *et al.*, 1970). The SLA complex is located on chromosome 7 and is disrupted by the

centromere (Geffrotin *et al.*, 1984; Rabin *et al.*, 1985). The first genetic map of the SLA complex was published by Velten *et al* (1998) and the entire region now has been sequenced. The organisation and content of the SLA complex has been remarkably well conserved during mammalian evolution, emphasising the fundamental importance of this region in immunity. For comparison, the content and organisation of human and swine MHC class I, II and III regions are shown in figure 1.1. In common with human and other animal species, the swine MHC consists of three major clusters or regions of genes: SLA class I, II and III. The class I region contains the classical and non-classical MHC class I genes as well as the MHC class I chain-related (MIC) genes. The class II region contains the MHC class II genes and several other genes that are important in the MHC presentation pathway such as TAP and LMP. The class III region also contains a number of genes with important immunological functions including the cytokines tumour necrosis factor alpha (TNF- α), lymphotoxin a (LTa) and lymphotoxin b (LTb). The most striking difference between pig and human MHC is that the pig MHC complex is much more compact than in human. This is most apparent in the class I region, in humans this region spans 1.8 Mb and contains some 80 genes whereas the swine class I region spans 1.1 Mb and contains 54 genes (Velten *et al.*, 1999). The genes encoded within the SLA class I region are the focus of this study, therefore the current understanding of the organisation, functions and expression patterns of SLA class I genes will now be discussed.

Figure 1.1 Comparative map of the major histocompatibility complex in humans and swine

The genetic map of human MHC is outlined above. The swine MHC, aligned using anchor genes that are perfectly conserved between human and swine, is depicted in red. Swine MHC genes whose positions have been mapped are shown. Human genes with immunological importance are shown in gold, class II sequences in cyan and class Ia genes in red. Porcine genes of immunological importance are typed in bold blue text while the class Ia sequences are in red font. Numbers represent genomic distances in kb. Reproduced with kind permission (Whitelegg, 2001).

1.2.1 The SLA class I region

Our present understanding of the SLA class I region has been gained by traditional serological methods and more recently, by molecular techniques such as the generation of YAC and BAC libraries (Rogel-Gaillard *et al.*, 1997; Rogel-Gaillard *et al.*, 1999) and the identification of YAC and BAC clones spanning the class I region (Velten *et al.*, 1999). Serological analyses of the SLA class I region has identified some 74 haplotypes and three class I series, A, B and C. Whilst the majority of haplotypes express these three series, exceptionally rare haplotypes have also been identified that express more than three SLA class I series and others which express a single locus (Renard *et al.*, 2001). Several pig haplotypes have been studied to date by a number of groups, the most detailed information available is from two haplotypes: H01 and H04 (or d/d). Recently, sequencing of the MHC class I region of the H01 haplotype pig has provided a more detailed understanding of this region with respect to gene content, intron and exon organisation and promoter structure, allowing predictions of expression to be made. Three classes of SLA class I genes have been identified: classical (or class Ia) genes, non-classical (class Ib) and MHC class I chain related (MIC) genes (or class Ic). A systematic nomenclature for the genes and alleles of the SLA based upon the World Health Organisation's nomenclature for human MHC was proposed recently (Smith *et al.*, 2005).

1.2.2 Classical (class Ia) genes

The classical SLA genes are highly polymorphic group of genes whose function is to present antigenic peptides to circulating CD8⁺ T cells. In

humans these genes are HLA-A, -B and -C. In pigs a total of seven classical SLA class I loci have been identified: *SLA-1*, *SLA-2*, *SLA-3*, *SLA-4*, *SLA-5*, *SLA-9* and *SLA-11*, the arrangement of these genes with respect to each other is shown in figure 1.2A. With the exception of *SLA-11*, all of these genes are transcribed in the same direction. Of the class Ia genes, only *SLA-1*, *SLA-2* and *SLA-3* are known to be functional and these genes correspond to the serologically defined SLA C, B and A series respectively. *SLA-1*, -2, -3 and -5 contain the "blueprint" organization of classical MHC class I molecules that is; a leader sequence, three exons encoding the extracellular $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains, a transmembrane region and three intra-cytoplasmic exons. *SLA-1*, -2 and -3 are expressed whereas the expression of *SLA-5* has not been determined. Despite comprising the correct intron/exon organization in order to be functional, the promoter region of *SLA-5* contains mutations that may prevent expression. The coding sequence of *SLA-9* contains a stop codon within exon three (encoding the $\alpha 2$ subunit), which would cause premature termination of transcription should this gene be transcribed. *SLA-11* and *SLA-4* have truncated sequences; *SLA-11* lacks exon 1 and *SLA-4* lacks exons 6 and 8 and encodes only fractions of exons 5 and 7. Therefore *SLA-4*, *SLA-9* and *SLA-11* are considered to be pseudogenes (Chardon *et al.*, 2001).

Overall, the structures of *SLA-1*, *SLA-2* and *SLA-3* genes are very similar and while these genes are ubiquitously expressed, the relative levels of expression of each gene differ. Given that these genes are highly

Figure 1.2 A: Schematic of SLA class Ia genes

The organisation of the seven SLA class Ia genes within the class I region on chromosome 7 of the pig is shown. Genes that are known to be expressed are shown as solid red blocks, genes for which expression is unknown are shown as checkered blocks and pseudogenes are striped boxes. Arrows represent the direction of transcription and the relative distance (in kilobases) between each gene is indicated.

Figure 1.2 B: Schematic of SLA class Ib and Ic genes

The organisation of the non-classical class Ib and Ic genes within the class I region on chromosome 7 is shown. Class Ib genes are depicted as yellow boxes, class Ic as green boxes and the *MIC-1* pseudogene as a checkered boxes. The the direction of gene transcription is indicated by the direction of the arrow point and approximate distances between each gene (in kilobases) are indicated.

homologous, analysis of the promoter regions of these genes may provide an explanation for the variability in expression observed.

1.2.2.1 Classical SLA class I gene promoter regions

The 5' flanking regions of *PD1* and *PD7* genes of the d/d haplotype pig have been examined in detail (Frels *et al.*, 1990, Maguire *et al.*, 1990). These studies permitted the promoter regions of the SLA class I genes *SLA-1*, *-2*, and *-3* genes of the H01 pig to be identified (Renard *et al.*, 2001). In each study a promoter domain of approximately 1 kb was identified upstream of each SLA class I gene. In the H01 pig, classical SLA promoter regions contain a series of conserved, putative regulatory motifs located within distal and proximal promoter domains. The distal domain contains enhancer and silencer sites. The proximal domain contains a region homologous to a functional motif within the mouse H-2k gene, a proximal enhancer, an Enhancer A and an interferon stimulated response element (ISRE). In addition, the proximal domain contains regulatory elements common to MHC class I, class II and β 2-microglobulin promoters, namely three transactivator motifs W, X and Y and a series of proximal or "core" elements including CAT and TATA boxes and a CAP binding motif. The functions of these regulatory motifs and the factors which interact with them have been not been tested experimentally in for the H01 haplotype. However, their human counterparts have been studied in detail, and will be discussed later.

1.2.3 Non-classical (class Ib) genes

A cluster of three non-classical class Ib genes have been identified within the class I region of the H01 haplotype pig: *SLA-6*, *SLA-7* and *SLA-8*. These genes are located adjacent to the junction between the class I and class III regions, approximately 500 kilobases upstream of the class Ia genes. Figure 1.2B shows the organization of these genes. *SLA-6* and -7 are transcribed in the same direction whereas *SLA-8* is encoded on the opposite strand. Six 'alleles' of *SLA-6* have been described which differ from H01 *SLA-6* by a single amino acid and would therefore give rise to a monomorphic product (Chardon *et al.*, 2001). The *SLA-6* gene of the H01 pig shares 99% sequence homology with PD6, a non-classical gene of the NIH d/d mini pig (Chardon *et al.*, 2001). The polymorphisms of *SLA-7* and -8 remain to be established, but it is assumed that they will have few polymorphisms in keeping with the monomorphic characteristic of non-classical genes in other mammals. Overall, the organization of SLA class Ib genes is consistent with the "blueprint" for MHC class I genes, and the products of the three genes are predicted to be membrane-anchored glycoproteins (Chardon *et al.*, 2001). Interestingly, *SLA-6*, -7 and -8 encode seven exons, in contrast to the 8 exons of class Ia genes. The functional implications of these differences remain to be established, as does the association of SLA class Ib with β 2-microglobulin.

Until recently, information regarding the expression of SLA class Ib genes was limited. *SLA-6* is ubiquitously expressed in all tissues and all breeds examined (Ehrlich *et al.*, 1987; Crew *et al.*, 2004). Similarly, Crew *et*

al (2004) reported constitutive expression of *SLA-8* on several tissues with the exception of the brain. *SLA-7* is expressed in a tissue specific manner; transcripts were reported at high levels in cultured swine fibroblasts, at very low levels in the spleen and testes and was not detected in the kidney or PBMCs (Chardon *et al.*, 1999; Crew *et al.*, 2004).

The exact functions of SLA class Ib genes are not well understood. Detailed analysis of coding sequences implies that they encode membrane anchored glycoproteins with the capacity to bind peptides, although the nature of these peptides is unknown (Chardon *et al.*, 2001). The restricted polymorphisms of class Ib genes means that the potential repertoire of peptides that could be presented by these molecules would be limited. There are no direct molecular homologues of non-classical SLA genes in the human or mouse; therefore predictions of functions of these genes cannot be extrapolated from sequence comparisons with human genes alone. However, given the importance of human and mouse class Ib genes in the regulation of NK and antigen-dependant T cell activities, and the relative conservation and/or co-evolution of MHC class I gene function during mammalian evolution, it is predicted that porcine class Ib genes will have functions in parallel with those of humans and mice, that are distinct from class Ia genes. In the human there are three non-classical HLA class I genes: HLA-E, -F and -G. All of these genes encode membrane-anchored glycoproteins and alternative transcripts of HLA-G encode soluble forms of this protein (Lefebvre *et al.*, 2001). HLA-E and -G are capable of binding peptides whereas HLA-F is expressed as an empty heterodimer devoid of

peptide (Wainwright *et al.*, 2000). The most unique features of these molecules are the nature of the peptides which they present, which are primarily derived from the leader sequences of classical HLA class I molecules, their roles as ligands for NK cell receptors and characteristic tissue specific expression patterns.

1.2.3.1 Non-classical SLA class I gene promoter regions

With the exception of SLA-8 which lacks an ISRE, the promoter regions of SLA class Ib genes contain the same regulatory motifs as SLA class Ia genes (Chardon *et al.*, 2001). However, there are some differences in the nucleotide sequences of these motifs which may affect the binding of transcription factors, these are discussed in more detail in results chapter 2.

1.2.4 MHC class I chain-related (class Ic) genes

In the pig, only two MIC-like genes, *MIC-1* and *MIC-2*, have been identified. These genes are located within the SLA class I region in close proximity to the class Ib genes (figure 1.2B, in green). The *MIC-2* gene encodes 6 exons and has a molecular structure compatible with that of a functional gene, whereas *MIC-1* has a truncated sequence and is therefore considered to be a pseudogene (Chardon *et al.*, 2001). Interestingly, the promoter region of *MIC-2* contains a consensus heat shock element, but lacks an ISRE, indicating that the mechanisms regulating *MIC-2* expression are distinct from those that regulate class Ia and Ib genes (Chardon *et al.*, 2001). Currently, knowledge of the transcription, expression and functions of

MIC-2 in the pig is limited, however, studies on human class Ic genes may allow some predictions of the expression patterns and functions of porcine *MIC-2* to be made.

In humans, the *MIC* genes were first described as a group of five genes related to, but highly divergent from class Ia and Ib genes (reviewed by Bahram, 2000). Two of these five genes *MICA* and *MICB* encode functional molecules and contain a genomic structure in parallel with the canonical MHC class I. Studies on human class Ic genes have highlighted four important differences between MHC class Ia and Ib and the *MIC* gene products. Firstly, expression patterns of *MIC* genes are different to class Ia and Ib genes. *MIC* expression is high in mucosal tumors, the epithelia of the subcapsular thymic cortex, endothelial cells and monocytes and relatively low or absent elsewhere (Zwirner *et al.*, 1998). Secondly, *MICA* and *MICB* lack an ISRE and accordingly, expression of *MIC* is not induced by interferons (Holmes *et al.*, 2002). Thirdly *MICA* and *MICB* gene promoters contain a region which shares homology to heat shock protein (HSP) gene promoters which is absent from classical and non-classical HLA class I promoters. The core of this region houses a heat shock response element (HRE) consisting of multiple repeats of the pentanucleotide motif 5'GAAn-3' which is common to a number stress-induced genes including the HSP70 gene (Groh *et al.*, 1996). The function of the HSE of *MICA* and *MICB* was demonstrated by heat shocking cells; incubation of cells at 42 - 42.5°C results in an increase in *MIC* expression at the mRNA level, suggesting that these genes play a role in the cellular response to stress (Groh *et al.*, 1996).

Another characteristic which sets MIC proteins apart from other class I proteins is that MIC-A and -B do not bind peptides or require β 2-microglobulin for correct folding, stability or expression at the cell surface. Therefore these genes do not play a role in antigen presentation to CD8⁺ T cells (Groh *et al.*, 2001). Instead they are ligands for the NKG2D receptor expressed on NK and $\gamma\delta$ T-cells. Engagement of MIC with the NKG2D receptor induces cytotoxic killing of cells expressing MIC and therefore increased MIC expression cells provides a stress signal to the immune system that is distinct to that of MHC class I and II molecules and is important in innate immunity for surveillance of virus infected cells or cells undergoing tumourogenesis. For example, Groh et al (2001) demonstrated that cell surface expression of MIC increases in cells infected with cytomegalovirus (CMV), which induced NK mediated lysis of infected cells.

1.3 Structure and functions of MHC class I proteins

1.3.1 Overview of the structure of MHC class I proteins

The MHC class I molecule is a type 1 transmembrane protein composed of two polypeptide chains; a 44-49 kDa heavy chain non-covalently associated with a 12 kDa light chain, β 2-microglobulin (Fruh & Yang, 1999). In humans, mice and pigs the heavy chain is encoded by the highly polymorphic MHC class I genes whereas the β 2-microglobulin gene is not MHC-encoded. The heavy chain polypeptide is approximately 350 amino acids in length and folds into three distinct extracellular domains α 1, α 2 and α 3 plus a transmembrane and cytoplasmic region (Bjorkman *et al.*, 1987),

Figure 1.3: Structure of MHC class I molecules and the peptide binding groove.

Figure A, shows a cartoon of MHC class I at the cell surface. The $\alpha 1$ and $\alpha 2$ domains of MHC fold together to form a peptide binding groove (peptide shown in red). B shows the structure of the peptide binding groove in detail; the structure comprises two segmented α helices resting on 8 anti-parallel β strands forming a cleft or groove. A peptide bound to the groove is shown in figure C (peptide in green) (Bjorkman *et al.*, 1987).

which associate with β 2-microglobulin (figure 1.3 A & B). The α 1 and α 2 domains fold together into a single structure comprising two segmented α helices resting on 8 antiparallel β strands. This results in the formation of a cleft or groove, the peptide-binding region of MHC (figure 1.3 C).

The α 1 and α 2 domains of the MHC protein contain the most variation in amino acid residues. Variation is concentrated in the amino acids forming the extremities of the groove region and the point of contact between MHC and the peptide. These variations dictate the nature of the peptide that can be bound. By contrast, the α 3 region is highly conserved between MHC molecules, this is important as this region contains a projection that interacts with the cytotoxic T cell CD8 co-receptor (Salter *et al.*, 1989). The light chain, β 2-microglobulin, is also highly conserved within the individual and between species. This region is not directly involved in peptide binding and presentation but is essential for maintaining the structural integrity of the MHC class I molecule.

1.3.2 Presentation of peptides to T cells

As discussed above, MHC class I molecules bind peptides that are derived from cytosolic proteins and display them at the cell surface. These peptides can be derived from endogenous ("self") or exogenous ("foreign") sources such as viruses, bacteria and parasites. Exogenous peptides are recognised as foreign by CD8⁺ T lymphocytes that initiate a specific immune response that ultimately results in killing of the infected cell, containment of

infection and generation of immunological memory. In comparison MHC class II proteins present to CD4⁺ T lymphocytes which are predominantly T helper cells.

The molecular structure of the MHC-antigen complex is arranged such that the T cell receptor (TcR) is in contact with both the peptide and some of the polymorphic residues of the MHC class I molecule. T cell activation requires recognition of both the antigen and the MHC molecule and is therefore said to be 'MHC restricted.' In order to prevent cytotoxic killing of healthy cells, it is essential that T cells with specificity for 'self' antigens are eliminated from the T cell repertoire. To prevent this, negative selection by endogenous antigen occurs in the thymus.

1.3.3 Requirements of peptides for MHC class I presentation

Efficient peptide presentation by MHC class I depends on a number of factors (reviewed in Palmer & Cresswell, 1998). Firstly, antigenic proteins must be degraded into peptides of an appropriate length for incorporation into the binding groove; 8 to 10 amino acids long. Secondly, peptides must contain appropriate amino acid motifs permissible for incorporation into the peptide-binding groove of polymorphic MHC molecules. Thirdly, peptide generation must be rapid, this is important during viral infections, where viral replication may be rapid. In this instance it is essential that peptides are presented to the immune system quickly, so that the spread of viral infection can be controlled. Some viruses replicate slowly and the amount of viral proteins available for degradation may be limited. In such cases protein

degradation must be highly efficient, in order to capitalise on the proteins available for presentation to cytotoxic T lymphocytes (CTLs). Most importantly, the whole process of antigen presentation and the gene products involved need to be inducible, so that under appropriate circumstances such as a viral infection, antigen presentation to T cells can be maximised.

1.3.4 The antigen presentation pathway

Intracellular antigen processing pathways determine which peptides are available for presentation on MHC class I and are therefore important for shaping the immune response. The antigen processing pathway involves multiple steps: proteasomal cleavage, transport into the ER, binding to MHC molecules and transport of MHC class I/peptide complexes to the cell surface (summarised in 1.4 and discussed in detail below). Each step of the pathway is a potential target for intervention by viruses. This will be discussed later.

1.3.4.1 Role of the proteasome in cytosolic proteolysis and peptide generation.

The proteasome is a barrel shaped cytoplasmic and nuclear organelle specialised for the elimination of cellular proteins marked for degradation and synthesis of peptides for presentation on MHC class I (Figure 1.4 A). Impaired proteasomal function causes a dramatic decrease in MHC class I peptide presentation at the cell surface, underlining the important role of this organelle in antigen presentation (Kessler *et al.*, 2003). The proteasome is made up of a proteolytic chamber, the 20S core subunit, capped at each end by a regulatory 19S subunit. The 20S core comprises four stacked rings,

Figure 1.4: Summary of the MHC class I presentation pathway.

Proteins are degraded by the proteasome into peptides of various lengths (A). Most importantly for MHC class I presentation, peptides of 9-12 amino acids in length are produced. These peptides are translocated into the ER lumen by a specialised pump, the transporter associated with antigen presentation (TAP) (B). Newly synthesised MHC class I heavy chains are also transported into the ER lumen where they bind to calnexin, β 2-microglobulin and ERp57. Calnexin disassociates from the complex and is substituted by calreticulin and together these proteins are known as the peptide loading complex (C). The peptide loading complex associates with TAP and peptides are loaded into the binding groove of the heavy chain, an interaction that requires the ER chaperone tapasin (D). The MHC class I is now fully assembled. MHC class I leave the ER and are transported via the trans-golgi network to the cell surface where they are displayed to circulating CD8⁺ T cells (E). Adapted from Palmer & Cresswell (1998).

each containing seven subunits. Active sites for proteolysis are located on the internal surfaces of the two innermost rings of the chamber and preferentially bind acidic, basic or hydrophilic residues within target proteins. The 19S subunits regulate proteolysis by guarding the entrance of the proteasome and preventing non-specific degradation of cytosolic proteins. They are composed of a ring of α subunits forming an axial channel which is able to control the influx and efflux of proteins by regulating the opening and closing of the entrance of the proteolytic chambers. The 19S subunits also contain six ATPases which unravel incoming proteins and feed them into the 20S core (Braun *et al.*, 1999). The speed and efficiency of peptide generation must be increased during viral infections, this is mediated by interferon gamma which induces the expression of genes encoding three additional proteasome subunits LMP-2, LMP-7 and MECL-1. These subunits are incorporated into the proteasome generating an "immunoproteasome" with enhanced peptide production capability (Aki *et al.*, 1994).

In order to prevent non-specific degradation of cellular proteins, several mechanisms exist that regulate degradation and targeting of proteins to the proteasome. Most commonly, proteins destined for destruction are tagged with ubiquitin, this involves conjugation of ubiquitin to lysine residues within the target protein sequence. Alternatively, the inherent nature of a protein can predispose it to rapid degradation; proteins containing regions rich in proline (P), glutamic acid (E), serine (S) and threonine (PEST sequences) are rapidly degraded and have shorter half lives than proteins lacking such sequences (Gromme & Neefjes, 2002). In addition, some

proteins contain 'destruction boxes' these are sequences that dictate degradation at specific points in the cell cycle, cyclin is the classic example of such a protein (Bastians *et al.*, 1999). The identity of the N-terminal amino acid of a protein also determines the rate of degradation. Proteins with N-terminal arginine are destabilised and rapidly degraded whereas methionine at this position stabilises proteins, this phenomenon is referred to as the N-end rule (Stryer, 2001).

As well as presenting peptides derived from proteins targeted for degradation, MHC class I presents peptides derived from metabolically stable proteins as well as peptides from proteins which are contained within sub-cellular compartments. The mechanism by which the proteasome has access to such proteins could not be explained until Schubert *et al.* (2000) proposed that peptides from metabolically stable or compartmentalised proteins are actually derived from defective ribosomal products (DriPs) produced during translation. Polyribosomes are fairly inefficient and frequently make translational errors resulting in the synthesis of proteins that contain errors or are incorrectly folded. Erroneous proteins are targeted to the proteasome for degradation and misfolded proteins are retrotranslocated from the ER through the translocon into the cytoplasm for degradation into peptides (Huppa & Ploegh, 1997).

1.3.4.2 Peptide trimming

Despite the proteasome being specialised for protein degradation it transpires that the generation of peptides compatible with MHC class I

binding is a relatively inefficient process. The proteasome produces peptide products that range between 3 and 22 amino acids in length, with around two thirds of these products being too short for MHC binding and a large proportion requiring further processing before being compatible with the MHC binding groove (Kisselev *et al.*, 1999). Peptides that are longer than 15 amino acids in length require trimming, this is carried out by cytoplasmic aminopeptidase tripeptidyl peptidase II (TPPII) and subsequently by cytoplasmic aminopeptidases such as leucine aminopeptidase (LAP) and thimet oligopeptidase (TOP) (Reits *et al.*, 2004). Some oversized peptides are trimmed in the ER by the ER-associated aminopeptidases ERAAP and ERAP1 (Saric *et al.*, 2002; Serwold *et al.*, 2002). Most peptides will be completely degraded into single amino acids by these peptidases, however, those which escape complete degradation may enter the MHC class I presentation pathway. In common with proteosomal degradation, the speed of digestion by aminopeptidases is increased by interferon (Reits *et al.*, 2004).

1.3.4.3 Translocation of peptides into the ER

Peptides must be transported into the lumen of the endoplasmic reticulum (ER) for loading onto MHC class I molecules. Targeting of proteins to the ER normally requires signal sequences however, in the case of peptides, this process is mediated by a specialised pump; the transporter associated with antigen processing (TAP) (figure 1.4 B). The TAP pump is a heterodimer of TAP-1 and TAP-2 proteins integrated into the luminal membrane of the ER which fold together to form an extra-luminal peptide

binding domain and a pore through which peptides translocate in an ATP-dependant manner (Reits *et al.*, 2000). In common with the proteosomal proteins LMP-2 and -7, the genes that encode TAP-1 and -2 are encoded within the MHC of humans and pigs and their expression is induced by interferons (Fisk *et al.*, 1994).

1.3.4.4 Folding of MHC class I heavy and light chains – the peptide loading complex.

Correct assembly of MHC class I molecules and binding of peptides requires the co-operative action of several accessory molecules and chaperones collectively known as the peptide loading complex (PLC) which includes; TAP1 and TAP2, tapasin, calreticulin and ERp57. Newly synthesised MHC class I heavy chains are transported into the ER where they bind calnexin. Subsequent binding of β 2-microglobulin is simultaneous with release of the MHC class I heavy chain, substitution of calnexin with calreticulin and recruitment of ERp57 (Lehner & Cresswell, 2004) (figure 1.4 C). Together calreticulin and ERp57 are an ER quality control mechanism that regulates correct folding of class I glycoproteins and ensures that incorrectly folded complexes are not transported to the cell surface. Calreticulin monitors the glycosylation status of MHC class I ensuring that immature heavy chains are not incorporated into the PLC. Finally, the ER chaperone tapasin binds to the α 2 and α 3 domains of MHC and to the TAP transporter, this interaction enables peptides to be loaded onto class I heavy chains and signifies the completion of MHC class I assembly (Sadasivan *et al.*, 1996). MHC class I proteins are then released from the PLC, exit the ER

and are transported, *via* the golgi, to the cell surface where they are displayed to CD8⁺ T cells (figure 1.4 E).

1.3.5 MHC class I molecules are ligands for NK cell receptors

In addition to recognition by CTLs, MHC class I proteins are recognised by Natural Killer (NK) cells and play an important role in innate immunity. NK cells are a heterogeneous population of bone marrow derived lymphocytes that are instrumental in the innate response against pathogens. They were first described as cells which could specifically recognise and kill tumour cells (Herberman *et al.*, 1975). We now know that their primary functions are (i) cytotoxicity; to seek out and destroy virally infected or tumour cells and (ii) cytokine production; to secrete IFN- γ .

NK cells recognise aberrant MHC class I expression. This is vital for recognition of tumour cells and during viral infections when class I expression may be down regulated, a phenomenon known as "missing self" (Ljunggren & Karre, 2002). It is important to note that NK cell killing is not by default when MHC expression is absent, since erythrocytes that do not express MHC are not targeted. A more appropriate description for targets of NK cytotoxicity is "altered" or "stressed self." The importance of NK cell recognition of MHC is reflected by the identification of several classes of genes encoding MHC receptors in several mammalian species. Interestingly, the function of MHC recognition by NK receptors appears to have evolved independently in different mammalian species; comparative studies of NK

genes sharing similar functions in different species are too distantly related to have a common ancestor (Parham, 2005).

1.3.6 Regulation of NK cell activity by inhibitory and activating receptors

NK cell activity is tightly regulated by the interactions of inhibitory and activating receptors on the surface of NK cells with receptors expressed on target cells, including two families of receptors that interact with MHC. The first are the Killer Immunoglobulin like receptors (KIR) in humans (section 1.3.7) and c-type lectin like receptors (Ly49) in rodents (section 1.3.8). The second are heterodimers of CD94 and members of the NK group 2A family (section 1.3.9). Each family includes activating and inhibitory forms of receptor, dictated by the presence of immunoreceptor tyrosine-based activation or inhibitory motifs "ITAM" or "ITIM" motifs in the cytoplasmic regions. ITAM and ITIM motifs mediate their effects *via* distinct biochemical pathways, summarised in figure 1.5. Inhibitory receptors such as KIR2DL contain one or more ITIM motif [(Ile/Val/Leu/Ser)-X-Tyr-XX-(Leu/Val) where x denotes any amino acid] (Lanier, 2003). Upon engagement of an inhibitory receptor with an MHC ligand the ITIM motif is phosphorylated and Src-homology (SH) containing phosphatases are recruited. These phosphatases inhibit NK cell activation by counteracting 'activation' signals, the exact mechanisms that mediate this effect are not fully understood. It is important to note that inhibitory receptors do not globally repress NK cell activation, rather it is the balance between activating and inhibitory signals that dictate the nature of the response generated.

Figure 1.5: Activating and inhibitory signals mediated by ITAM and ITIM motifs upon NK receptor-MHC engagement.

This figure shows NK cell receptors that interact with MHC class I (blue), their associated adaptor molecules (green) and the mechanism by which they initiate and inhibitory or activating signal. Inhibitory receptors contain immunoreceptor tyrosine-based inhibition motifs (ITIM) in the cytoplasmic domain [(Ile/Val/Leu/Ser)-X-Tyr-XX-(Leu/Val) where x denotes any amino acid, shown in blue]. Upon engagement of an MHC class I ligand, ITIM motifs are phosphorylated by Src-homology containing phosphatases (SHP), resulting in an inhibitory signal (left panel). The DAP adaptor molecules associated with activating receptors contain immunoreceptor tyrosine-based activation motifs (ITAM) [YxxL/Ix6-8YxxL/I or YxxM (where x is any amino acid), shown in red]]. The activating receptors KIR2DS, NKG2C, -E and -H (centre panel) associate with DAP12 (centre panel). Upon engagement with MHC class I ligands, protein tyrosine kinases of the Src family are activated and phosphorylate tyrosine residues of the DAP12 ITAM leading to recruitment and activation of the tyrosine kinases Zap70 or Syk. NKG2D associates with DAP10 (right panel). Interaction of this receptor with class Ic ligands results in phosphorylation of the DAP10 ITAM by phosphatidylinositol-3 kinase (PI3 kinase) and activation of NK cell activity. Adapted from Lanier, 2003.

In the case of activating receptors, ITAM motifs contain at least one activation motif (YxxL/Ix₆₋₈YxxL/I where x is any amino acid) with one exception, the adaptor molecule DAP10, which contains an YxxM motif. Engagement of activating receptors with their corresponding ligands activates protein tyrosine kinases of the Src family which phosphorylate ITAMs on tandem tyrosine residues. This leads to the recruitment and activation of the tyrosine kinases Zap70 and Syk. In the case of NKG2D (section 1.3.10), phosphorylation of the ITAM motif of the adapter molecule DAP10 causes recruitment and activation of phosphatidylinositol-3 kinase (PI3 kinase) (Lanier, 2003). In addition, an alternative (short form) transcript of NKG2D is expressed in activated NK cells which can associate with DAP10 and DAP12 (Diefenbach *et al.*, 2002). This means that under appropriate circumstances, NKG2D can stimulate both signal transduction pathways representing both stimulation and co-stimulation by the same receptor.

1.3.7 Killer Immunoglobulin like receptors (KIR) and MHC class I

KIRs are members of the immunoglobulin (Ig) superfamily, expressed on NK cells and a subset of T cells. The KIR receptors vary in the number of extracellular Ig domains (two or three) and the length of their cytoplasmic tails (long or short). Long cytoplasmic domains contain ITIMs and are inhibitory receptors; short cytoplasmic domains lack ITIMs and are activating receptors. KIRs interact with the $\alpha 1$ and $\alpha 2$ domains of a number of classical and non-classical HLA class I molecules. The interactions between KIR receptors and their respective class I ligands are summarised in figure 1.6.

Figure 1.6: HLA class I alleles are inhibitory ligands for NK killer immunoglobulin receptors (KIRs).

In the human, classical and non-classical MHC class I are recognised by specific members of the killer immunoglobulin-like (KIR) family of receptors. KIRs contain up to three extracellular domains (blue circles) and four immunoreceptor tyrosine-based activation motifs (red squares). This figure shows individual KIR receptors and their HLA class I ligand. Engagement of KIR/HLA provides an inhibitory signal which blocks NK cytotoxicity (Moretta *et al.*, 1996).

KIR3DL2 recognises HLA-A3, KIR3DL1 recognises HLA-Bw4 and the KIR2DL subfamily recognises distinct groups of HLA-C alleles (Dohring *et al.*, 1996, Gumperz *et al.*, 1995, Moretta *et al.*, 1996). In addition, KIR2DL4 has been shown to bind HLA-G, although there is conflicting data suggesting that HLA-G is the ligand for ILT2 and ILT4 rather than KIR2DL4 (Allan *et al.*, 1999; Navarro *et al.*, 1999, Rajagopalan & Long, 1999).

1.3.8 Leucocyte immunoglobulin-like receptors (LIR or ILT)

The LIR are a family of 13 Immunoglobulin-like receptors. Predominantly expressed on B and T cells, some members of this group are expressed on NK cells. LIR-1 has four extracellular domains and four ITIMs within the cytoplasmic domain and is a receptor for classical HLA molecules as well as the non-classical molecule HLA-G (Gonen-Gross *et al.*, 2003). LIR-1 recognises the highly conserved $\alpha 3$ domain of HLA-class I which may explain its broad reactivity for classical and non-classical HLA class I. LIRs also recognise UL16 binding proteins (ULBP) and the human cytomegalovirus virus MHC class I homologue UL18 (reviewed by Radaev *et al.*, 2002). Interestingly LIR-1 has a higher affinity for UL18 than for HLA class I and since CMV down-regulates MHC class I expression, UL18 may be important for avoiding NK cell recognition of virus infected cells (Chapman *et al.*, 1999).

1.3.9 Murine Ly49 and MHC class I

Ly49 are a polymorphic family of 9 inhibitory receptor genes (LyA-I) encoded on the mouse NK complex. Ly49 homologues have been identified in the rat (Dissen *et al.*, 1996), and in the pig (Gagnier *et al.*, 2003) but are absent in humans. These receptors are type II transmembrane glycoproteins and are expressed as homodimers at the cell surface. In common with other members of the c-type lectin superfamily, the extracellular region of Ly49 contains carbohydrate recognition domains and it is through these domains that Ly49 recognise the $\alpha 1$ and $\alpha 2$ domains of murine MHC class I molecules. This interaction is dependent upon association of $\beta 2$ -microglobulin (Lanier, 1998).

1.3.10 NKG2 family

The NKG2 family (NKG2A-F) are c-type lectins encoded on chromosome 12 of the human and 6 of the mouse. The family includes both activating and inhibitory receptors. NKG2 are expressed at the cell surface as heterodimers with the invariant chain CD94, with the exception of NKG2D that is expressed as a homodimer in association with the adaptor protein DAP10. NKG2A, -B and -C are inhibitory receptors that recognise the non-classical MHC molecule HLA-E (Posch *et al.*, 1998). In the mouse NKG2A recognises the non-classical MHC class I Qa-1 (Vance *et al.*, 1998). HLA-E and Qa1 are unusual in that they bind and present signal peptides derived from the leader sequences of MHC class I heavy chains, therefore monitoring these MHC molecules is used to measure "normal" class I expression.

NKG2D is a divergent member of the c-type lectin-like family of NK receptors and shares only 20 to 30 % homology with other members of the NKG2 family. It is an activating receptor expressed on NK cells, CD8⁺ $\alpha\beta$ T cells and CD8⁺ $\gamma\delta$ T cells as a homodimer in association with the adaptor molecule DAP10. The ligands for NKG2D in the human are the stress-inducible MHC class I chain related proteins MICA and MICB and the Cytomegalovirus encoded UL16 binding proteins (ULBP) (Bauer *et al.*, 1999; Cosman *et al.*, 2001). In the mouse, ligands for NKG2D are the retinoic acid early inducible gene (Rae1) and H60 (Cerwenka *et al.*, 2000; Diefenbach *et al.*, 2000). Expression of MIC, Rae-1 and H60 is usually restricted to tumour cells, suggesting a specialised role for NKG2D in tumour recognition. Recently, a single porcine UL16 binding protein (PULBP) was identified that, with the exception of the brain, is ubiquitously expressed (Garcia-Borges *et al.*, 2005). Increased expression of PULBP is induced by retinoic acid and heat stress. PULBP does not bind human NKG2D, presumably because of the cross species difference between pig and human NKG2D, and its interaction with porcine NKG2D has not been confirmed.

1.4 Regulation of MHC class I expression by cytokines and transcription factors.

Given the pivotal role of MHC in the immune system, it is vital that the transcription and expression of MHC class I gene expression is tightly regulated and can be induced under appropriate circumstances, such as during a viral infection. Over the past two decades the mechanisms by which cellular factors such as cytokines and transcription factors regulate MHC

class I expression have been unravelled. Focus will now be placed on the cytokines that orchestrate class I transcription, the transcription factors that these cytokines induce and subsequently how these transcription factors interact with the MHC class I promoter to direct transcription. For simplicity, this section will consider the human MHC class I promoters.

1.4.1 Interferon

Interferons were first described as a substance which protected cells against viral infection (Isaacs & Lindenmann, 1957). Following intensive investigation since their discovery, the interferon family of cytokines are now recognised as major components of innate and adaptive immune responses, of particular importance during viral infections. Upon interaction with their receptor, IFNs stimulate the expression of a variety of interferon sensitive genes (ISGs) which, among other effects, culminates in the induction of the 'anti-viral state' where viral replication and protein synthesis are inhibited. Most importantly for this study, interferon stimulation induces the expression and activation of interferon regulatory factors, which bind the ISRE of MHC class I genes, stimulating transcription. During a viral infection, increasing surface class I expression facilitates immune surveillance of foreign peptides by CD8⁺ T cells, promoting cytotoxic killing of virus infected cells. As well as inducing the transcription of MHC class I heavy chains, interferon induces the expression of β 2-microglobulin, peptide transporters, chaperones and the proteosomal subunits essential for the maturation of newly synthesised class I molecules. The IFN families, the induction of IFN gene expression by viral

factors and how IFN signalling pathways culminate in inducing MHC class I gene transcription will now be discussed.

There are two types of interferon, type I and type II, which interact with different cellular receptors to initiate both distinct and overlapping biochemical pathways. Type I IFNs are produced by most cells, are induced by viral infection and include the extensively studied IFN- α and IFN- β . IFN- α is a family of at least 12 members. Each IFN- α species appears to have a distinct profile of biological activities, for example some are more potent stimulators of NK and T cell activity whereas others seem to be more important as antiviral and antiproliferative agents (Bogdan *et al.*, 2004). Several other type I IFNs have been identified including IFN- ϵ and IFN- κ (humans and mice), IFN- ω (humans), IFN- τ (sheep), IFN- δ (pigs) and limitin (mice) [reviewed by (Pestka *et al.*, 2004)]. Recently, a new family of interferon-like cytokines have been discovered in humans: IFN- λ 1, IFN- λ 2 and IFN- λ 3 (also known as IL-28, IL-28B and IL-29 respectively) (Kotenko *et al.*, 2003). Like type I IFNs, IL-28 and -29 induce an antiviral state in cells and induce gene transcription through ISRE motifs, however, in contrast to type I IFNs these cytokines do not appear to have antiproliferative effects (Kotenko *et al.*, 2003; Sheppard *et al.*, 2003).

Type II interferon is IFN- γ . It was first described as an interferon-like virus inhibitor produced by mitogen stimulated lymphocytes (Wheelock, 1965). In general, IFN- γ secretion is restricted to specialised cells of the

immune system including activated NK cells, macrophages and effector T cells although low level production of IFN- γ has been detected in non-immune cells such as fibroblasts and erythroid cells (Rady *et al.*, 1995). IFN- γ is a potent activator of macrophages and induces a T_H1 response by inducing differentiation of naïve T cells into T_H1 helper cells and inhibiting proliferation of T_H2 cells.

1.4.1.1 Induction of type I interferon during viral infections

Type I IFNs are rapidly produced during a viral infection when viral factors such as envelope glycoproteins, CpG DNA, double stranded RNA (dsRNA) and single stranded RNA are recognised by cellular pattern recognition receptors, namely the toll-like receptors (TLRs) and the mannose receptor (MR). These receptors are important on a range of cell types including antigen presenting cells and have been shown to bind to a variety of ligands enabling recognition of a wide selection of pathogens. During a viral infection, the MR and TLRs each recognise distinct intermediates of viral replication and initiate a series of cell signalling cascades which activate the transcription factors ATF2, c-JUN, IRF-3 and NF- κ B. All of the TLRs studied to date have been shown to induce NF- κ B activation in response to viral stimuli *via* signalling cascades that require adaptor proteins, of which MYD88 is the most well understood example (see figure 1.7 for details). By contrast, IRF-3 activation appears to be restricted to TLR3 in viral infections and TLR4 in bacterial infections (Bowie & Haga, 2005). In both cases, activated transcription factors translocate to the nucleus and bind to promoter regions of IFN- α and - β genes, inducing transcription (Figure 1.7).

Figure 1.7: Induction of type I IFNs during a viral infection

This figure shows a summary of the viral factors recognised by individual Toll like receptors (TLRs) and the mannose receptor (MR) and how the transcription of type I interferons is induced by activation of the transcription factors IRF-3, ATF-2/c-Jun and NF- κ B (reviewed by Takeda and Akira, 2005; Moynagh, 2005). PKR and TLR3 recognise dsRNA. The dsDNA-dependant protein kinase (PKR) (green ovals) recognises dsRNA (Alexopoulou *et al.*, 2001). When TLR3 binds dsRNA, TRAF-6, TAK1 and TAB2 (solid gold) form a complex which translocates into the cytosol and interacts with PKR resulting in phosphorylation of the TAK1 subunit. TAK1 then activates IKK and MKK6, which in turn activate NF- κ B and MAP kinase (Takeda *et al.*, 2004). In addition, ATF/c-Jun, IRF-3 and NF- κ B are activated by TLR3 signalling, these pathways are not shown but commonly involve the recruitment of adaptor proteins such as MYD88 (purple square) and activation of IRAK. TLR9 and TLR7 recognise viral DNA in intracellular endosomal compartments (Heil *et al.*, 2004). TLR9 recognises CpG dsDNA. TLR7 recognises ssRNA (Lund *et al.*, 2003). The MR recognises terminal mannose and fucose groups on viral glycoproteins (Milone & Fitzgerald-Bocarsly, 1998).

In response to TLR or MR signalling, IRF-3 forms homodimers, ATF-2 forms a heterodimer with c-Jun and the regulatory subunit of NF- κ B, $\text{I}\kappa\text{B}\alpha$, is lost. These activated transcription factors translocate to the nucleus and bind the promoter regions of type I IFN genes. Transcription of the IFN- β promoter is induced when ATF2/c-Jun complexes bind the PRDIV region, IRF-3 binds the ISRE motif within the PRDIII region and NF- κ B bind the κ B motifs within the PRDII region. IFN- α 1 transcription is induced when IRF-3 binds the ISRE motif.

Figure 1.7: Induction of type I IFNs during a viral infection

Both TLR3 and the dsRNA-dependant protein kinase (PKR) recognise dsRNA. Viruses produce dsRNA during replication as an intermediate in RNA synthesis (RNA viruses) or as a by-product of symmetrical transcription of DNA virus genomes. Interestingly, PKR knockout mice can still induce IFN in response to synthetic analogue of dsRNA, poly(I:C) (Goh *et al.*, 2000). This observation lead to the search for additional receptors that recognise dsRNA and subsequently, Alexopoulou *et al.* (2001) identified TLR3, which binds dsRNA and induces the activation of NF- κ B and IRF-3. This research group also used a series of dominant negative signalling components to demonstrate that the IL-1 receptor-associated kinase (IRAK) and TRAF6 mediate TLR3 signalling. A direct link between TLR3 and viral dsRNA was finally demonstrated by Tabeta *et al.* (2004) using TLR3 $-/-$ mice in which TLR3 was shown to contribute to type I IFN production in response to murine CMV infection.

Both TLR9 and TLR7 recognise DNA sequences contained within intracellular endosomal compartments. TLR9 recognises DNA sequences containing unmethylated CpG dinucleotides released from endocytosed viruses (Lund *et al.*, 2003) and induces type I IFN *via* a MYD88-dependent pathway. TLR7 (mouse) and TLR8 (human) recognise GU-rich ssRNA in endosomal compartments (Heil *et al.*, 2004). Since TLRs 7, 8 and 9 can also recognise host DNA *in vitro*, it is assumed that their function is to recognise abnormal localisation of nucleic acids rather than 'non-self' sequences or motifs. To put this point into context, host DNA is rarely contained within

endosomal compartments, whereas many viruses enter cells by hijacking cellular receptor mediated endocytosis and releasing DNA into endosomes.

Mannose receptors (MR), expressed on natural interferon producing cells (NIPC), bind to glycosylated viral envelope proteins and mediate IFN synthesis (Milone & Fitzgerald-Bocarsly, 1998). Studies using viruses containing mutated N-linked glycosylation sites and viruses treated with glycosylase enzymes demonstrated that correct glycosylation of envelope proteins is required for recognition by the MR (Charley *et al.*, 1991; Laude *et al.*, 1992). Human glycoproteins typically contain N-acetylglucosamine and sialic acid groups and therefore the MR can differentiate between self and non-self glycoproteins.

1.4.1.2 Induction of IFN- γ

As previously discussed, the majority of IFN- γ produced originates from activated immune effector cells. In common with the IFN- α and - β genes, the IFN- γ gene contains complex arrangement of positive and negative regulatory elements that bind transcription factors in a co-operative manner and orchestrate the transcriptional activation of this gene. Among these transcription factors are NF- κ B, members of the nuclear factor activating transcription factor (NFAT) family (Sica *et al.*, 1997), signal transducer and activator of transcription (STAT), T-bet (Szabo *et al.*, 2000), AP-1, CREB-ATF, GATA-3 and yin-yang (Ye *et al.*, 1996).

1.4.1.3 IFNs induce the transcription of ISGs - the 'delayed' response

Following transcription, secreted IFN can have an autocrine effect, acting on the secreting cell or a paracrine effect, stimulating neighbouring cells and thus preventing viral replication and further spread of the virus. In both cases, IFNs binds to specific cellular receptors: IFN- α and IFN- β bind IFNAR-1 and IFNAR-2 and IFN- γ binds IFNGR-1 and IFNGR-2. Receptor binding initiates signal transduction cascades involving the Janus activated kinases (JAKs) and members of the signal transducer and activator of transcription (STAT) family transcription factors. JAKs phosphorylate tyrosine residues of STATs which then form hetero or homodimers, translocate to the nucleus and bind the regulatory promoter elements of IFN sensitive genes. STATs drive the transcription of an array of genes, many of which encode transcription factors and proteins with anti-viral properties. Most importantly for this study, IFNs induce the activation of members of interferon regulatory factor (IRF) family of transcription factors and NF- κ B, which in turn induce the transcription and expression of MHC class I.

1.4.2 The Interferon regulatory factor (IRF) family of transcription factors

IRFs are a family of seven transcription factors that are activated in response to IFNs and viral infections and induce the expression of IFN inducible genes including MHC class I. IRFs mediate diverse transcriptional responses to IFN stimulation that depend on the cell type, interactions with other transcription factors and post-translational modifications of the IRF involved. Some IRFs are transcriptional inducers while others have been

shown to be transcriptional repressors. Characteristically, each IRF contains a 155 amino acid DNA binding domain comprising five repeated tryptophan residues and it is through this motif that they bind the ISREs of IFN sensitive genes. In humans, IRF-1, -2, -3 -7 and -9 bind the consensus sequence G/A G/C TTTC of HLA class I promoters (Chang *et al.*, 1992; Drew *et al.*, 1995b; Drew *et al.*, 1995a; Massa & Wu, 1995; Gobin & van den Elsen, 2000; Lefebvre *et al.*, 2001).

IRF1 and IRF-2 were the first members of the IRFs to be characterised. They are constitutively expressed at low levels in most cell types and their expression and activation is induced by protein kinase A (PKA), protein kinase C (PKC) or casein kinase II which in turn are induced by IFN- α , - β , - γ , TNF- α and interleukin -1 and -6. Upon activation IRF-1 is a transcriptional activator. It binds the ISRE of HLA-A, -B, -C and -F promoters and the distal ISRE of the HLA-G promoter driving transcription in response to IFN- γ stimulation (van den Elsen *et al.*, 2004). IRF-1 can induce transcription in co-operation with other transcription factors; following co-stimulation with TNF- α and type 1 IFNs it forms a complex with NF- κ B and mediates synergistic induction of class I expression (Drew *et al.*, 1995b; Johnson, 2003). In addition to HLA class I genes, IRF-1 induces the transcription of TAP1 and LMP2, demonstrating the importance of this transcription factor in global regulation of antigen presentation (White *et al.*, 1996).

IRF-2 competes for ISRE binding sites with IRF-1 and was first described as a transcriptional repressor of the IFN- β gene (Harada *et al.*, 1989). We now know that IRF-2 is a transcriptional repressor of a variety of ISGs and has been shown to bind the ISRE of HLA-A, -B -C and -F and repress transcription (Gobin *et al.*, 1999). It transpires that IRF-2 can also function as a transcriptional activator and has been shown to induce the transcription of a functionally diverse group of genes including vascular cell adhesion molecule -1 (VCAM-1), the human histone 4 gene and the MHC class II transactivator (CIITA) (Jesse *et al.*, 1998, Masumi *et al.*, 2003, Xi & Blanck, 2003). Induction of CIITA is not only unusual because this gene is actually an ISG, which are usually repressed by IRF-2, but also because in this instance IRF-2 co-operates with IRF-1 to facilitate CIITA transcription (Xi & Blanck, 2003).

IRF-3 is particularly important in the immediate, early response to dsRNA and the induction of IFNs during viral infections. It expressed constitutively and becomes activated during a viral infection when dsRNA is detected by TLR3, -4 and PKR as described above (Au *et al.*, 1995). Phosphorylated IRF-3 translocates to the nucleus where it can bind to and induce transcription of ISGs including the cytokines IFN- α , IFN- β (reviewed in Malmgaard, 2004) and Regulated on Activation Normal T-cell Expressed and Secreted (RANTES) (Lin *et al.*, 1999). Direct binding of IRF-3 to HLA promoters has not been described in the literature but is assumed, given the homology of the IRF-3 binding domain to that of IRF-1 and -2.

IRF-7 is only constitutively expressed at low levels in lymphoid tissues but becomes expressed at high levels and activated by phosphorylation following IFN stimulation (Zhang & Pagano 2002). Activated IRF-7 can either form a homodimer or a heterodimer with IRF-3. IRF-7/3 heterodimers are important in the delayed response to viral infection where they provide positive feedback and potentiate IFN induction. IRF-9 is constitutively expressed in an inactive form and lacks the intrinsic ability to induce transcriptional responses as a mono- or homo-dimer. However, IFN signalling induces this transcription factor to form a heterotrimeric complex with STAT1 and STAT2, called ISGF3 γ which binds to the promoters of ISGs (Kraus *et al.*, 2003).

1.4.3 Nuclear Factor kappa B

Nuclear factor kappa B (NF- κ B) is considered to be a central mediator of cellular stress responses because of the diversity of the signals that activate it and the genes that it regulates. It was first identified in B lymphocytes as a nuclear transcription factor that binds to a 10 bp DNA element in the kappa immunoglobulin light-chain promoter enhancer motif (Sen *et al.*, 1986). We now know it modulates the expression of a spectrum of genes involved in cell adhesion, cell proliferation, apoptosis, cell migration, angiogenesis and the immune response where it activates transcription of cytokines, chemokines, interleukins, immunoglobulins, acute phase proteins and MHC class I genes. Of particular relevance to this study is the activation

of NF- κ B by the cytokine tumour necrosis factor alpha (TNF- α) and by TLR signalling during a viral infection (described above).

Five distinct NF- κ B subunits have been identified including NF- κ B1 (p50/p105), NF- κ B2 (p52/p100), RelA (p65), RelB, and c-Rel. NF- κ B1 and NF- κ B2 undergo proteolytic processing to liberate p50 and p52, respectively. Each NF- κ B subunit contains a conserved 300 amino acid domain, called the rel homology domain (RHD), which is responsible for subunit dimerisation as well as DNA binding. In resting cells, inactive NF- κ B homo- or heterodimers are constitutively present in the cytoplasm with I κ B, the intracellular inhibitor of NF- κ B. Upon activation by cytokines I κ B is inactivated and degraded releasing NF- κ B, which translocate to the nucleus where it binds promoter regions containing the κ B motif GGGGATTCCCC. Classical HLA class I promoters contain two κ B motifs within the Enhancer A region: κ B1 and κ B2. Gobin *et al.*, (1998) tested these motifs in classical and non-classical promoters for transcription factor binding and activation, using HLA class I κ B sequences to drive the expression of a luciferase reporter. In HLA-A, both κ B sites bind to p65, p50 and c-Rel subunits of NF- κ B and co-transfection of NF- κ B strongly induced luciferase activity driven by the HLA-A Enhancer A region. By contrast, only the κ B1 site of HLA-B binds NF- κ B and accordingly, transactivation by NF- κ B was weaker than for HLA-A, demonstrating locus specific activation by this transcription factor. Neither of the κ B sites of HLA-C bind, or are induced by NF- κ B, although an alternative transcription factor Sp1 was shown to bind this region. The non-classical

HLA are also differentially transactivated by NF- κ B. The κ B1 site, but not κ B2, of HLA-F binds p50 and p65 and the κ B sites of HLA-E do not bind any NF- κ B species. The κ B sites of HLA-G, the most divergent of all HLA class I, only bound p50 homodimers. None of these genes were transactivated by NF- κ B in co-transfection studies. The locus-specific variation in NF- κ B mediated regulation of HLA correlates to nucleotide differences in κ B motifs and explains in part, different constitutive and cytokine-induced levels of classical and non-classical HLA expression. More recently, a putative κ B site in the promoter region of MICA was shown to bind p50 and p65 homo-dimers and mediate an increase in MICA mRNA in activated T-lymphocytes following inhibition of I κ B (Molinero *et al.*, 2004).

1.4.4 The SXY region and the MHC class I enhanceosome

The SXY module of MHC class I promoters consists of four regulatory elements namely the S (or W) box, the X1 and X2 boxes and the Y box which are co-operatively bound by a multi-protein complex which functions as an enhanceosome and drives transcription of class I (and class II) genes. The enhanceosome comprises; regulatory factor X5 (RFX5), RFXB and RFXAP, collectively known as regulatory factor X (RFX), cyclic AMP response element binding protein (CREB)/activating transcription factor (ATF) and nuclear factor Y (reviewed by van den Elsen *et al.*, 2004). These transcription factors are collectively known as the MHC class I enhanceosome. An additional transcription factor, the MHC class II transactivator (CIITA) also directs MHC class I transcription *via* the SXY

module. CIITA was first described as a transcription factor required for constitutive and IFN- γ induced expression of MHC class II, however it transpires that it can also transactivate MHC class I genes (Martin *et al.*, 1997). CIITA does not bind directly to promoter DNA sequences. It interacts with RFX/CREB/NFY transcription factors bound to the SXY box region of the class I and class II promoters. The crucial importance of the SXY module in MHC class I and II transactivation is demonstrated in patients lacking one of the RFX subunits or CIITA, who suffer from a severe combined immunodeficiency called Bare Lymphocyte Syndrome (BLS), in which patients completely lack expression of MHC class II and have reduced levels of expression of MHC class I (Reith & Mach, 2001).

1.4.5 Summary of Human MHC class I promoter regulation

MHC class I expression in humans is regulated by the interaction of transcription factors with regulatory motifs within the promoters of MHC class I genes. These mechanisms of transcriptional regulation provide a model for comparative studies of porcine MHC class I regulation. Our current understanding of transcription factor-promoter interactions in the human is summarised in Figure 1.8. The key motifs in class I promoters that regulate transcription are; an Enhancer A which binds NF- κ B, an ISRE which binds members of the interferon regulatory factor (IRF) family of transcription factors (shown in the figure 1.8 as IRF-1 for simplicity) and an SXY domain which binds RFX/CREB/NFY and CIITA. Overall, the expression of classical class I genes is (HLA-A –B and –C) is regulated by these transcription

Figure 1.8: Transcriptional regulation of HLA class I genes

The regulation of classical and non-classical HLA class I genes at the promoter level is summarised above in the context of four core promoter regions: the distal ISRE, Enhancer A, ISRE and SXY. STAT1 homodimers bind the distal ISRE of HLA-E following IFN- γ stimulation, a similar motif in the HLA-G promoter binds IRF-1 in response to IFN- α . The Enhancer A regions of HLA-A, -B, -E and -F bind NF- κ B in response to multiple stimuli, including TNF- α . The ISRE of HLA-A, -B, -C and -F bind IRFs in response to IFNs. With the exception of HLA-G, HLA class I SXY motifs bind RFX CREB and NF-Y (green) and are further transactivated by co-operative binding of CIITA. Comparison of HLA promoters with the β 2-microglobulin promoter shows that the same regulatory motifs and transcription factors control expression of this gene, although the arrangement of these motifs differs in the β 2-microglobulin promoter. Adapted from van den Elsen *et al.*, 1998; 2004).

factors, with the exception of HLA-C, which does not bind NF- κ B. Nucleotide variation in Enhancer A and ISRE sequences directly influences the binding of transcription factors to these sites and therefore the level of promoter activation and class I expression induced by IFN and NF- κ B pathways differs between MHC class I loci.

The non-classical genes are divergent from classical HLA genes both in sequence and function. This is reflected in the transcriptional regulation of these genes. HLA-G is expressed on trophoblast cells at the materno-foetal interface and was also shown to be expressed in melanoma tumours, and may be involved in immunotolerance of tumour cells. Accordingly, the tight transcriptional regulation of this gene is reflected at the promoter level. HLA-G is not induced by NF- κ B, IRF-1 or CIITA *via* the usual motifs. Instead, HLA-G transcription is induced by IFNs *via* binding of IRFs to a hybrid distal GAS/ISRE element located upstream of the core regulatory motifs (Lefebvre *et al.*, 2001). The Enhancer A of HLA-G binds p50 and the generic transcription factor Sp1, but is not transactivated by NF- κ B p65 (Rousseau *et al.*, 2000). In addition, there is no apparent regulation by transcription factor binding to the SXY region and HLA-G is not transactivated by CIITA. This is because the X1 box does not bind RFX5 and therefore CIITA binding is prevented (Rousseau *et al.*, 2004). HLA-G expression is also up regulated by cell stress and contains a heat shock element that binds to heat shock factor 1 (HSF-1) (Ibrahim *et al.*, 2000)

Like HLA-G, HLA-E, is not induced by NF- κ B or IRF-1 but is induced in response to IFN- α *via* STAT1 binding to a distal GAS upstream of the ISRE that mediates classical HLA responses to type 1 IFN (Gobin & van den Elsen, 2000; Gustafson & Ginder, 1996). HLA-E is also responsive to CIITA through the SXY regulatory module.

Interestingly, the β 2-microglobulin promoter shares the same regulatory motifs as HLA class I promoters, although the organisation of these motifs differs slightly, the ISRE is 5' to the Enhancer A.

1.4.6 Regulation of expression of MHC class I genes of the d/d pig

In 1982, Singer *et al* used an HLA cDNA to probe genomic DNA obtained from pigs homozygous at the d locus (d/d pig) and isolated a genomic clone encoding a major SLA antigen, PD1 (paralogous to SLA-1). When transferred into mouse L cells this gene directed the synthesis of SLA heavy chains which associated with murine β 2-microglobulin, were expressed at the cell surface and were induced by mouse IFN- α (Satz & Singer, 1984). Similar results were obtained with a second SLA antigen, PD14 (Satz *et al.*, 1985). Both of these studies provide evidence that the mechanisms regulating constitutive and IFN-induced expression of MHC class I are conserved in the mouse and pig. Therefore the region which controls transcription, the MHC class I promoter, was examined. Preliminary sequencing of PD1 and PD14 identified a series of conserved regulatory domains in the 5' regions upstream of the transcriptional start site including a

CAT and TATA box. Subsequent studies identified a positive and negative regulatory element within the promoter of PD1 that controlled expression (Ehrlich *et al.*, 1988). The negative element was unique as it shared no homology with other known regulatory elements and its activity was dependent upon a downstream enhancer. A transgenic mouse expressing PD1 was made, skin grafts from this transgenic mouse were rejected in the parental strain, demonstrating that SLA was expressed and recognised as a functional transplantation antigen (Frels *et al.*, 1985). The tissue distribution of PD1 expression in the transgenic mouse was similar to that observed in the pig, indicating that the cellular factors that regulate porcine MHC class I expression are conserved between mouse and pig species. This was confirmed by treatment of transgenic mice with IFN- α/β , which increased PD1 expression. Interestingly the magnitude of SLA class I responses to type I IFN induction varied between tissues; IFN-induced PD1 expression in the brain was 6 times higher than in lymph nodes (Satz & Singer, 1984). The conclusions of this study were that both constitutive and IFN induced expression of PD1 is controlled in a tissue specific manner and that regulation is controlled by a 1 kb region 5' to the PD1 coding sequence. A putative ISRE motif was subsequently identified in the promoter region of PD1. Analysis of a series of truncated PD1 promoter constructs demonstrated that deletion of the ISRE region abolishes IFN- α/β induced expression *in vitro* and that deletion of two upstream silencers (-894 and -516) increases constitutive PD1 expression in transfection studies *in vitro* and *in vivo* in a second transgenic mouse expressing PD1 containing a truncated 5' flanking region (Maguire *et al.*, 1990). Further analysis of a 589

bp region containing the novel silencer domain region -503 to -1091 upstream of PD1 coding sequence identified two functional motifs associated with this region: an enhancer (-731 to -771) and silencer (-676 to -731). Distinct cellular factors are associated with each motif and the binding sites for each group of factors overlap. It was demonstrated that levels of MHC class I expression are directly associated with the levels of silencer binding factors present in a particular cell type. Although enhancer binding factors were shown to be preferentially expressed, binding of these factors to the enhancer motif is inhibited in the presence of silencer binding factors (Weissman & Singer, 1991). For example, PD1 expression is low in the kidney, where high levels of silencer binding factors are expressed and high in B cells where silencer binding factor expression is low.

Concurrently, the expression of a second classical MHC class I gene PD7, with high sequence homology to PD1, was examined in pig and transgenic mouse tissues (Frels *et al.*, 1990). Interestingly, PD7 was expressed at much lower levels in tissues than PD1. In order to find an explanation for this observation two strains of transgenic mice which differed in the length of the 5' flanking region upstream of PD7 revealed a silencer element with tissue specific activity, which regulates PD7 expression in lymphoid tissues and skin and contributes to the tissue specific expression of this gene. Sequence comparison of the Enhancer A region of PD1 (GGGGAGTCCCC) with PD7 (G**AGG**AGTCCCC) reveals a mutation in the second residue (highlighted in bold). This correlates with observations in humans where mutations in second and tenth residues of Enhancer A

prevents binding of a nuclear factor and abolishes Enhancer A activity (Johnson, 2003), therefore it was hypothesised that this nucleotide difference may account for reduced expression of PD7 in a number of tissues in vivo compared to PD1.

1.4.7 Regulation of expression of MHC class I genes of the H01 pig

Recently, the MHC class I region of the H01 pig was fully sequenced revealing 8 class I genes (SLA-1, -2, -3, -5, -6, -7, -8 and MIC2) with the correct intron/exon organisation in order to be functional (Chardon *et al.*, 2001; Renard *et al.*, 2001). Analyses of 5' flanking regions of these genes reveal promoter elements that share homology with the promoter regions of human and mouse MHC class I genes and those of the d/d pig already discussed here. The functions of the regulatory motifs encoded within class I genes of the H01 pig have not been investigated experimentally before and are the focus of this study.

1.5 Viruses and MHC class I

Given the importance of MHC class I in anti-viral immunity, it comes as no surprise that many viruses have evolved ways of circumventing class I mediated antigen presentation. To date, an impressive number and variety of mechanisms of disrupting MHC class I have been documented. There are viral mechanisms that target almost every step and component of the MHC class I antigen presentation pathway, from inhibition of transcription through to endocytosis and degradation of cell surface MHC class I. Figure 1.9

Figure 1.9: Viruses target the MHC class I antigen presentation pathway in order to avoid immune recognition.

A summary of the MHC class I presentation pathway from gene transcription through to cell surface expression is shown. At each step in this pathway examples of viral intervention, leading to reduced or impaired peptide presentation are indicated in bold black text and the corresponding virus and viral protein (where known) are indicated in green text. Specific organelles are labelled in blue text. In an uninfected cell, peptides are produced by the proteasome and transported into the ER by the TAP transporter, where they are loaded onto newly synthesised MHC class I heavy chains. Fully assembled MHC class I is transported to the cell surface via the trans golgi network.

provides an overview of the MHC class I presentation pathway and some examples of viruses and specific viral proteins that disrupt it.

Adenovirus type 12 down regulates MHC class I expression by targeting the transcription of MHC class I at the promoter level. In Adenovirus type 12 transformed cells, nuclear NF- κ B is hypophosphorylated and cannot bind and drive transcription of the MHC class I promoter (Hou *et al.*, 2002). In addition, a transcriptional repressor COUP-TFII remains associated with the MHC class I promoter, preventing binding and transactivation by NF- κ B.

Several viruses use cell surface MHC class I as a target for immune evasion. Varicella Zoster Virus (VZV) has been shown to down regulate surface expression of MHC class I molecules on human fibroblasts and T cells early in infection. This is mediated by the product of the early gene ORF66. Further biochemical analyses demonstrated that MHC class I molecules were retained in the golgi following synthesis in the ER but the mechanism for this remains unclear (Abendroth *et al.*, 2001).

HIV-1 exploits the cellular phosphoferin acidic cluster sorting protein-1 (PACS) in order to disrupt MHC class I cell surface expression (Piguet *et al.*, 2000). PACS protein is a 'coat' protein that organises the transport of furin protein between transport endosomes and the golgi. The HIV Nef protein bridges together the cytoplasmic domains of PACS protein and MHC class I this results in the removal of MHC class I from the cell surface by

endocytosis. MHC class I is then transported to the trans golgi network (TGN) where they are retained. Similarly, the K3 protein of Kaposi's Sarcoma-associated Herpesvirus (KSHV) removes MHC class I from the cell surface by increasing endocytosis. Class I molecules are then targeted to lysosomes (*via* the TGN) where they are destroyed (Hou *et al.*, 2002).

Herpes simplex virus type 1 (HSV-1) and type 2 (HSV-2) are able to inhibit expression of MHC class I by a different mechanism. Both these viruses encode ICP47, an immediate early protein that blocks MHC class I assembly by inhibiting the transporter associated with antigen presentation (TAP), thereby preventing the transport of peptides into the ER (Jugovic *et al.*, 1998). MHC class I molecules cannot be assembled without peptides and therefore no new MHC class I molecules are transported to the cell surface.

While it is advantageous to block some expression of MHC class I, complete blockage of MHC class I expression ('missing self') would attract the attention of NK cells and would not be advantageous for viruses. Therefore, in order to evade destruction by both NK and CD8⁺ T cells a virus must employ strategies to disrupt both MHC class I expression of peptides as well as prevent recognition by NK cells. Human cytomegalovirus (HCMV) is a well documented example of a virus that is proficient in immune evasion, particularly in modifying MHC class I expression. HCMV encodes a number of unique short (US) viral proteins dedicated to the disruption of classical HLA class I expression at several stages of the peptide presentation pathway

resulting in the prevention of peptide presentation to CD8⁺ T cells. Interestingly, HLA-A expression is inhibited early during infection whereas disruption of HLA-G expression occurs later (48hpi), and to a lesser extent than HLA-A alleles. It is hypothesised that differential regulation of HLA-G occurs so that a continuous interaction with the inhibitory ligand KIR2DL4 is maintained, ensuring that NK cell activation and secretion of IFN- γ is prevented (Pizzato *et al.*, 2004).

Given the fundamental importance of MHC class I in the immune response to viruses it is anticipated that disruption of SLA class I would be an important immune evasion strategy for viruses that infect pigs, particularly for viruses that cause persistent infections, such as Classical Swine Fever virus (CSFV). The immune evasion strategies of CSFV are not well understood, a greater understanding of these mechanisms could provide valuable information with applications in vaccine design and target identification for therapeutic intervention.

1.6 Classical Swine Fever

Classical Swine Fever (also known as hog cholera) is a highly contagious, often fatal disease that affects domestic pigs and wild boar (*Sus scrofa*) with the latter acting as a natural reservoir for the disease. Inactivated virus can be used safely and efficiently as a vaccine, however vaccinated animals cannot be serologically distinguished from animals infected with field isolates and therefore vaccination was banned by the European Union in 1990. Until a better vaccine or medicinal therapeutic is

developed, slaughter remains the only truly efficient means of controlling spread of this disease when an outbreak occurs. Outbreaks of CSF carry a high economic cost; the 1997-1998 epidemic in The Netherlands is estimated to have cost around US\$ 2.3 billion and required some 11 million pigs to be slaughtered (Meuwissen *et al.*, 1999).

1.6.1 Clinical perspective and causative agent

CSF causes a variety of clinical disease states ranging from acute to chronic. Disease symptoms vary for each state but are generally characterised by fever, leukopenia, haemorrhage and mortality of up to 90% in cases of severe acute infection. Infection during early stages of pregnancy often results in abortion and stillbirths whereas infection during late stages of pregnancy results in the birth of persistently infected animals. These animals become reservoirs for infection.

The causative agent of CSF is classical swine fever virus (CSFV). CSFV is a member of the *Pestivirus* genus in the *Flaviviridae* family, which also includes bovine viral diarrhoea virus (BVDV) and border disease virus (BDV). The CSFV genome is a single positive-stranded RNA of about 12.5 kb in length, comprising a single open reading frame which encodes a 4000 amino acid polyprotein. The polyprotein is co- and post-translationally processed into 12 proteins, summarised in figure 1.10. The N terminal protease N^{pro} is a cysteine autoprotease and is of particular interest for this study because of its recently demonstrated role in host immune evasion.

Normally during a viral infection, IRF-3 is activated and translocates to the nucleus where it binds to IFN- α and - β promoters, initiating transcription of these genes. During CSFV infection, N^{pro} inhibits IFN- α production by depleting IRF-3 (Larocca *et al.*, 2005). In addition, CSFV has been shown to inhibit activation and IFN- α production by dendritic cells (DCs), although the mechanism and involvement of N^{pro} in the inhibition of type 1 IFNs has not been determined in this system (Carrasco *et al.*, 2004).

As mentioned above, CSFV causes a persistent infection and therefore must be able to avoid recognition by the host immune response. One important question that remains to be answered is how does CSFV achieve a persistent infection and does this virus affect MHC class I expression in order to achieve this? One aim of this thesis is to address this question, but in order to study MHC class I expression one must first understand the mechanisms that regulate the expression of this family of genes. Understanding class I expression and the mechanism that CSFV uses to disrupt it would provide valuable information for vaccine design and may identify targets for therapeutic intervention.

Objectives

The aims of this project were:

- To characterise cell lines and antibodies that can be used as tools to study SLA class I expression.
- To identify cellular factors that regulate MHC class I expression at the cell surface.
- To produce tools that can be used to investigate SLA class I expression at the promoter level.
- To determine which SLA class I gene promoters are constitutively active.
- To examine sequence variation in putative regulatory motifs identified by Chardon *et al* (2001) and make predictions about which transcription factors will bind these motifs and drive transcription.
- To evaluate transcription factor activation of MHC class I expression experimentally and compare promoter responses of classical and non-classical SLA class I genes.
- To relate variation of classical and non-classical SLA class I gene responses to transcription factors to sequence variations within SLA class I promoter regions.
- To determine the effect of CSFV on SLA class I expression and understand the mechanism of this immune evasion strategy.

Chapter 2. Materials and Methods

2.1 Reagent suppliers

Unless otherwise stated chemicals were supplied by Sigma-Aldrich Company (Dorset, England), buffers and kits for molecular biology were purchased from Qiagen (Germany) and enzymes were purchased from Promega (U.K.).

2.2 Media, buffers and solutions

All media were autoclaved or filter-sterilised before use. All solutions were prepared in sterile or distilled water unless otherwise stated.

2.2.1 Cell culture media

RPMI: Roswell memorial park institute medium (RPMI) (Invitrogen) containing 20% (v/v) foetal calf serum (FCS), 100 U/ml streptomycin, 100 U/ml penicillin and 20mM L-glutamine. Only FCS that has been certified free from bovine viral diarrhoea virus (BVDV) was used. Certified sera is free from BVDV specific antibodies and viral particles following propagation in cell culture for several passages. Prior to use, FCS was heated to 56°C for 30 mins to inactivate complement.

IMDM: Iscoves Modified Dulbecos Medium (IDIM) (Invitrogen) supplemented with 10 % (v/v) FCS (certified BVDV free), 100 U/ml streptomycin, 100 U/ml penicillin and 20 mM L-glutamine. Serum free IMDM was as above but did not contain FCS.

EDTA-trypsin: 136 mM NaCl, 5.3 mM KCl, 5.5 mM NaHCO₃, 0.02% (w/v) trypsin, 0.01% (w/v) ethylenediaminetetraacetic acid (EDTA), and 0.1% (w/v) phenol red.

2.2.2 Bacterial culture media

Luria Broth (LB) medium: 1% (w/v) tryptone, 0.5% (w/v) yeast extract, 0.5% (w/v) NaCl in dH₂O.

Agar plates: 1.5% (w/v) Bacto agar (Oxoid N° 1) in LB medium.

Ampicillin resistance: 50 µg/ml ampicillin.

2.2.3 Immunofluorescence microscopy buffers

Paraformaldehyde fixative: 4% (w/v) paraformaldehyde (PFA) in PBS, pH 7.5-8.0.

Methanol/Acetone Fixative: 50% (v/v) methanol, 50% (v/v) acetone.

Blocking buffer: 50mM Tris (pH 7.4), 150mM NaCl, 0.2% (w/v) gelatine, 1% (v/v) Igepal, 30% (v/v) 30% normal goat serum (NGS) (Harlan Sera Labs UK).

Mouse block buffer: 2% (w/v) normal mouse serum (Harlan Sera Labs UK) in blocking buffer.

Wash buffer: 0.1% (v/v) Tween-20 in PBS.

Reporter assay buffers

Z buffer: 60 mM Na₂HPO₄, 40 mM NaH₂PO₄, 10 mM KCl, 1 mM MgSO₄

ONPG substrate buffer: 60 mM Na₂HPO₄, 40 mM NaH₂PO₄, adjusted to pH 7.0 and containing 4 mg/ml o-nitrophenyl- β -D-galactopyraniside (added immediately prior to use).

2.2.4 FACS Buffers

FACS diluent: 0.6% bovine serum albumin (w/v) 0.6% sodium azide (v/v) in phosphate buffered saline and 2% (v/v) horse serum, adjusted to pH 7.4.

FACS fixative solution: 1.0 to 1.5% PFA in PBS.

2.2.5 DNA manipulation buffers

Agarose gel DNA electrophoresis buffers

50X TAE buffer: 2 M Tris-base, 5.5% (v/v) glacial acetic acid, 50mM EDTA.

Ethidium bromide solution: 1.5% (w/v) ethidium bromide in dH₂O.

2.2.6 Plasmid and Bacterial Artificial Chromosome buffers

DNA preparation buffers

Suspension medium: 50 mM glucose, 25 mM Tris/HCl (pH 8.0), 10 mM EDTA.

Lysis buffer: 0.2 M NaOH, 1% (w/v) SDS.

Potassium acetate buffer: 3 M potassium acetate, 11% (v/v) acetic acid.

TE Buffer: 10 mM Tris-HCl, pH 8.0 and 1 mM EDTA, pH 8.0.

2.3 Cell lines and viruses

2.3.1 Cell lines

Max cells: Max cells (swine kidney cells that express the SLA d/d haplotype) were obtained from the Federal Research Centre for Virus Diseases of Animals (Tübingen, Germany) and have been described previously (Pauly *et al.*, 1995). Max were grown at 37°C in IMDM supplemented with 10% FCS.

Porcine aortic endothelial cells (PAEC): PAECs were isolated from swine aorta from outbred pigs of unknown haplotypes (referred to as outbred PAECs). d/d PAECs were obtained from a d/d minipig (referred to as d/d PAECs). Animal tissues were donated by H. Takamatsu (IAH, Pirbright) from animals kept under Home Office regulations and in accordance with the necessary ethical considerations. Both PAECs and d/d PAECs were prepared using the protocol described previously (Vallee *et al.*, 2001). Briefly, the aorta was removed from an outbred or d/d minipig (aged 6-8 weeks). Minor blood vessels were ligated and the aorta was washed thoroughly by flushing with PBS and RPMI media. The aortal lumen was filled with collagenase IV solution (0.1 % in PBS, Worthington) and incubated at 37°C for 4 minutes. The aorta was then massaged and rinsed with 50 ml of RPMI media. Luminal cells were pelleted by centrifugation (500 g, 10 min), resuspended in RPMI and plated onto 25 cm² tissue culture flasks pre-coated with gelatine (10% in PBS). PAECs were maintained on gelatine-coated flasks and cultured at 37°C in RPMI supplemented with 20% FCS.

Shimozuma cells: The Shimozuma cell line was provided by Dr. H. Takamatsu, IAH Pirbright. These cells were derived from a pig with malignant lymphoma producing porcine endogenous retrovirus (PERV), both the retro virus and PRV reverse transcriptase have been isolated and well characterised (Suzuka *et al.*, 1986). Shimozuma cells were grown in IMDM under standard cell culture conditions.

2.3.2 Virus and infection protocols

2.3.2.1 CSFV stock and preparation

The virulent isolate of CSFV Brescia, used for all infections, was provided by the Institute for Animal Science and Health (Lelystad, The Netherlands) and adapted for cell culture by serial passage on Max cells in the manner described by Van Gennip *et al.*, (2004). Virus was isolated by freeze-thaw lysis and centrifugation at 1500g for 10 minutes and titrated by immunostaining with anti-E2 antibody WH303 (Bensaude *et al.*, 2004) and used in experiments at a multiplicity of infection (moi) of 2 TCID₅₀. The virus containing supernatant was stored at 4°C or -70°C.

2.3.2.2 Virus infection

For CSFV studies, Max cells were infected for 1 hour at 37°C, washed in PBS then supplemented with IMDM (containing 10% FCS) or RPMI (containing 20% FCS) respectively. Infection was allowed to progress for the time indicated in each experiment. Flasks of uninfected cells were maintained in cell culture for the same time period as infected cells and were used as negative controls in FACS and transfection studies.

2.4 Antibodies

2.4.1 Antibodies recognising MHC class I

The monoclonal antibodies 2.27.3 (Ivanoska *et al.*, 1991), 74.11.10 (Pescovitz *et al.*, 1984), 2.12.3 and 2.32.1 [Ivanoska *et al.*, unpublished data cited in (Lunny & Pescovitz, 1988)] and 16.7E4 (Lunney, 1994) were supplied as hybridoma cell lines to the Institute for Animal Health, Pirbright. Antibody preparations were supplied by Brenda Jones (Institute for Animal Health, Compton). Immunoglobulins were isolated from the hybridoma using the miniperm system according to manufacturers protocol (Sartorius Ltd, Epsom, UK). The monoclonal antibody PT-85 was purchased from VMRD (Pullman, WA) and has been described previously (Davis *et al.*, 1987). Briefly, thymocytes from outbred pigs were used to immunise Balb/c (H2-d) mice and hybridomas were produced, those that lysed 97% of porcine cells were selected.

2.4.2 Conjugated antibodies and dyes

Goat anti-mouse and anti-rabbit Alexa⁴⁸⁸ (green) and Alexa⁵⁹⁴ (red) conjugates were purchased from Molecular Probes (The Netherlands). Goat anti-rabbit and anti-mouse horseradish-peroxidase (HRP) conjugates were purchased from Promega (UK). Goat anti-mouse IgG, IgG2a, IgG2b and IgM antibodies conjugated to phycoerythrin were purchased from Biosource (UK). 4'-6-diamidino-2-phenylindole (DAPI) was purchased from Sigma. Phalloidin⁵⁹⁴ was purchased from Molecular Probes (UK).

2.5 Plasmid DNA tools

2.5.1 Transfection control plasmids

The vector pJATLacZ, a β -galactosidase reporter gene under the control of the rat β -actin promoter (Didcock *et al.*, 1999) was used as a control for transfection efficiency and normalisation of luciferase assay data.

The pGL3-Control vector (Promega) contains the firefly (*Photinus pyralis*) luciferase gene under the control of SV40 promoter and enhancer sequences and was used as a positive control for luciferase activity in transfection experiments.

2.5.2 Transcription factor activity control plasmids

The control plasmids used to monitor transcription factor activity were kindly provided by Dr. Stephen Goodborn (St. George's Hospital Medical School, Tooting, London).

The reporter plasmid p(9-27 ISRE)4tk Δ (-39) lucifer contains 4 copies of the ISRE consensus sequence (AGGAAATAGAAACTG) arranged in tandem upstream of the firefly luciferase reporter gene of the ptk Δ (-39) Lucifer plasmid (Reid *et al.*, 1989; King & Goodbourn, 1998) and was used as a control to monitor responses to IFN- α .

The luciferase reporter plasmid p(IRF-1*GAS)6tk Δ (-39) Lucifer contains 6 copies of the gamma activated sequence (GAS) consensus sequence (TTTCCCCGAAA) arranged in tandem upstream of the firefly

luciferase reporter gene of the ptk Δ (-39)lucifer plasmid (King & Goodbourn, 1994; 1998).

The luciferase reporter plasmid p κ B6tkluc, contains 6 NF- κ B binding sites upstream of the luciferase gene under the control of the Herpes Simplex thymidine kinase promoter, and was used as a positive control to monitor cellular responses to TNF- α and NF- κ B p65 co-transfection.

A control reporter p[(AAGTGA)4]5tk Δ (-39)lucifer (HEX-4) was used to monitor IRF-1 activity. This plasmid contains a luciferase gene under the control of a synthetic promoter containing the herpes simplex virus thymidine kinase TATA box and 5 tandem repeats of an IRF-1 and 2 binding sequence (AAGTGA).

2.5.3 Transcription factor expression plasmids

IRF-1 and IRF-2 expression plasmids pEFPlink2.IRF-1 and pEFPlink2.IRF-2 are based on the mammalian expression vector pEFPlink2 (Marais *et al.*, 1995) which was derived from pEFBos (Mizushima & Nagata, 1990) that uses the elongation factor 1 α promoter to direct high levels of expression of cloned cDNAs. These plasmids contain full length human cDNA encoding IRF-1 and IRF-2 respectively and were kindly provided by Dr. S. Goodbourn, (St. George's Hospital Medical School, Tooting, London).

The IRF-3 expression plasmid containing full length human cDNA encoding IRF-3 within the pcDNA3.1 backbone was provided by Oanh Dang

and M. David (USCD, CA). Similarly, pcDNA3.1-IRF-7 contains the full length human cDNA encoding IRF-7 and was kindly provided by S. Goodbourn, (St. George's Hospital Medical School, Tooting, London).

The IRF-9 expression vector pcDNA3-IRF-9.vp16 encodes a constitutively active form of IRF-9 comprising a hybrid cDNA containing the DNA binding domain of IRF-9 fused to the transactivator domain of the HSV1 VP16 protein and has been previously described (Kraus *et al.*, 2003). Briefly, cDNAs encoding full length IRF-9 and 80 C-terminal amino acids of the vp16 transactivator domain were produced by PCR. A second round of PCR was then used to generate IRF-9.VP16 hybrid cDNA inserts which were then cloned into the mammalian expression vector pcDNA3 (Invitrogen). This plasmid was kindly provided by Dr. C Horvarth (The Mount Sinai School of Medicine, New York, USA).

NF- κ B expression vectors pRc/CMV.p50 and pRc/CMV.p65 contain human p50 or p65 cDNAs respectively within the vector pRc/CMV and were a gift from Heike Pahl, University of Freiberg.

The CIITA expression vector pcDNA3.1-CIITA (1315) contains CIITA cDNA derived from human B-cells in the pcDNA3 backbone and was a gift from Prof. Cheong-Hee Chang, Indiana University School of Medicine, USA.

2.6 DNA manipulation techniques

2.6.1 Small scale preparation of plasmid DNA (miniprep)

Single colonies were picked from agar plates, added to 5 ml of LB containing the appropriate antibiotic in a 20 ml universal tube and incubated overnight in a 37°C orbital shaker at 2000 rpm. 1.5 ml of the culture was removed and the bacteria were pelleted for 2 minutes at 16,000g. Pellets were re-suspended in 100 µl of suspension medium and placed on ice for 5 minutes. 200 µl of lysis buffer was added; the suspension was mixed by inverting the tube 4-5 times and placed on ice for 5 minutes. 150 µl of potassium acetate buffer was then added. The solution was mixed by inverting and placed on ice for a further 5 minutes. The suspension was centrifuged for 5 minutes at 16,000g. The supernatant was carefully decanted into 500 µl of phenol/chloroform solution and mixed by vortexing. The mixture was centrifuged for 10 minutes at 16,000g and the top layer was pipetted off. The DNA was precipitated with 1 ml of ice-cold absolute ethanol and incubated at room temperature for 2 minutes. DNA was pelleted by centrifugation at 16,000g for 10 minutes, rinsed in 70% ethanol and air-dried for 10 minutes. The resulting DNA pellet was redissolved in 20 µl of TE and stored at 4°C prior to use.

2.6.2 Large scale preparation of plasmid DNA (maxiprep)

For a preparation of a large concentration of DNA plasmid, a small-scale culture was prepared as above and diluted 1:500 in LB containing an appropriate antibiotic for selection. Cultures were incubated overnight at 37°C in an orbital shaker. Bacterial cells were harvested and pelleted at

6000 *g* for 20 minutes at 4°C. Cell pellets were re-suspended in 10 ml of suspension medium and left at room temperature for 5 minutes. Twenty ml of lysis buffer was added; the suspension was mixed by inverting the pot 4-5 times and placed on ice for 5 minutes. Fifteen ml of potassium acetate buffer was added; the solution was mixed by inverting and placed on ice for a further 15 minutes. The suspension was centrifuged for 15 minutes at 15,000 rpm in Sorval GSA rotor and the supernatant was removed by filtration through muslin cloth. Supernatants were then added to an equilibrated Qiagen-tip 500 column (Qiagen, UK) and allowed to pass through the resin by gravity flow. The column was washed twice with Qiagen wash buffer and the DNA was eluted with 15 ml of Qiagen elution buffer. The DNA was precipitated with isopropanol and immediately centrifuged for 20 minutes at 15,000g. The pellet was rinsed in 70% ethanol, air-dried at RT and redissolved in 300 µl of TE. DNA concentrations were determined with an Ultrospec 2100 UV spectrophotometer (Pharmacia biotech, UK).

2.6.3 Purification of DNA fragments from agarose

DNA bands were purified using the Geneclean Turbo kit (BIO 101, USA) according to the manufacturers instructions. Briefly, DNA bands were visualised on a UV light source, excised and melted in an equal volume of an aqueous solution of a guanidine chaotropic binding salt and transferred to a DNA binding column. The column was washed twice with a salt solution containing ethanol and the DNA was eluted with RNase/DNase-free elution solution or RNase/Dnase-free dH₂O.

2.6.4 Restriction enzymes digestion

Digestion reactions were set up as follows: 2 µl 10X digestion buffer, 2 µl RNase, 0.5 µl of each enzyme, 100 ng DNA template, made up to a final reaction volume of 20 µl with dH₂O. DNA digestion was performed at 37°C in a water bath for at least 1 hr. All enzymes and enzyme reaction buffers were purchased from Promega.

2.6.5 Ligation

Insert:vector molar ratios of 3:1, 1:1 and 1:3 were used for ligation of SLA promoters (insert) into pGL3-basic (vector). Each ligation reaction consists of 50 ng of linearised dephosphorylated vector, SLA promoter insert, 1 µl of 10X ligation buffer, 1-3 U of T4 DNA ligase and distilled water to a final volume of 20 µl. Ligation reactions were performed overnight at 16°C. To calculate the appropriate amount of SLA promoter insert to include in the ligation reaction, the following equation was used;

$$\frac{\text{Ng of vector} \times \text{kb size of insert}}{\text{Kb size of vector}} \times \text{insert:vector molecular ratio} = \text{ng insert}$$

2.6.6 Transformation of competent bacteria

1-10 ng of plasmid DNA or 2 µl of ligation reaction was added to 50 µl of frozen *E. coli* JM109 (Promega) or TOP10 (Invitrogen) competent cells (thawed on ice). The mixture was flicked gently, placed on ice for 20 minutes and heat-shocked for 45 seconds at 42°C in a water bath. 950 µl LB broth was added to the mixture and the tube was incubated for 1.5 h at 37°C with

shaking. Cells were pelleted at approximately 1000 g for 10 minutes and re-suspended in 200 µl of LB broth. Finally, 50-150 µl of the mixture was plated onto an agar plate containing the appropriate antibiotic for selection and incubated overnight at 37°C.

2.6.7 Generation and small-scale preparation of Bacterial Artificial Chromosomes

Bacterial artificial chromosomes (BACs) 207G8 (AJ251829), 490B10 (AJ131112) and 493A6 (AJ251914) used in this study were kindly provided by Christine Renard and Patrick Chardon (LREG, INRA, France). Small scale production of these BACs is described in detail in (Chardon *et al.*, 2001). These BACs were derived from a genomic library comprising some 107000 clones spanning the entire haploid genome of a male, Large White pig homozygous for the MHC H01 haplotype, their generation has been described previously (Rogel-Gaillard *et al.*, 1999). Screening of the entire BAC library using primers designed for amplification of conserved regions of MHC class I sequences (exons 1 to 4) identified 18 BAC clones containing SLA class I genes, including those used in this study (Velten *et al.*, 1999). A precise map of the spatial arrangement of SLA class I genes was elucidated by sequence and southern blot analysis of these BAC clones which also confirmed the validity of the first physical map of the SLA class I region elucidated from a yeast artificial chromosome (YAC) library (Velten *et al.*, 1998).

2.7 Production of SLA class I promoter reporter constructs

2.7.1 Primer design

Sequence data for full-length promoter regions of each SLA class I gene (defined in Renard *et al.*, 2001) were downloaded from the Entrez Nucleotide website (National Centre for Biotechnology Information, U.S. National Library of Medicine, Bethesda, USA), and screened for restriction sites complimentary to those present in the multiple cloning region (MCR) of pGL3-basic using GCG software. Restriction sites that could potentially be used for cloning SLA class I promoters into pGL3 basic are summarised in table 1. *NheI* and *HindIII* sites were selected for cloning because they are absent from the coding sequence of each SLA class I promoter and are present in the MCS of the pGL3 basic vector. To enable directional cloning, forward primers and reverse primers were designed to incorporate *NheI* and *HindIII* sites respectively. In addition, three additional bases were included 5' to the restriction site to facilitate restriction enzyme interaction and digestion. Finally, Primer 3 software was used to screen each primer and primer pair in silico for hairpin loop formation, primer dimerisation and melting temperature. Primers were then purchased from MWG Biotech (Germany). Table 2 lists the primers used for PCR amplification of full-length promoter regions (from the upstream enhance silencer sites, to the ATG) for each SLA class I gene.

Table 1: Summary of restriction sites within SLA promoter

<i>Restriction site</i>	<i>Gene and position of restriction enzyme sites</i>							
	<i>SLA-1</i>	<i>SLA-2</i>	<i>SLA-3</i>	<i>SLA-5</i>	<i>SLA-6</i>	<i>SLA-7</i>	<i>SLA-8</i>	<i>MIC-2</i>
<i>KpnI</i>	-	-	1424	-	1430	151	-	347, 1728
<i>SacI</i>	-	1047	-	-	1620	-	-	1333
<i>MluI</i>	-	-	-	-	-	-	-	-
<i>NheI</i>	-	-	-	-	-	-	-	-
<i>SmaI</i>	-	-	1428	-	-	-	-	1724
<i>XhoI</i>	-	-	-	-	-	-	-	-
<i>BglI</i>	164	166	-	1165	-	-	-	-
<i>HindIII</i>	-	-	-	-	-	-	-	-

Table 2: Primers used for PCR amplification of SLA promoters

Primer	Target gene	5' to 3' Primer sequence	Anticipated product size (bp)
sla-1-NheI_F	SLA-1	GGGGCTAGCAAGGAAAATCATTCTCTGAGC	1069
sla-1-HindIII_R	SLA-1	GGGAAGCTTGATCCTCAGCCTCGGGGTCT	
Promsla-2b-NheI_F	SLA-2	CTAGCTAGCTACATACCAAACTCCT	1330
Promsla-2b-HindIII_R	SLA-2	CCCAAGCTTTGCGTGGAGAGTTATAA	
sla-3-NheI_F	SLA-3	GGGGCTAGCATACTTTAATAGTATCTTCTATG	1402
sla-3-HindIII_R	SLA-3	AGAAAGCTTTACCAGTCAGGGCCAGGG	
sla-5-NheI_F	SLA-5	GGGGCTAGCGTTTAAATAGTATCTTCTCTG	1093
sla-5-HindIII_R	SLA-5	AAAAAGCTTCCCGAGAGCAGCAGGAAGAGG	
Promsla-6b-NheI_F	SLA-6	CTAGCTAGCAGAGTCACAGTTTGGGAA	988
Promsla-6b-HindIII_R	SLA-6	TTTAAGCTTCTGAGTTTGGCTGGTCTG	
Promsla-7b-NheI_F	SLA-7	CTAGCTAGCTCCCTGTCTCTGGTGTGA	997
Promsla-7b-HindIII_R	SLA-7	TTTAAGCTTAACGCACCCTCGGGGT	
Promsla-8a-NheI_F	SLA-8	CTAGCTAGCGCTGCCTCCATAATGATC	962
Promsla-8a-HindIII_R	SLA-8	TTTAAGCTTGACCGAATCCTGTGGGT	
promMIC2-NheI_F	MIC2	CTAGCTAGCAACCCATGGGATACTGATTC	1228
promMIC-2-HindIII	MIC2	ATAAAGCTTGCCAGAAACGACAGGTAACGG	

2.7.2 Polymerase chain reaction

For gradient PCR, reaction mixtures were set up as follows: 10 µl 10x High Fidelity Buffer (Eppendorf), 2 µl 10 mM dNTPs, 1 µl 200µM forward primer, 1 µl 200µM reverse primer, 80 ng BAC DNA template and 0.5 µl Triple Master Enzyme (contains Taq polymerase and *pfu* Eppendorf), made up to a final reaction volume of 100 µl with filtered water. The reaction

mixture was divided into 10 samples and placed in a Eppendorf thermocycler.

Optimum PCR annealing temperatures for each primer pair were determined by performing PCRs across a temperature gradient of 45 to 72°C using a gradient thermocycler (Eppendorf). PCR product size was determined by gel electrophoresis on a 0.8% (w/v) agarose gel. Optimum annealing temperatures, defined by PCR product size and yield, were used in all subsequent PCR reactions. PCR products were extracted from agarose gels and subcloned into pcDNA3.1/V5-His/TOPO (Invitrogen) as a shuttle vector. The insert was excised with *NheI* and *HindIII* and analysed by gel electrophoresis to confirm the size of SLA promoter inserts.

The following PCR cycling parameters were used:

Denature 94°C for 30 sec

Denature 94°C for 30 sec

Anneal at gradient temperature for 30 sec

Elongation for 2 minutes

Final elongation at 72°C for 10 minutes

} 30 cycles

After completion of the program, PCR products were mixed with 6X gel loading dye and the resulting mixture was run on a 1% (w/v) agarose gel. The DNA fragments were visualised by ethidium bromide staining and viewed on a UV light source. Appropriate PCR products were excised from gels and isolated as described below.

2.7.3 Sub-cloning of SLA class I promoters

SLA class I promoter sequences were amplified by PCR and isolated as described above. First, PCR products were cloned into the T-tailed vector pcDNA3.1/V5-His/TOPO (Invitrogen) following the manufacturer's protocol. Promoters inserts were then excised with *NheI* and *HindIII* and analysed by gel electrophoresis on a 0.8% (w/v) agarose gel to confirm the size of each insert. The DNA containing each SLA promoter was extracted and purified as above and ligated into the promoter-less vector pGL3-basic (Promega, UK) upstream of the firefly luciferase gene. Plasmids were then analysed by digestion with *NheI* and *HindIII* followed by gel electrophoresis to re-confirm the size of promoter inserts. The sequences of SLA promoter inserts within pGL3-basic were confirmed by DNA sequencing. Table 3 summarises the SLA promoter constructs generated in this study.

Table 3: Summary of the SLA promoter reporter constructs produced

Construct name	SLA promoter insert	insert size
pGL3-SLA1	SLA-1	1069
pGL3-SLA2	SLA-2	1330
pGL3-SLA3	SLA-3	1402
pGL3-SLA-6	SLA-6	988
pGL3-SLA-7	SLA-7	997
pGL3-MIC-2	MIC-2	1228

2.7.4 Sequencing

2.7.4.1 Sequencing primers

pGL3-SLA promoter constructs were sequenced with the forward and reverse primers specific for individual SLA class I promoters listed in table 2. Primers specific for pGL3-basic were also used, these were as follows:

forward primer RVfor 5' -CTAGCAAATAGGCTGTCCC- 3'

reverse primer GLrev 3'-CTTTATGTTTTTGGCGTCTTC-5'.

2.7.4.2 Sequencing reaction

Sequencing templates were approximately 350 ng of small scale preparation of plasmid DNA diluted in 10 µl of deionised water. The DNA was heat-treated for 1 minute at 96°C. The samples were prepared using the Dye Terminator Cycle Sequencing Quick-Start kit (Beckman Coulter, USA). The following reagents were added to the DNA sample: 3.2 pmoles of sequencing primers, 4 µl of reaction buffer, 4 µl of Quick-start solution and dH₂O to a final reaction volume of 20 µl. The following cycling parameters were used:

Denature 96°C for 20 sec	}	30 cycles
Anneal 50°C for 20 sec		
Polymerise 60°C for 4 minutes		
Hold at 4°C		

The PCR products were then precipitated with 2 volumes of 3 M NaOAC, pH 5.2. The DNA pellet was cleaned with 3 washes of 70% ethanol. Forty µl of sample loading solution was added and samples were analysed with an automatic capillary sequencer CEQ 8800 Genetic Analyses System (Beckman Coulter, USA). Sequence trace data were analysed using Chromas (version) and compared with the reference BAC sequences using DNAsp (version and programme name). A minimum of two clones of each construct were sequenced four times.

2.7.5 Computational analysis of DNA sequence data

Sequence analysis was carried out using components of the Wisconsin Package version 10.1 (Genetics Computer Group (GCG), Madison, USA). Sequence alignments were carried out using DNAsis software using standard multiple sequence alignment criteria. Properties of primers such as predicted annealing temperature, hairpin loop formation and internal purine-pyrimidine bond formation were assessed using the predictive software Primer3 (Rozen & Skaletsky, 2000). The program TFSEARCH version 1.3 searches highly correlated sequence fragments against TFMATRIX transcription factor binding site profile database in the TRANSFAC databases by GBF-Baunschweig (Heinemeyer *et al.*, 1998). This program was used to search for transcription factor binding sites in SLA class I promoters.

2.8 Cell culture techniques

2.8.1 Standard transfection protocol

Cells were grown to 70-80% confluency (either on 24 well plates or on Petri dishes) and were transiently transfected with plasmid DNA using a liposomal "Lipofectamine" transfection system (Invitrogen, UK). Lipofectamine was used at a ratio of 1 μ g plasmid DNA to 2.5 μ l lipofectamine (the optimum ratio for pGL3-SLA promoter constructs were determined experimentally, data not shown) and according to the manufacturer's protocol. For transfections on 24 well plates, 1.25 μ l of Lipofectamine and 25 μ l of serum-free IMDM were mixed with 0.5 μ g plasmid in 25 μ l of serum-free IMDM (per plasmid per well). The mixture was vortexed and left for 45 minutes at RT. In the meantime, cells were washed three times with serum-free IMDM. 50 μ l of each plasmid DNA/Lipofectamine/IMDM mixture was then added to each well and topped up to a final volume of 400 μ l with serum free IMDM. Cells were incubated with the transfection mixture at 37°C for one hour. Next, 400 μ l IMDM supplemented with 20% FCS was added to each well and cells were incubated for a further 48 hrs (unless stated otherwise). For cytokine stimulation experiments, cells were transfected as above and incubated in IDIM for 32 hours. Cytokine was then added at the appropriate dilution for the time indicated in each experiment, usually 16 hours. Cells were then harvested and processed for beta-galactosidase and luciferase assays as standard.

2.8.2 Co-transfections with transcription factor expression plasmids

The optimum amount of transcription factor expression vectors to be co-transfected with pGL3-SLA promoter constructs was determined by titration of each transcription factor expression vector against luciferase activity driven by the SLA-1 and SLA-7 promoters. A combination of 0.1 µg transcription factor expression vector, 0.4 µg pcDNA3 was found to be optimal, greater amounts of transcription factor expression vector were found to inhibit SLA-1 driven luciferase activity. Max cells were transiently transfected with 0.5 µg of SLA promoter constructs, 0.5 µg pJat-lac, 0.4 µg of pcDNA3 and 0.1 µg of one of the following expression vectors: pEFPlink2.IRF-1, pEFPlink2.IRF-2, pcDNA3-IRF-3, pcDNA-IRF-7, pcDNA3-IRF-9.vp16, pRc/CMV.p50, pRc/CMV.p65 or pcDNA3.CIITA(1315). For control experiments, 0.5 µg of the following control plasmids was used: pGL3-control, p(9-27 ISRE)4tkΔ(-39)Lucifer, p(IRF-1*GAS)6tkΔ(-39)Lucifer, pkB6tkluc, p[(AAGTGA)4]5tkΔ(-39)Lucifer. Transfections were carried out as above, using plasmid DNA and Lipofectamine at a ratio of 1:2.5, see figure 2.1 for a summary.

Figure 2.1: Protocol for co-transfection of Max cells with multiple plasmids

Typically co-transfection experiments included 0.5 μ g of SLA promoter reporter construct, 0.5 μ g pJatLacZ control, 0.1 μ g of transcription factor expression plasmid and 0.4 μ g of a backbone plasmid corresponding to that of the latter. Each plasmid was incubated with lipofectamine and serum free media media at a plasmid DNA to lipofectamine ratio of 1.25 as shown above.

2.8.3 Luciferase and β -galactosidase assays

For transfections carried out in 24 well plates, 80 μ l of lysis buffer (Promega) was added to each well. Cells were scraped and lysates were transferred into 1.5 ml tubes and vortexed briefly. Supernatants were transferred into 1.5 ml tubes and placed on ice prior to use in luciferase and β -galactosidase assays. 20 μ l of cell lysate was used for both types of reporter assay. Luciferase assays were performed according to manufacturers instructions and luciferase activity was measured using a Wallac 1205 Betaplate luminometer (PerkinElmer lifesciences, Pharmacia).

For β -galactosidase assays, cell lysates were diluted in 0.5 ml of Z buffer, vortexed briefly and 200 μ l of substrate buffer (o-nitrophenyl- β -D-galactopyraniside) was added. Samples were vortexed briefly and incubated for 1 hour at 37°C. 200 μ l 1M Na₂CO₃ was added to inactivate β -galactosidase enzyme and stop the reaction. 200 μ l of sample was then transferred to a 96 well plate and absorbance was read at 420 nm using a Labsysteme Multiscan plus plate reader spectrometer (Labsystems, UK).

2.8.4 Cytokine treatments

Porcine interferon alpha (pIFN- α): For FACS and transfection experiments d/d PAECs or Max cells were incubated with 200 U/ml of recombinant porcine IFN- α (PBL Biomedical labs, USA) in RPMI or IMDM respectively, for 16 hrs at 37°C.

Porcine interferon gamma (pIFN- γ): d/d PAECs or Max cells were incubated with 0.2 μ g/ml recombinant porcine IFN- γ (Biosource, USA) in RPMI or IMDM respectively, for 16 hrs at 37°C.

Tumour necrosis factor alpha (TNF- α): d/d PAECs or Max cells were incubated with 0.5 μ g/ml recombinant human TNF- α (R&D Systems, USA) in RPMI or IMDM respectively, for 12-16 hrs at 37°C. TNF- α is referred to as hTNF- α in the text.

2.9 Indirect Immunohistochemistry

2.9.1 Indirect immunofluorescence

PAECs and Max were cultured on 13-millimetre glass coverslips in RPMI or IMDM respectively to 70-80% of confluency. Cells were washed with PBS three times and fixed with 100% methanol for 10 minutes or 4% PFA for 10 minutes with shaking at RT. Cells were rinsed with PBS three times and blocked in blocking buffer for 20 minutes with shaking at RT. Cells were rinsed three times in PBS then incubated with the primary antibody diluted in blocking buffer for 20 minutes with shaking. See table 4 for dilutions used for each primary antibody. Cells were then washed in PBS and incubated with the Alexa⁴⁸⁸ or Alexa⁵⁹⁴ conjugated secondary antibody at 1:500 dilution of a 1 mg/ml stock in blocking solution for 20 minutes in the dark. Cells were washed three times in PBS before being dyed for 5 minutes with a 1:20,000 dilution of a 5 mg/ml stock of DAPI in PBS. Finally cells were washed in

distilled water and coverslips were mounted onto Superfrost slides (BDH, UK) using Fluoromount G (Southern Biotechnology, USA).

2.9.2 Indirect immunofluorescence analysis of MHC class I

PAECs and Max were cultured on 13-millimetre glass coverslips in RPMI or IMDM respectively to approximately 80% of confluency. For surface staining, cells were washed three times in PBS and incubated with detergent free blocking buffer for 20 minutes at RT°C with shaking. Cells were then incubated with PT-85 (1 µl in 1000 µl of blocking buffer) for 30 mins at RT°C. Cells were then washed in PBS and incubated with the Alexa⁴⁸⁸ or Alexa⁵⁹⁴ conjugated secondary antibody at 1:500 dilution of a 1 mg/ml stock in blocking solution for 20 minutes in the dark and then fixed in acetone/methanol for 2 minutes. Cells were washed three times in PBS-Tween-20 then incubated with DAPI (1:20,000 dilution of a 5 mg/ml stock) for 10 minutes to stain nuclear DNA. Finally cells were washed in distilled water and coverslips were mounted onto Superfrost slides (BDH, UK) using Fluoromount G (Southern Biotechnology, USA).

For internal staining of MHC class I, cells were washed in PBS three times then fixed for 60 seconds with methanol/acetone. Cells were incubated with blocking buffer (containing 0.5% NP-40) for 20 minutes at RT°C then permeabilised with 1% (v/v) NP40 at 37°C for 15 minutes. Cells were incubated with PT-85 (dilution as above) for 45 minutes at RT°C, washed three times in PBS-Tween 20 and incubated with goat anti-mouse Ig

conjugated to Alexa⁴⁸⁸ (1:500 in blocking buffer containing 0.5% NP40) for 30 minutes in the dark. Cells were then washed three times and nuclear DNA was stained with DAPI and cells were mounted on glass slides as above.

2.9.3 Image analysis

Preparations were viewed at 60X/1.4 NA with a Nikon E800 microscope and images were captured with a Hamamatsu C-4746A CCD camera. Captured images were analysed using Improvision Openlab 3.1 software (Improvision, UK).

2.9.4 Analysis of MHC class I surface expression by FACS

Following virus infection, cytokine or control treatment for the time period indicated for each experiment, cells were washed in PBS, incubated with 2mls of versine/trypsin until detached and collected by gentle centrifugation. Cell pellets were resuspended in FACS diluent, allowing 25 µl of cell suspension per sample. Twenty five µl of cell suspension was added to each well of a 96 well, round bottomed plate as required. To wash the cells, plates were centrifuged at 500g for 2 minutes at 4°C, supernatants disposed of and pellets dispersed by gentle shaking. Cells were incubated with 25 µl of primary antibody at an appropriate dilution (given in table 4) and 150 µl of FACS diluent in darkness at 4°C for 20 minutes. Control samples for autofluorescence and non-specific binding of secondary antibody were incubated with 25 µl of FACS diluent in place of primary antibody. Cells were

washed twice as above and 25 µl of phycoerythrin-conjugated goat anti-mouse IgG_{2a}, 2b, or IgM (Southern Biotechnology Associates) was added to each well, including control samples for non-specific binding of secondary antibodies, to detect bound primary antibodies. 150 µl of FACS diluent was added to each well and plates were incubated at 4°C in darkness for 20 minutes. Finally, cells were washed twice, re-suspended in 100 µl FACS fixative, sealed and stored at 4°C prior to analysis. Cells surface MHC class I expression was analysed by FACS (FACScan, Becton Dickson) using the CellQuest program. The degree of cell membrane MHC class I expression was expressed as mean fluorescence intensity (MFI).

Table 4: table of antibodies and dilutions used in FACS experiments

Antibody	Specificity	Ig subclass	Dilution
PT-85	MHC class I	IgG2a	1:1000
2.27.3	SLA class I	IgG2a	Neat
74.11.10	SLA class I	IgG2b	Neat
2.32.1	SLA class I, d/d haplotype	IgG2b	Neat
2.12.3	SLA class I, d/d haplotype	IgM	Neat
16.7E4	SLA class I, c/c haplotype	IgM	Neat

2.10 Statistical analyses

All statistical analyses were performed using the statistical software package SPSS version 10.4 for Windows according to the manufacturers instructions.

Chapter 3: Cellular characterisation of SLA class I expression in porcine cells.

3.1 Introduction

In the pig, the polymorphic nature of MHC class I and the variation in allele expression by individuals makes studying cell surface expression of MHC class I complicated. However, characterisation of SLA class I expression is simplified with the use of the NIH inbred mini pig, which are homozygous on both alleles for the whole SLA locus (Pennington et al., 1981). In addition, several mouse antibodies that are specific for pig SLA have been produced, including antibodies that can distinguish the d/d haplotype from other haplotypes. The principle aims of this chapter were to explore the tools available for studying SLA class I, namely antibodies and porcine cell lines and to use these tools to characterise SLA class I expression on a series of porcine cell lines and in response to cytokine stimulation and viral infection. In addition, this study aimed to expand the technical applications of the antibodies available for SLA class I studies.

In order to address these aims, firstly cell lines from NIH d/d inbred pigs which express high levels of SLA class I were isolated. Secondly, a panel of SLA class I specific antibodies were used to investigate surface and intracellular expression of SLA class I on these cells, using FACS and immunofluorescence. Finally, changes in surface expression in response to cytokine stimulation or infection with a pathogen, classical swine fever virus,

were investigated. In the human, mouse and pig it is well established that MHC class I expression is induced by interferon and pro-inflammatory cytokines. It has been shown that modulation of MHC class I expression is an important immune evasion mechanism for many viruses. In this study, the porcine virus CSFV was used to determine whether infection can modulate expression of SLA class I. There has been little work done on the mechanism of inhibition of SLA class I expression by a pig virus. This has important implications for vaccine design and mechanism of T cell killing in the pig. Isolation of primary cells from pigs also provides a host cell model to study field strains of virus, which may not replicate in cultured cell lines and since these cells are highly differentiated, they should rapidly respond to stimulation by cytokines or viruses.

3.2 Expression of SLA class I on primary cells

3.2.1 Porcine aortic endothelial cells and responses to interferon

Porcine aortic endothelial cells (PAECs) were isolated from the aortas of both Landrace cross Large White pigs of unknown haplotype (referred to herein as 'outbred' PAECs) and from NIH minipig inbred at the MHC locus, expressing the d/d (H04) haplotype (referred to herein as 'd/d' PAECs). PAECs were incubated with pIFN- α (200 u/ml for 16 hours) or media alone, harvested and incubated with a panel of antibodies specific for MHC class I. The panel of antibodies (listed in table 4) comprises a series of antibodies raised in mice against SLA class I including locus-specific antibodies for the H04 haplotype (d/d). Binding of each antibody to SLA class I was detected

with isotype specific goat anti-mouse immunoglobulin antibodies conjugated to phycoerythrin. In control experiments for non-specific binding, PAECs were incubated with goat-anti mouse immunoglobulin isotype specific antibodies alone. SLA class I expression was then examined by FACS for each antibody and condition. Figure 3.1 shows FACS profiles of SLA class I expression on outbred and d/d PAECs from a single experiment, representative of several repeats. For each antibody, histograms representing constitutive (black histograms) and IFN- α induced (red histograms) SLA class I expression are compared. Constitutive SLA class I expression was detected on outbred PAECs by PT-85 and 2.27.3 and on inbred PAECs by PT-85, 2.27.3 and 2.12.3 antibodies. PAECs from d/d pigs express higher constitutive levels of SLA class I than PAECs from outbred pigs. Interestingly, the antibody 2.12.3 did not detect SLA class I on outbred PAECs (compared to isotype control), but did detect SLA class I expression on d/d PAECs, showing that the epitope for this antibody is specific for the d/d haplotype and either not a common epitope among outbred pigs or not expressed by the individual examined.

Interferon induced MHC class I to a greater extent on the cell surface of inbred PAECs compared to outbred PAECs. Interestingly, interferon had no effect on the PT-85 or 2.12.3 epitopes on outbred cells and caused only a marginal increase on the epitope recognised by 2.27.3. However, on inbred pig cells, interferon induced an increase in the SLA epitopes recognised by each antibody tested, especially that of 2.12.3, where the MFI increased by 3 logs. Serological analyses have deduced that 2.12.3 has an exclusive

Figure 3.1: Comparison of SLA class I expression and IFN responses on PAECs derived from inbred and outbred swine.

PAECs were isolated from the aortas of an outbred pig and an inbred NIH mini pig expressing the d/d haplotype. Cells were incubated with media alone (black histograms) or pIFN- α (red histograms) for 16 hours, harvested and incubated with antibodies specific for MHC class I: PT-85, 2.27.3a and 2.12.3. Antibody binding was detected with goat anti-mouse isotype specific antibodies conjugated to phycoerythrin and cells were analysed by flow cytometry (FACScan, Beckman). Controls for autofluorescence and non-specific secondary antibody binding were also included. In each experiment the secondary antibody control did not exceed a mean fluorescence intensity of 10 (data not shown). This figure shows a single experiment, representative of at least three repeats. Cells were untreated (black histograms) or cultured in the presence of IFN- α for 16 hours (red histograms).

specificity for the H04 (d/d) haplotype. These results also suggest that interferon induction of SLA class I can be masked in cells expressing a mixed population of SLA molecules. There appears to be haplotype or allele specific interferon induction, indicating that some alleles of SLA class I are more sensitive to interferon induction than others. For this reason, d/d PAECs should be used in future studies.

The antibodies PT-85 and 2.27.3 were then used to detect SLA class I by indirect immunohistochemistry. The antibody 2.12.3 was not used for two reasons; results from this study demonstrated that MHC class I is not always detected by 2.12.3 on resting cells by FACS and secondly, this antibody is an IgM and therefore has limited technical applications. Since the antibodies PT-85 and 2.27.3 have not been used for this technique previously, a technique to analyse cell surface and intracellular SLA class I by indirect immunohistochemistry required development. Preliminary experiments showed that PT-85 and 2.27.3 do not recognise SLA class I on cells that had been fixed with methanol or paraformaldehyde. Therefore, surface MHC class I was stained with PT-85 or 2.27.3 on unfixed PAECs at room temperature. Cells were incubated with each of these antibodies, washed in PBS then incubated with a goat antibody conjugated to Alexa⁴⁸⁸ specific for mouse IgG to detect antibody binding to MHC class I and then fixed. For intracellular staining of SLA class I cells were fixed (60 secs) with methanol and acetone, washed and permeabilised with 0.5% NP40 and stained with PT-85 or 2.27.3. Figure 3.2 shows SLA class I detected by PT-85 at the cell surface (panel A) and inside the cell (panel B). Intracellular class I was found

Figure 3.2: Cell surface and intracellular MHC class I expression in PAECs

A technique for analysing SLA class I expression by indirect immunofluorescence was developed. For cell surface staining (panel A), PAECs were incubated with PT-85, a mouse monoclonal antibody specific for MHC class I in the absence of detergent. Antibody binding to SLA class I was detected with goat anti-mouse antibody conjugated to Alexa⁴⁸⁸ (green). Cells were then fixed in Methanol/Acetone (1:1) and cellular DNA was labeled with DAPI stain (blue). For intracellular staining (panel B), PAECs were briefly fixed with metanol/acetone, incubated for 15 minutes with 0.5 % NP40 at 37°C and incubated with PT-85. Nuclear DNA was stained with DAPI (blue). Cells were viewed at 100 (A) or 60X (B) with a Nikon E800 microscope and images were captured with a Hamamatsu C-4746A DCC camera as described in the materials and methods. Internal SLA class I is located in peri-nuclear vesicles presumed to be within the Golgi and ER.

as anticipated in a peri-nuclear location within vesicular structures that resemble the endoplasmic reticulum or golgi, staining for ER/golgi markers would be required to confirm this. Using this method, cell surface class I could not be detected by PT-85, presumably because chemical fixation denatures surface class I. The antibody 2.27.3 produced very similar results to PT-85 and therefore is not shown.

3.2.2 Primary pig lymphoma cells (Shimozuma cells) and responses to cytokine induction

Shimozuma cells are a swine malignant lymphoma-derived cell line in which SLA class I expression has not previously been characterised. Shimozuma cells were cultured in the presence of cytokines hTNF- α , pIFN- α or pIFN- γ for 16 hours, harvested, incubated with antibodies specific for MHC class I (PT-85, 2.27.3, 74.11.10 and 2.12.3) and analysed by FACS. Constitutive expression of MHC class I on Shimozuma cells was not detected by any of the antibodies tested (data not shown). Interestingly, this primary B-cell line, which might have been expected to express SLA class I, have lost this surface expression, a common characteristic of transformed cells which underlines the value of using primary cell line in these studies.

3.2.3. Primary Max pig kidney cells and responses to interferon

Max cells isolated from the kidney of a NIH d/d mini pig and immortalised by transfection with SV40 were a gift from A. Saalmuller (Federal Research Centre for Virus Diseases of Animals, Germany). These cells have the added advantage that they survive in cell culture for 20 passages before SLA class I cell surface expression is lost, compared to PAECs that can only be used for 5-10 passages (data not shown). In order to determine constitutive and interferon-induced SLA class I expression, Max cells were stained with a panel of antibodies recognising inter-species, porcine-specific or d/d haplotype specific antibodies PT-85, 2.27.3 and 2.12.3 antibodies described above and one additional antibody, 2.32.1 and analysed by FACS. Figure 3.3 shows that Max cells express high constitutive levels of SLA class I, which was detected by PT-85, 2.27.3 and 2.32.1 but not 2.12.3 (open black histograms). These results support the findings for PAECs from the d/d haplotype.

Max cells were treated with either pIFN- α (figure 3.3 left panel) or pIFN- γ (right panel) for 16 hours and processed for FACS analysis as above. Both IFN- α (red histograms) and IFN- γ (blue histograms) increase cell surface SLA class I expression. Expression of the epitope recognised by PT-85 is more weakly induced by IFN- α / γ than the epitopes recognised by 2.27.3 and 2.32.1. As seen in outbred PAECs, SLA class I recognised by 2.12.3 is not constitutively expressed on Max cells and like inbred PAECs, a large increase in expression is seen after treatment with IFN- α / γ . Overall,

Figure 3.3: MHC class I expression is up regulated by IFN- α and IFN- γ on Max cells .

Max cells were grown in the presence of IFN- α (red histograms) or IFN- γ (blue histograms) or media alone (open black histograms) for 16 hours, harvested and incubated with mouse antibodies specific for a highly conserved element of MHC class I (PT-85), SLA class I (2.27.3) or d/d haplotype SLA class I (2.32.1 and 2.12.3). Solid grey histograms represent isotype controls used for each antibody and unstimulated Max cells are shown as open black histograms. These data show a single experiment but are representative of trends observed in multiple repeats.

SLA class I was induced to a greater extent by IFN- γ than IFN- α . This suggests that IFN- γ , which is secreted by activated T cells, is a more potent inducer of cellular responses such as IRF activation than IFN- α .

The immunofluorescence microscopy procedure developed above was used to detect surface SLA class I expression on Max cells and determine the effect of IFN- α /- γ on SLA class I expression. Max cells were grown on 13 mm glass coverslips to 70-80% confluency and incubated with pIFN- γ in media or media alone for 16 hours. Cell surface MHC class I was then stained with PT-85 antibody as described above. In figure 3.4, constitutive expression of MHC class I on the cell surface of Max cells can be visualised by immunofluorescence microscopy (panel A, in green), cellular DNA is labelled with DAPI (blue). An increase in cell surface expression of MHC class I was detected with PT-85 antibody following IFN- γ treatment (panel B, in green). These results demonstrate that indirect immunofluorescence microscopy can be used to monitor gross changes in cellular expression of MHC class I.

Figure 3.4: IFN- γ induces cell surface expression of MHC class I.

Max cells were grown at 37° on cover slips for 24 hours then incubated with media alone or pIFN- γ for a further 16 hours. Cell surface MHC class I was stained with PT-85 followed by Alexa⁴⁸⁸ conjugated goat anti mouse antibody (green). Cells were then fixed in acetone/methanol (1:1 v/v) and cellular DNA was labelled with DAPI (blue). Cells were viewed at 60X/1.4NA with a Nikon E800 microscope and images were captured with a Hamamatsu C-4746A DCC camera as described in the materials and methods. This figure shows un-stimulated Max cells (panels a to c) and IFN- γ stimulated (panel d to f).

3.3 Effect of CSFV infection on MHC class I expression in porcine cells

The continuous interactions between host and pathogens during their co-evolution have shaped the immune system, but also the counter measures used by pathogens. Down-regulation of cell surface MHC class I expression is a common immune evasion strategy employed by viruses to avoid recognition by CTLs. Selective disruption of specific MHC class I alleles also allows a virus to avoid NK cell lysis. CSFV causes a persistent infection in pigs and therefore must be able to circumvent detection by the immune system. This thesis investigated whether CSFV infection affects MHC class I expression. Since CSFV infection of Max cells has not been previously reported, it was first necessary to determine if Max cells could support CSFV infection and replication and be used in such an investigation.

3.3.1 Max cells can be infected with field isolates of Classical Swine Fever Virus.

Max cells were infected with the Brescia strain of CSFV for 24 hours, fixed with methanol and incubated with a mouse monoclonal antibody specific to the E2 protein of CSFV as a marker of cellular infection. Nuclear DNA was stained with DAPI and uninfected cells were used as a negative control. Figure 3.5 shows that Max cells infected with CSFV express the CSFV E2 protein in the cytosol (green). Uninfected cells were negative for E2. Immunofluorescence staining of E2 protein was used as a positive control for virus infection in FACS experiments. This was important for SLA class I expression studies in CSFV infected cells (see below), since the E2

Figure 3.5: Max cells can be infected with field strains of CSFV

Max cells were incubated at 37°C with the Brescia strain of CSFV for 24hrs. Control cells were uninfected. Cells were fixed with methanol and incubated with a mouse monoclonal antibody specific for the E2 protein of CSFV. Antibody binding to E2 was detected using a goat antibody specific for mouse Ig conjugated to Alexa⁴⁸⁸ (green). Cells were viewed at 60X/1.4NA with a Nikon E800 microscope and images were captured with a Hamamatsu C-4746A DCC camera as described above. Cellular DNA was labeled with DAPI (blue) in uninfected control cells (A) and cells infected with CSFV (B). The E2 protein of CSFV can be detected in the cytosol of infected Max cells (B) but is absent in uninfected cells (D).

antibody cannot be easily used in FACS analysis for double staining with 2.27.3 or PT-85 as it is also a mouse IgG_{2a} antibody.

3.3.2 CSFV down-regulates constitutive SLA class I surface expression.

Next, the effect of CSFV infection on cell surface SLA class I expression was investigated. Max cells were infected with CSFV Brescia for 24, 48, 72 or 96 hours, control cells were incubated with media alone. Cells were harvested, incubated with the panel of antibodies listed in table 4 and cell surface expression of SLA class I was then measured by FACS. Antibody binding to MHC class I was detected with an appropriate isotype control conjugated to PE and as in previous experiments, a conjugate control was included for non-specific binding of these secondary antibodies. As a control for CSFV infection, cells were stained for E2 expression by immunofluorescence as described above.

In the experiments depicted in figure 3.6, 100% of cells were positive for CSFV E2. Graphs show data generated from a series of repeats of the experiment described above and individual graphs represent the mean average percentage decrease in MHC class I expression, obtained from pooled mean fluorescence intensity data, at each viral infection time point compared with uninfected control cells (normalised to 100%). Error bars represent the standard deviation from this mean percentage decrease. Table 5 summarises the raw data (MFIs) from these experiments including the mean MFI $\pm$ standard deviation. Finally, Mann U Whitney tests were performed on unpaired raw (MFI) data to determine statistical significance.

Figure 3.6: CSFV down-regulates MHC class I surface expression on Max cells.

Cell surface MHC class I expression on Max cells was measured at 24, 48, 72 and 96 hpi with CSFV Brescia by flow cytometry using a panel of antibodies specific for MHC class I. Graphs show the percentage of class I expression relative to an uninfected control (normalised to 100%) at each time point obtained with antibodies PT-85, 74.11.10 and 2.27.3. Error bars indicate standard deviation from the mean relative percentage. Mann U Whitney tests were performed on unpaired raw data, significant ($p < 0.05$) results are indicated by an asterisk. A significant decrease in MHC class I expression was detected with PT-85 (blue). 74.11.10 (red) showed a trend of down-regulation whereas no decrease in MHC class I expression was detected using the antibodies 2.27.3 (yellow) or 2.12.3 (data not shown). Raw MFI data are summarised in table 5 below.

Antibody	Uninfected	24 hpi	48 hpi	72 hpi	96 hpi
PT-85	180.1 ± 45.5 n=5	104.4 ± 28.1 n=5	95.8 ± 31.8 n=5	79.7 ± 28.1 n=4	122.4 ± 26.6 n=4
2.27.3	63.3 ± 73.1 n=5	45.4 ± 44.6 n=5	52.5 ± 52 n=5	46.1 ± 41.3 n=4	52.7 ± 36.7 n=4
74.11.10	86.6 ± 58.8 n=6	56.5 ± 30.6 n=6	56.5 ± 46.6 n=6	39.0 ± 21.8 n=5	53.3 ± 25.9 n=4
2.12.3	10.7 ± 6.9 n=6	5.3 ± 1.9 n=6	6.5 ± 3.0 n=6	6.5 ± 3.38 n=5	6.8 ± 2.7 n=4
16.7E4	3.4 ± 0.4 n=6	3.5 ± 0.3 n=6	3.3 ± 0.5 n=6	4.1 ± 0.6 n=5	5.135 ± 3.0 n=4

Table 5: Summary of raw MFI data

The table shows the mean average MFI recorded for each antibody, at each time point ± standard deviation and the number of experimental repeats represented by these data (n).

In each sample the MFI of the isotype controls and the negative control 16.7E4 did not exceed 10 (data not shown) at all time points. The data presented shows that CSFV significantly inhibits MHC class I expression detected by PT-85 (blue) at 24 hpi ($p = 0.009$), 48 hpi ($p = 0.009$) and 72 hpi ($p = 0.014$). Overall, the decrease in MHC class I expression progresses with viral infection. However at 96 hpi, the magnitude of the decrease in MHC class I expression is less than at earlier time points. The results using the antibody 74.11.10 show a similar trend; MHC class I expression appears to decrease, although large standard deviations prevented statistical significance being reached. Similarly, the mean percentage decrease in class I expression recorded with the antibody 2.27.3 (yellow) shows a trend of down regulation. In this instance significance could not be achieved, presumably the large variations in standard deviation at each time point is responsible for this.

By contrast, the antibody 2.12.3 did not record an increase or decrease in SLA class I expression (results not shown) and the MFI of the negative control 16.7E4 remained below 10.

Figure 3.7 shows a series of representative histograms showing data from one repeat of this experiment. FACS profiles of isotype controls (black histograms), uninfected control cells (open black histograms) and CSFV infected Max cells are shown for each antibody and at each time point.

The interesting result from these experiments is that CSFV infection down-regulates SLA class I expression significantly and therefore these results describe a previously undocumented immune evasion strategy used by this virus. The exact identity of the alleles that are down-regulated following CSFV infection could not be determined, since the identity of the specific class I alleles recognised by the panel of antibodies used is not known.

Figure 3.7: CSFV down regulates MHC class I expression on Max cells - representative histograms.

Max cells were infected with the Brescia strain of CSFV for 24, 48, 72 and 96 hpi (as indicated) Figure 6A: CSFV down-regulates MHC class I surface expression on Max cells. Cell surface MHC class I expression was measured by flow cytometry using a panel of antibodies specific for MHC class I. This figure shows the individual results from each antibody recognising MHC class I and is a representation of a single experiment from figure 6A. Each graph shows an isotype control (solid black histogram), MHC class I on uninfected Max cells (open black histogram) and MHC class I on CSFV infected cells (open red histograms). This figure is continued overleaf.

Figure 3.7: CSFV down regulates MHC class I expression on Max cells continued

3.4 Discussion

3.4.1 Summary of results

This chapter has examined SLA class I expression on a number of porcine cell lines using the analytical tools currently available and has identified two porcine cell lines that express high constitutive levels of MHC class I which can be induced by IFN- α /- γ : Max d/d kidney cell line and primary d/d PAECs. In addition, a cell line that lacks MHC class I expression has been identified, Shimosuma cells. SLA class I expression on PAECs from outbred and d/d animals was compared, demonstrating that SLA epitopes differ in their responses to IFN- α /- γ which underlines the importance of using inbred animals expressing a restricted set of SLA class I genes for SLA class I expression studies. In addition, SLA antibodies localised intra and extracellular SLA class I in PAECs and Max cells expressing high levels of MHC class I by immunofluorescence. This immunofluorescence protocol can be used to monitor gross changes in SLA class I expression on individual cells, such as IFN induction, in future studies. After having characterised MHC class I expression on Max cells, the effects of infection with CSFV on SLA class I expression were examined. CSFV infection caused a dramatic down-regulation in the constitutive levels of SLA class I cell surface expression measured by two antibodies. This represents a novel immune evasion strategy for this virus which has not previously been reported.

3.4.2 General considerations for studying MHC class I

When interpreting experimental data from MHC class I expression studies a number of factors pertaining to the specificity of the SLA antibodies being used and the inherent nature of MHC class I polymorphisms and expression patterns must be considered. These will now be discussed in detail.

3.4.2.1 Antibody specificity

The specificity of the antibodies used in this study were originally defined by serology. To date, antibodies raised against SLA class I have identified and characterised 75 haplotypes (in all breeds tested), indicating that there is considerable variation between individuals in which and how many alleles are expressed (Lunney, 1994). Although serology is a cheap and reliable means of haplotyping pigs and defining the specificity of a particular antibody there are some limitations to consider. Firstly some haplotypes may appear to have the same specificity, but may in fact be subtly different. For example, if gene duplication arose recently the allelic series would look the same by serology but is actually quite distinct. Serology cannot differentiate these 'pseudoalleles' whereas molecular techniques such as RT-PCR, can. Alternatively, the same antibodies can be used to immunoprecipitate MHC class I which can then be differentiated by iso-electric focussing. Secondly, SLA haplotyping is based on the expression of classical SLA genes and does not consider non-classical expression.

The antibody PT-85 has the broadest specificity of all the antibodies used here, having specificity for MHC class I in a range of species with distant evolutionary relationships it can be described as a 'pan-class I' antibody. It must therefore recognise a highly conserved epitope within the MHC class I molecule. The SLA-specific antibody 2.27.3 is serologically defined as poly-loci specific and recognises a common determinant shared among the pig haplotypes H04, H07, H08, H10, H12 and H14 (C. Renard, personal communication). By contrast, the antibodies 2.12.3 and 2.32.1 have an exclusive specificity for the H04 (d/d) haplotype. This has been confirmed by recognition of mouse L cells transfected with the d/d cDNAs encoding PD1 (SLA-1) and PD14 (SLA-2) by these antibodies (Satz & Singer, 1984).

Consideration of regions of structural homology within SLA class I molecules is essential when defining antibody specificity. Antibodies raised against a group of polymorphic molecules with structural homology such as SLA class I may be reactive against both 'public' and 'private' epitopes. The antibodies available to study SLA class I were raised in mice so they could potentially be specific for any part of the SLA class I molecule. To put this point into context, highly conserved regions such as exon 4 are virtually identical in all class I molecules, including the non-classical SLA. Recent studies characterising non-classical SLA expression by RT-PCR have demonstrated that some non-classical genes, namely SLA-8, are ubiquitously expressed, although only a limited number of tissues were studied (Crew *et al.*, 2004). Therefore, when interpreting the FACS data in this study it cannot

be assumed that the SLA antibodies raised against SLA class I have exclusive specificity for classical genes and do not recognise the exon 4 region of this non-classical molecule.

3.4.2.2 Antibody affinity

The affinity of an antibody for an epitope and the concentration of antibody used for staining are also important considerations for FACS analysis. Use of high affinity antibodies at high concentrations is not only wasteful but may lead to non-specific binding. Use of low affinity antibodies at low concentrations will only allow a proportion of the total epitope density to be detected and therefore may not provide a truly accurate representation of the level of epitope expression. For these reasons, antibodies were titrated wherever possible in order to determine an appropriate working dilution for SLA class I detection, that is the concentration of antibody which gives good sensitivity with a high signal:noise ratio. The SLA class I specific antibodies were used undiluted. Higher concentrations may provide better sensitivity, to achieve this affinity purification would be necessary.

3.4.2.3 Tissue specific expression

In the human, it is well established that expression of classical MHC class I molecules is ubiquitous whereas the expression of non-classical genes is tissue specific (reviewed by van den Elsen *et al.*, 2004). Where expressed, the basal level of expression of each class I molecule may also vary between tissues. In the pig this has been demonstrated for the classical gene PD1. PD1 mRNA was detected in all tissues examined, but the degree

of expression varies between tissues (Singer *et al.*, 1987). Expression is highest in mature lymphoid tissues such as the spleen and lymph nodes and peripheral blood lymphocytes whereas the thymus, kidney and testis express relatively low levels of PD1 mRNA. Comparison of two individuals revealed variable levels of PD1 expression in the heart and liver, although lymphocyte contamination could not be ruled out as the cause of this variation (Ehrlich *et al.*, 1989). This is an important consideration when using primary cells, and ideally cells derived from several individuals should be compared in order to minimise the impact of individual variation upon results. Also, use of a panel of antibodies and a comparative analysis of MHC class I expression in a range of cells derived from different tissues would allow tissue specific expression to be explored more comprehensively.

The porcine non-classical genes SLA-6 and SLA-7 are also expressed in a tissue specific manner and the level of class I transcription varies between tissues (Crew *et al.*, 2004). Most importantly for this study, SLA-7 is expressed in the kidney. This has implications for studies in Max cells where cross reactivity of SLA antibodies with the exon 4 region of SLA-7 is a possibility. As discussed above, constitutive expression of SLA-7 and SLA-8 means that FACS profiles may represent both classical and non-classical gene expression. Ideally, future studies should combine serological and molecular techniques such as RT-PCR to study tissue specific expression and rule out cross reactivity of SLA antibodies.

3.4.2.4 Non-genetic factors

The influences of non-genetic factors such as immune status, sub-clinical disease, diet, psychological and physiological stress on SLA class I expression are difficult to define, quantify and explain. However, these factors are particularly relevant in experiments using primary cells such as PAECs, since they have an impact on gene transcription and therefore SLA class I expression. These factors have the potential to contribute to differences in FACS profiles seen between individual pigs. Although primary cells do provide a realistic representation of MHC expression *in vivo*; a dynamic system where expression changes in response to invading pathogens which is influenced by the state of health of the animal, results using these cells can be difficult to explain. The use of cell lines derived from a single individual such as Max cells, can help rule out variation caused by these factors. Ideally, comparative studies using d/d PAECs from a large population of individuals would be used, in practise this is both financially and technically demanding.

Experimental manipulation of cell samples can also affect the density of SLA class I molecules expressed at the cell surface, which in turn influences antibody recognition. Minute differences in technical manipulation during sample preparation for FACS analysis may also affect SLA class I expression and contribute to the variations in levels of class I expressed between experiments observed using cells derived from the same individual.

3.4.2.5 Summary of the limitations of SLA class I antibody use

In summary, the SLA antibodies used in this study can be used to determine the haplotype and allelic series of an individual. However, they may not recognise individual class I molecules or differentiate between pseudoalleles and may recognise both classical and non-classical SLA class I. While these antibodies are important tools for serological analysis, they cannot be used for molecular identification of discrete SLA class I antigens. For these reasons it is important to use a panel of antibodies recognising SLA class I in order to have a clear and unbiased picture of patterns of SLA class I expression. It is also important to examine SLA class I expression on a series of cells originating from different tissues, so that tissue specific expression patterns are considered for both constitutive and cytokine-induced expression. For the reasons discussed above cells originating from both d/d inbred and outbred pigs of unknown haplotypes were chosen to compare SLA class I expression and emphasises the importance of using porcine cells derived from individuals of known haplotype wherever possible.

3.4.3 SLA class I expression in primary cells

SLA class I expression was examined on PAECs from two groups of pigs; large White 'outbred' minipigs of unknown haplotype and inbred NIH minipigs that express the d/d haplotype. Clearly, there will be variation between the alleles expressed constitutively by d/d and outbred individuals and this is reflected in the different FACS profiles seen using the same panel of antibodies for both types of PAEC in figure 3.1. In particular, no staining of

SLA class I on outbred PAECs was observed with 2.12.3 antibody, suggesting that the SLA recognised by this antibody is either not present on outbred populations or not expressed by the individuals examined. The specificity of 2.12.3 antibody was defined by serology; it recognises an SLA determinant that is unique to d/d cells. The results in figure 3.1 therefore correlate with the serological definition of this antibody. Another important consideration when analysing cells from pigs of unknown haplotypes using a small panel of SLA antibodies is that the combined specificities of the antibodies being used may not be broad enough to recognise every single class I allele expressed. This is another argument which supports the use of cells derived from inbred animals of a known haplotype and antibodies with a defined specificity for a particular haplotype, such as d/d locus specific antibodies.

3.4.4 Interferon induction of SLA class I expression

The importance of increasing SLA class I expression in response to a viral infection has already been discussed. Stimulation of outbred PAECs with interferon induced a minimal increase in SLA class I expression compared to IFN- α/γ induction observed in d/d PAECs. Since it is well established that interferon is a potent inducer of MHC class I expression, this observation could be explained by the specificities and concentrations of the antibodies used; it is possible that the increases in class I expression cannot be seen despite having used a panel of SLA class I antibodies at high concentrations. In future studies, the SLA antibodies could be further

concentrated by using affinity columns and tested at a range of higher concentrations than used in this study. The physiological state of the cells would also affect SLA class I expression and impaired response to interferon may be a result of cell culture conditions or the health of the individual. It is conceivable that activated cells express a maximum amount of MHC class I which cannot be further induced by interferon. There are two pieces of evidence which indicate that outbred PAECs have not lost their ability to respond to IFN- α/γ , firstly stimulation of PAECs with IFN- α induces an increase in IFN- α mRNA (Bensaude *et al.*, 2004). Secondly, IFN- α strongly induced SLA class I expression on d/d PAECs, which have been subjected to the same cell culture conditions as outbred PAECs.

The most intriguing observation in the experiments using inbred PAECs stimulated with interferons was the dramatic increase in expression of the SLA recognised by the antibody 2.12.3. As previously stated, 2.12.3 and 2.32.1 have specificity for the H04 haplotype and recognise both PD1 (SLA-1) and PD14 (SLA-2) on mouse L cells stably transfected with these genes (Singer *et al.*, 1987, Ehrlich *et al.*, 1989). This indicated that both antibodies recognise common determinants of these SLA molecules. However, FACS profiles of IFN- α/γ stimulated Max cells and inbred PAECs using these two antibodies were quite distinct, indicating that these two antibodies actually recognise different epitopes. On Max cells the SLA recognised by 2.32.1 is constitutively expressed whereas the epitope recognised by 2.12.3 is not. However, on d/d PAECs, 2.12.3 gave a weak positive signal on untreated cells that was dramatically induced by IFN- α ,

further evidence that these two antibodies recognise distinct SLA gene products. The exact identity of which SLA gene products 2.12.3 and 2.32.1 recognise remains to be determined. What is certain is that whatever 2.12.3 antibody recognises, is very strongly induced by IFN- α/γ . This is interesting in the context of immune response to pathogens and indicates an important allele for response to viral infection. This allele could be targeted in vaccination programmes and therefore it warrants further investigation. Unfortunately, attempts to isolate the SLA class I protein that is recognised by 2.12.3 were limited by the technical applications of this IgM antibody.

3.4.5 CSFV down-regulates MHC class I expression on porcine cells

Importantly, this study has demonstrated that infection with CSFV down-regulates constitutive SLA class I expression on Max cells by FACS. This could be a novel immune evasion strategy for CSFV. The decrease in SLA class I expression was only detected with PT85, 74.11.10 and 2.32.1, but not by 2.27.3, or 2.12.3. As previously discussed, allele-specific expression needs to be controlled by the virus in order to prevent both CTL and NK cell activity therefore targeting CTL specific MHC ligands/SLA class I alleles rather than those which are inhibitory receptors for NK cells would provide CSFV with a distinct advantage. The identity of the class I alleles recognised by the panel of antibodies used in this study remains to be elucidated, therefore specific alleles affected by CSFV infection could not be examined. Since, the exact functions of each porcine SLA class I allele,

particularly non-classical SLA class I are not fully understood at present, it is not possible to determine the implications of SLA allele specific MHC class I down-regulation at this time.

The epitope recognised by the antibody 2.12.3 was not up-regulated by viral infection. This is interesting and suggests that CSFV does not induce an IFN response. Figures 3.1 and 3.3 show that IFN- α strongly induces the expression of SLA class I molecule recognised by this antibody, therefore 2.12.3 can be used as a marker for measuring SLA class I responses to IFNs. During a viral infection, IFN- α transcription and synthesis is induced when members of the Toll Like receptor family of receptors recognise intermediates of viral replication (figure 1.7). Taking this into account, one would anticipate that CSFV would induce IFN- α secretion and stimulate an increase in SLA class I expression, which would be reflected by an increase in the allele recognised by 2.12.3. This hypothesis is supported by previous data which demonstrated that IFN- α secretion is inhibited in CSFV infected cells (Bensaude *et al.*, 2004 & Larocca *et al.*, 2005).

The next challenge is to elucidate the mechanism by which CSFV down-regulates MHC class I. Key questions to address are does this virus down-regulate SLA class I expression by disrupting MHC class I transcription, synthesis, assembly or transport and which viral proteins are involved. As discussed in the introduction, viruses can disrupt the MHC class I presentation pathway at many stages. In chapter 5 the effect of CSFV on SLA class I expression at the level of transcription will be examined.

3.4.6 Summary

In order to understand the mechanisms that regulate SLA class I expression in more detail, I chose to investigate SLA class I expression at the transcriptional level by studying SLA class I gene promoters and study the effects of cytokines, transcription factors and CSFV upon SLA promoter activity. The next chapter will describe how promoters regions of classical and non-classical SLA genes derived from the H01 haplotype pig were cloned into a luciferase reporter vector. Investigation of the transcriptional control of six SLA class I genes is described in Chapter 5.

Chapter 4: Physical and Structural Characterisation of the SLA Class I Locus

4.1 Introduction

A series of overlapping Bacterial Artificial Chromosomes (BACs) that span the entire MHC class I locus of an HO1 haplotype pig were produced and sequenced by the LREG laboratory at INRA, France in collaboration with Genoscope, France (Velten *et al.*, 1998; Rogel-Gaillard *et al.*, 1999). Sequence analyses identified three classical class Ia genes (SLA 1, 2 and 3), three non-classical class Ib genes (SLA-6, -7 and -8) and one class Ic gene (MIC-2) with the correct intron/exon organisation and promoter elements that suggest that they encode functional SLA class I molecules (Chardon *et al.*, 2001; Renard *et al.*, 2001). The first aim of this work was to identify which genes could be transcribed from these DNA sequences by functionally analysing the upstream promoter elements. The sequences upstream of the MHC class I gene coding regions of human, mouse and the pig PD1, PD7 and PD14 class I genes contain interferon Sensitive Response Elements (ISRE), Enhancer A, NF- κ B and SXY sites (Ehrlich *et al.*, 1998). Analysis of the individual SLA class I gene promoter regions of the H01 pig identified the same conserved regulatory elements including; an ISRE containing putative IRF binding sites, an Enhancer A containing putative NF- κ B binding sites and an SXY region (Chardon *et al.*, 2001, Renard *et al.*, 2001). Chapter 5 will determine which of these elements are functional in the pig and explore the interactions of these motifs with the relevant transcription factors.

This chapter includes a summary of the physical information obtained from DNA sequence analysis, including alignment of pig and human HLA promoters, and an alignment of the promoter elements for seven of the classical and non-classical SLA class I genes. It also describes the cloning strategy used to express SLA class I promoters in luciferase reporter plasmids to analyse their transcriptional activity.

4.2 Description of the porcine MHC class I locus and regulatory promoter elements.

Figure 4.1 A shows a diagram of the major histocompatibility complex (SLA) of the pig located on chromosome 7. It has been divided into three regions according to gene content; class II, class III and class I regions. The position of the centromere is indicated by a red circle. The regions containing postulated open reading frames (ORFS) for SLA class I are shown in blue, SLA class II genes are shown in yellow and other functional genes of interest are shown in white. A genomic library of BACs spanning the entire class I region of the Large White haplotype H01 was produced and a BAC contig spanning the entire class I region was built. The BAC clones corresponding to the minimum tiling path were sequenced (Chardon et al., 2001, Renard et al., 2001). Three BACs were identified which, among other genes, collectively contain all of the SLA class I genes.

Figure 4.1: Physical structure of SLA promoter regions on chromosome 7 and identification of SLA Class I genes from Bacterial Artificial Chromosomes (BACs).

A: The MHC class I locus of the pig is located on chromosome 7 of the pig and spans approximately 2 Mega bases (Velten *et al.*, 1998). The class I region contains SLA class Ia, Ib and Ic genes, shown in blue. The class II region contains SLA class II genes, shown in yellow. Other functional genes of interest are depicted as white boxes.

B. The gene content of three BACs which include all of the SLA class I genes is shown. Functional genes are shown in blue, pseudogenes are shown as hatched boxes and the direction of transcription is designated by the arrows. BAC 493A7 contains the non-classical class Ib genes SLA-6, -7 -8 and the class Ic gene MIC-2. BAC 490B10 contains the classical class Ia genes SLA-2 and -3, the putative class Ia gene SLA-5 and three pseudogenes SLA-11, -4 and -9. BAC 207G8 contains SLA-1.

C: BAC sequence data were used to design primers for the amplification of full length promoter regions of SLA-1, -2, -3, -6, -7, -8 and MIC-2 genes (red circles). The promoter regions amplified and the nucleotide positions of primers is shown with respect to their positions on each BAC (red rectangles).

D: Alignment of SLA promoter regions reveals a series of conserved regulatory domains in class Ia and Ib genes including a distal domain containing enhancer and silencer sites, a proximal domain containing a region homologous to a functional motif within the mouse H-2k gene, a proximal enhancer, two putative Sp1 binding sites, Enhancer A and an interferon stimulated response element (ISRE) (Chardon *et al.*, 2001; Renard *et al.*, 2001). Three transactivator elements common to MHC class I, class II and b2-microglobulin promoters, w, x1 and y are located within the proximal region. A series of proximal or "core" elements including CAT and TATA boxes and a CAP binding motif are also present.

Figure 4.1 B shows the content and physical organisation of the genes identified within each BAC (accession numbers for sequence data are indicated). Individual genes are shown as blue arrows and hatched arrows indicate pseudogenes. The orientation of each gene and proposed direction of transcription is indicated by the direction of the arrow. BAC 207G8 contains the class Ia gene SLA-1. BAC 490B10 contains the class Ia genes SLA-2 and SLA-3, the putative class Ia gene SLA-5 and the pseudogenes SLA-4, -9 and -11. BAC 493A6 contains the class Ib genes SLA-6, -7 and -8 and the class Ic gene MIC-2.

Figure 4.1 C shows the nucleotide positions, with respect to each BAC, that were selected for cloning out full length SLA class I promoters from the distal enhancer silencer sites to the translation initiation sites identified previously (Chardon *et al.*, 2001, Renard *et al.*, 2001). Figure 4.1 D shows an alignment of the promoter regions of SLA class Ia and Ib and indicates the key regulatory domains thought to be implicated in the regulation of SLA class I expression. SINE elements within the promoter regions of SLA-2 and SLA-3 have been excluded from this alignment. When compared to human and mouse MHC class I the DNA sequences of the seven SLA class I genes revealed conserved regulatory domains, highlighted in red. A series of regulatory domains were identified within the 5' flanking regions of SLA class I genes including several distal enhancer silencer sites, a region homologous to that found in mouse H-2k and an enhancer. In addition, a proximal domain was identified containing functional motifs implicated in cytokine-induced regulation of HLA expression including two putative Sp1 binding sites, an

Enhancer A region containing putative binding sites for NF- κ B family transcription factors and an ISRE region which may bind IRF-1 and IRF-2. A class I transactivator region containing W/S, X and Y elements was deduced from comparison of SLA promoters with other vertebrate class I and class II promoter sequences that have already been characterised (van den Elsen *et al.*, 1998). In humans, the S, X1, X2 and Y domains bind RFX, CREB and NF-Y respectively and control constitutive HLA class I expression. CIITA acts as a co-activator of transcription, interacting with RFX, CREB and NF-Y bound to SXY motifs and drives HLA class I expression in response to IFN- γ . A CAT box, TATA box and CAP binding domain were also identified. A detailed comparison of individual SLA class I promoter sequences reveals locus-specific differences in promoter elements that may be responsible for differential levels of constitutive and cytokine-induced expression of the individual SLA class I genes. The functional motifs within SLA-1, -2 and -3 promoters are highly conserved whereas those of non-classical promoters contain some nucleotide differences which may affect their ability to bind transcription factors. For example, since the SLA-8 promoter lacks an ISRE, type I IFN may not stimulate the transcription of this gene.

In addition to the regulatory motifs described above, putative STAT binding motifs were identified in the promoter regions of SLA-1, -2, -3 and -6. These previously undescribed motifs and their position with respect to the transcriptional initiation site (ATG) are listed in table 6.

Promoter	Motif sequence	Position
SLA-1	TTACTGGAA	-952
SLA-2	TTACTTCAA	-501
SLA-3	TTACTGGAA	-1268
SLA-5	TTCCTGGAA	-1233
SLA-6	TTCAGGGAA	-327
SLA-7		
SLA-8		
MIC-2		

Table 6: Putative STAT binding motifs in SLA class I promoters

Analysis of promoter sequences using TFSEARCH software reveals putative STAT binding sites within the promoters of SLA-1, -2, -3, -5 and -6. The table shows the sequence of each STAT motif and the location within each promoter region with respect to the ATG. These motifs have not been previously described or tested experimentally.

The functions of promoter regulatory domains and the specific sequences that bind transcription factors within these domains have been studied in detail for human MHC class I genes (summarised in figure 1.8). The information from these human studies can be used as a comparative model for studying the molecular regulation of SLA class I expression. In figure 4.2, the promoter region of SLA-1 is aligned with the promoter regions of HLA-A, -B and -C and a consensus sites for each regulatory element is shown. Since SLA-1, -2 and -3 share highly homologous sequences in Enhancer A, ISRE and SXY domains only SLA-1 is shown here.

4.3 Cloning of SLA class I promoter regions

Sequence data were used to design primers for the amplification of full length SLA class I promoter regions from the 5' distal enhancer silencer sites to the 3' ATG from BAC DNA. Promoter regions of approximately 1000 nucleotides in length were cloned into the luciferase reporter vector pGL3-basic (Promega), listed in table 7.

The DNA containing each SLA promoter was subcloned into pcDNA3.1/V5-His/TOPO (Invitrogen), excised, then ligated into the promoter-less reporter vector pGL3-basic upstream of the luciferase gene. Plasmids were then analysed by digestion with *NheI* and *HindIII* followed by gel electrophoresis to re-confirm the size of promoter inserts (figure 4.3 and table 7). The sequences of SLA promoter inserts within pGL3-basic were confirmed by DNA sequencing.

Figure 4.2: Comparison of porcine SLA-1 promoter sequence with human HLA-A1, -B7 and -Cw7 alleles.

This figure shows an alignment of the promoter region of SLA-1 with three common human classical HLA class I alleles. The 5' flanking sequence of SLA class Ia share homology with human class I a genes within regions containing functional motifs including enhancer A, ISRE, w/s, x and y domains. Purine and pyrimidine residues are shown in coloured boxes: arginine (yellow), thymidine (red), guanine (green) and cystine (blue). The functions of these promoter domains have been characterised for human class I genes but not for H01 SLA class I genes. Consensus sites for each motif are indicated below each section of the alignment .

Target gene	Anticipated PCR product size
SLA-1	1069
SLA-2	1330
SLA-3	1402
SLA-6	988
SLA-7	997
SLA-8	962
MIC-2	1228

Table 7: Anticipated sizes of SLA class I gene promoter regions to be cloned into pGL3-basic

Figure 4.3: Analysis of DNA amplified from BACs and cloned into pGL3 basic.

Promoter DNA was cloned into pGL3 basic and the resulting plasmids were digested with *Hind*III and *Nhe*I and run on an analytical agarose gel to confirm molecular weight of inserts. The predicted size of the promoter regions of SLA-1, -2, -3, -6, -7 and MIC-2 genes amplified from BACS by PCR in table 7 corresponds with the sizes of promoter regions cloned into pGL3-basic. These results were also confirmed by sequencing.

All SLA promoter constructs were sequenced with both pGL3-basic vector-specific and SLA sequence specific primers to confirm the presence of SLA promoter sequences upstream of the luciferase gene. Two clones of each construct were sequenced four times and compared to the reference BAC sequences (see Figure 4.1C for accession numbers). The promoters of SLA-1, -2, -3, -6, -7, and MIC-2 genes were successfully ligated into pGL3-basic. Cloning of SLA-8 was unsuccessful and therefore this promoter is excluded from this study. Sequence data for pGL3-SLA-1, -6, -7 and MIC-2 matched the BAC reference sequences with 100 % identity. Analysis of sequence data for SLA-2 and SLA-3 revealed several differences to the published sequence, these are reported in figure 4.4. Nucleotides that are at variance with the published sequence are shown in bold type. Nucleotide positions are given with reference to the cloned region of SLA-1 and -2 promoters (refer to figure 1 C for address on BAC). None of the base changes reported here are within the Enhancer A (blue), ISRE (red), S (yellow), X (green) or Y (orange) regions of the SLA-2 and SLA-3 promoters therefore are not regarded to be of significance for the experiments described in Chapter 5.

4.4.1 Summary of results

This chapter has described the production of reporter plasmids containing the whole of the regulatory region for six SLA class I genes, including classical genes SLA-1, -2 and -3, non-classical 1b genes SLA-6, and -7 and the class Ic gene MIC-2. Evidence from restriction digests and

SLA-2

SLA-3

Figure 4.4: Sequence analysis of SLA promoters

Sequencing of SLA-2 and SLA-3 constructs revealed several modifications to the published sequences within BAC 490B10. Two base changes, one insertion and one deletion in the sequence of the SLA-2 promoter and one base change in the SLA-3 promoter are shown in bold. Importantly, none of the base changes are located within the enhancer A, ISRE or SXY promoter modules.

DNA sequencing confirmed that SLA promoter regions were cloned into each reporter construct.

4.4.2 The sequences of the SLA-2 and SLA-3 promoter differ to the published sequences.

Comparison of our sequence data with the published sequence data identified several differences in the sequences of SLA-2 and SLA-3. The original BAC sequence data provided by Chardon *et al* (2001) and Renard *et al* (2001) were generated from sequencing of each BAC clone at least 10 times. BAC DNA is considered as very stable and it is therefore unlikely that the base changes reported here arose during manipulation and preparation of BAC DNA or errors introduced by the *pfu* polymerase used for PCR amplification. Crucially, none of the base changes reported here are located within the regulatory domains of the promoter that will be studied here and it is anticipated that these changes will not affect promoter activity or be of consideration for the functional studies performed in chapter 5.

4.4.3 Functional analysis of SLA class I promoters from the H01 pig

The BAC sequences used to generate SLA promoter reporter constructs were derived from a male, Large White pig expressing the H01 haplotype. At the time this study commenced, little was known about which SLA class I genes are expressed by this haplotype. In the recent paper by Crew *et al* (2004), the expression of the non-classical genes has been

clarified. What remains unclear is how expression of SLA class I genes is regulated at the molecular level. Variations in the promoter regions of SLA genes affect levels of constitutive expression; this has been demonstrated in the d/d pig where variation in PD7 and PD14 expression may be explained by base changes in the enhancer region (Ehrlich *et al.*, 1998). In addition, nucleotide variations in promoter regions of SLA class I genes of different haplotypes may effect regulation through transcription factor binding. The response of the SLA genes of the H01 pig to cytokines and transcription factors is unknown. Since the promoter regions of HLA class I have been well characterised, these human genes can be used as a model for the pig class I genes. In the next chapter I will determine the constitutive activities of SLA class I promoters and investigate the functional activity of each SLA class I promoter in respect to responses to cytokines, transcription factors and viral infection.

Chapter 5: Transcriptional regulation of classical and non-classical SLA class I genes.

5.1 Introduction

This chapter will investigate the mechanisms of regulation of SLA class I gene expression using the SLA class I promoters cloned into reporter plasmid described in the previous chapter. Key questions to address are: which promoters are constitutively active? and which SLA promoters are induced by pro-inflammatory cytokines? Induction by cytokines will be examined at the molecular level by examining promoter responses to transcription factors that are induced by cytokines and predicted to interact with the ISRE, Enhancer A and SXY core regulatory motifs within SLA class I promoters. In addition, the effects of CSFV infection on SLA gene expression will be investigated. In chapter 3 it was demonstrated that CSFV down-regulates the expression of SLA class I molecules at the cell surface. In this chapter, SLA promoter constructs will be used to determine if CSFV targets SLA class I at the level of gene transcription and which SLA class I genes are involved.

5.2 Constitutive activity of classical and non-classical SLA class I promoters in porcine cells

To determine the constitutive activity of SLA-1, -2, -3, 6, -7 and MIC-2 promoters in porcine cells in culture, classical and non-classical promoters were cloned in front of a luciferase reporter gene (as described in chapter 4).

They were transiently transfected into the porcine Max cell line isolated from an inbred NIH minipig expressing the d/d haplotype (described in chapter 3). As a control for transfection efficiency, Max cells were co-transfected with the plasmid pJATLacZ, a β -galactosidase reporter gene under the control of the rat β -actin promoter (Didcock *et al.*, 1999). Normalised luciferase activities were calculated for each sample as shown below, this calculation was used in all subsequent transfection experiments unless otherwise stated.

$$\text{Normalised LUC} = \frac{\text{luciferase reading}}{(\beta\text{-gal reading} - \text{background})}$$

The constitutive activities of SLA class I promoters 48 hours post transfection of Max cells is shown in Figure 5.1. The data shown here represent the mean average RLU from three repeats and is a qualitative representation of the trends observed throughout this study. The trend observed was that SLA-1 was most active of the three classical SLA promoters, SLA-2 had less activity and SLA-3 activity was the lowest of the three classical genes. Of the non-classical promoters, SLA-7 was most active, and in this cell type was always higher than SLA-1. In this example SLA-7 activity is approximately three times that observed for SLA-1. Constitutive SLA-6 activity was also high, to a similar magnitude as SLA-1. MIC-2 has a constitutive activity comparable to SLA-2.

Figure 5.1: Constitutive activity of SLA class I promoters in porcine cells

Max cells were transfected with the SLA class I promoter-driven luciferase reporter plasmids pGL3-SLA-1, -2, -3, -6, -7 -8 or -MIC-2 and co-transfected with the β -galactosidase control pJATLacZ for normalisation. After 48 hrs, luciferase and β -galactosidase assays were performed. Luciferase readings were normalised with β -galactosidase readings to obtain normalised relative light units (RLU). This graph shows the mean average normalised RLU in untreated cells for each SLA promoter construct from a single experiment containing three repeats. The trend observed is representative of that observed throughout this study. Of the classical SLA promoters, SLA-1 was the most active, followed by SLA-2 and SLA-3. The non-classical SLA-7 promoter was the most active of all promoters tested. SLA-6 was also strongly active. The MIC-2 promoter is constitutively active, with an activity comparable to SLA-2.

5.3.1 Regulation by interferons

In order to determine whether the ISRE and putative STAT binding motifs identified within the promoter regions of SLA class I genes are functional, the responses of SLA class I promoter reporters to stimulation with porcine interferon alpha and gamma were examined. First it was necessary to set up a series of controls. The reporter vector pGL3-control, which contains the firefly luciferase gene under control of the SV40 promoter, was used as a positive control for the luciferase assay. The reporter plasmid pISRE contains 4 copies of the ISRE consensus sequence (AGGAAATAGAACTG) arranged in tandem upstream of the firefly luciferase reporter gene of a ptk Δ (-39)lucifer plasmid (King & Goodbourn, 1998, Reid et al., 1989). This plasmid was used as a positive control to monitor cellular responses to IFN- α . The reporter plasmid pGAS contains 2 copies of the IRF-1 GAS consensus sequence (TTTCCCCGAAA) arranged in tandem upstream of the firefly luciferase reporter gene of the ptk Δ (-39)lucifer plasmid (King & Goodbourn, 1994, King & Goodbourn, 1998). This plasmid was used as a positive control for IFN- γ activation. In control experiments, Max cells were transfected with pISRE or pGAS and co-transfected with pJATLac for 32 hours, followed by treatment with pIFN- α or pIFN- γ for 16 hours. Cell lysates were analysed for luciferase and β -gal activity, and luciferase was normalised against β -gal as described above and normalised β -galactosidase activity was calculated. Statistical significance was evaluated for each condition using Wilcoxon signed ranked tests on paired raw data.

Stimulation of Max cells transfected with pISRE with IFN α and cells transfected with pGAS with IFN- γ caused a 2.1- and 3.2-fold increase in luciferase synthesis respectively (Figure 5.2 A i and ii). Interestingly, there was a significant decrease in pGL3-control plasmid activity when treated with IFN- α (Figure 5.2A iii) ($p=0.028$). The addition of IFN- γ also caused a significant decrease in luciferase production (0.6) by pGL3-control ($p=0.028$).

The classical SLA class I promoter constructs pGL3-SLA-1, -2 and -3 were transiently transfected into Max cells together with the control pJATLacZ and stimulated with IFN- α or IFN- γ for 16 hours. The normalised fold inductions of luciferase activity as a consequence of IFN- α and IFN- γ stimulation are shown in Figure 5.2 (i) and (ii) respectively. The classical SLA promoters SLA-1, -2 and -3 are more sensitive to stimulation with IFN- γ than IFN- α . IFN- α induced the activity of the SLA-1 promoter by mean fold induction of 1.6, a value which is close to statistical significance ($p=0.068$). Induction of the SLA-2 promoter by IFN- α is negligible, a mean fold induction of 1.1 is reported here ($p=0.068$).

IFN- γ increased SLA-1 promoter activity by 4.1 fold ($p=0.015$). SLA-2 and -3 also appear to be induced by IFN- γ , a mean fold induction of 2.5 and 2.0 was observed and these data are approaching significance ($p=0.068$ and

The responses of the non-classical SLA promoters of SLA-6, -7 and MIC-2 genes to IFN- α and IFN- γ were investigated by transient transfection

Figure 5.2: Effect of interferon stimulation on classical SLA class I promoters.

Max cells were transfected with pJATLacZ and SLA promoter luciferase reporter plasmids (SLA-1, -2 and -3) or the control plasmids pISRE, pGAS or pGL3-control for 32 hours. Transfected cells were incubated with IFN- α or IFN- γ for a further 16 hrs. These graphs show the mean normalised fold induction $\pm$ standard deviation of the mean of normalised luciferase following IFN stimulation compared with untreated controls for each condition and plasmid tested. Wilcoxon signed ranked tests were performed on paired raw data. Statistically significant p values are marked with asterisks ($p < 0.05$). The number of repeats (n) is indicated below each graph. Panel A shows the results for the control plasmids. In (i), the positive control for IFN- α activity, pISRE is induced by IFN- α , in (ii), the positive control for IFN- γ activity, pGAS is induced by IFN- γ and in (iii) the positive control pGL3-control is repressed by IFN- α and IFN- γ . In panel B (i), IFN- α appears to induce SLA-1 promoter activity. Induction of SLA-2 is negligible. In B (ii), IFN- γ significantly induces SLA-1 activity and also appears to induce SLA-2 and SLA-3 activity.

of Max cells with the promoter constructs pGL3-SLA-6, pGL3-SLA-7 or pGL3-MIC-2 for 32 hours. Cells were then treated with pIFN- α or pIFN- γ for 16 hours as described above. Promoter responses to IFN- α and IFN- γ stimulation are shown in figure 5.3 [(i) and (ii) respectively]. The SLA-6 promoter is neither induced nor repressed by IFN- α or IFN- γ . For SLA-7, a significant repression was observed with IFN- α ($p=0.021$) but IFN- γ does not affect the transcription of this gene. Finally, MIC-2 promoter activity appears not to be repressed by IFN- α or - γ although statistical significance could not be measured these data due to the small sample size.

Pro-inflammatory cytokine such as TNF- α and interleukins have been shown to up-regulate expression of MHC class I genes in the human (Israel *et al.*, 1989, Johnson, 2003). In order to determine whether TNF- α induces SLA class I expression, the classical SLA class I promoter construct pGL3-SLA-1 and the non-classical promoter construct pGL3-SLA-7 were transiently transfected into Max cells together with the normalisation control pJATLacZ for 32 hours then stimulated with human TNF- α for 16 hours. The reporter vector pGL3-control was used as a positive control for luciferase activity as described before. Statistical significance was calculated using Wilcoxon's signed ranked tests on paired raw data. In figure 5.4a (yellow bars), SLA-1 activity increased by a mean fold induction of 1.9. A small, but significant fold induction (1.3) of the SLA-7 promoter was also observed ($p= 0.008$) (figure 5.4c yellow bar). The luciferase activity of the control plasmid pGL3-control was not affected by TNF- α (figure 5.4c, yellow bar).

Figure 5.3: Effect of interferon stimulation on non-classical SLA class I promoters.

Max cells were transfected with pJATLacZ and SLA promoter luciferase reporter plasmids or the control plasmids pISRE, pGAS or pGL3-control for 32 hours then incubated with pIFN- α , pIFN- γ or media alone for a further 16 hrs. The plasmids pISRE, pGAS and pGL3 control were used as controls (refer to figure 2). Graphs show the average fold induction $\pm$ standard deviation of normalised luciferase following pIFN- α (red) and pIFN- γ (green) stimulation compared with unstimulated controls for each condition and plasmid tested. The number of repeats (n) is indicated below each graph. Wilcoxon signed ranked tests were performed on paired raw data and asterisks indicate statistically significant data ($p < 0.05$). The SLA-6 promoter was not affected by IFN treatment. SLA-7 is repressed by IFN- α but is not affected by IFN- γ . The MIC-2 promoter was not affected by IFNs (statistical test not performed).

Figure 5.4: Stimulation of SLA-1 and SLA-7 promoters with TNF- α and co-stimulation with IFN- α /- γ and TNF- α

Max cells were transfected with pJATLacZ and pGL3-SLA-1 or pGL3-SLA-7 promoter reporter plasmids or the control plasmid pGL3-control for 32 hours. Transfected cells were then incubated with (i) TNF- α (yellow bars), (ii) TNF- α plus IFN- α (orange bars) or (iii) TNF- α plus IFN- γ (blue bars) for a further 16 hrs. Graphs show the average fold induction $\pm$ standard deviation of normalised luciferase following cytokine stimulation compared with untreated controls for each condition and plasmid tested. The number of repeats (n) and standard deviation (SD) from the mean fold induction is indicated below each graph. Wilcoxon signed ranked paired tests were performed on raw RLU data, asterisks indicate statistical significance ($p < 0.05$).

Previous studies have shown that co-stimulation with TNF- α and IFN- α or IFN- γ synergistically increase HLA class I promoters (Drew *et al.*, 1995b, Johnson, 2003). Therefore, the synergistic induction of SLA class I expression by TNF- α and IFNs was investigated using the classical promoter construct pGL3-SLA-1 and the non-classical promoter construct pGL3-SLA-7. Max cells were transiently transfected with pGL3-SLA-1 or pGL3-SLA-7 and pJATLacZ for 32 hrs then treated with a combination of TNF- α and IFN- α or IFN- γ for 16 hours, as in previous experiments, pGL3-basic was used as a negative control for cellular responses to IFN- α/γ and TNF- α . Mean fold inductions for each reporter plasmid are shown in figure 5.4. As above, statistical significance was determined using Wilcoxon's signed ranked tests on paired raw data.

In control experiments, TNF- α did not affect luciferase production in cells transfected with the control pGL3-control (figure 5.4a yellow bar). Combined treatment with TNF- α and IFN- α represses luciferase activity of pGL3 compared with the untreated control and compared with TNF- α alone ($p=0.028$), shown in figure 5.4a (orange bar) as a fold induction of 0.3. IFN- γ also represses pGL3 control, although to lesser extent than IFN- α , shown in figure 5.4a as a fold induction of 0.6 ($p=0.028$).

The data presented in figure 5.4b show a trend indicative of SLA-1 induction by TNF- α . A mean fold induction of 1.9 is reported although in this experiment, these data approach, but do not reach significance ($p=0.068$).

Co-stimulation of SLA-1 with TNF- α and IFN- α does not cause a synergistic induction of the SLA-1 promoter compared with TNF- α alone (figure 5.4b, orange bar). Co-stimulation of SLA-1 promoter with TNF- α and IFN- γ resulted in a mean fold induction of 4.4 which was higher than that reported for IFN- γ alone (4.1) or TNF- α alone (2.0). This observation could not be confirmed statistically ($p=0.066$) and therefore may reflect the variation in induction by IFN- γ , which is mirrored in figure 5.2b (iii), rather than synergistic induction by these two cytokines.

5.4 Regulation of SLA promoters by transcription factors

In order to determine which promoter elements are involved in transcriptional regulation of SLA class I in response to cytokines at the molecular level, SLA class I promoter constructs were co-transfected into Max cells with a number of transcription factors that are involved in the interferon and NF- κ B signal transduction pathways. Many of these transcription factors have been shown to bind specific promoter motifs of HLA class I promoters and induce transcription of these genes. The regulation of three core promoter motifs within SLA class I promoters: ISRE, Enhancer A and SXY will now be considered.

5.4.1 The interferon-stimulated response element (ISRE)

The previous experiments showed that classical SLA class I promoters were regulated by interferon. Interferon alpha and gamma induce the expression of members of the interferon regulatory factor family (IRFs)

through the JAK/STAT pathway of signal transduction. IRFs then bind to the consensus sequence G/A G/C TTTC within ISRE elements of interferon sensitive genes, including human MHC class I promoters and modulates expression of these genes.

ISRE were found within all SLA class I promoters with the exception of MIC-2 and SLA-8 (Chardon *et al.*, 2001, Renard *et al.*, 2001). Figure 5.5 shows an analysis of the nucleotide sequences of the ISRE elements of SLA-1, -2, -3, -6, and -7 and a comparison with HLA-A, -B, -C, -E, -F and -G. The ISREs of SLA-1, -2 and -3 are identical to the ISRE of HLA-A1. It is therefore likely that the ISRE motifs of these SLA promoters will bind and respond to IRFs. By contrast, the sequences of ISRE in non-classical SLA-6 and SLA-7 promoters not only differ from classical HLA and SLA class I, but also from the three human non-classical genes, HLA- E, -F, and -G. The ISRE of SLA-6 contains an insertion of 2 nucleotides that may disrupt transcription factor binding or affect the spatial arrangement of transcription factor complex binding. Overall, the ISRE of SLA-7 is very similar to that of classical SLA and HLA. However, there are three nucleotide variations up- and downstream of the consensus IRF binding motif which may have an impact on transcription factor binding.

In humans, IRF-1 is a transcriptional activator of MHC class I whereas IRF-2 functions primarily as a transcriptional repressor (Reis *et al.*, 1992, Drew *et al.*, 1995a). To determine the effect of IRF-1 and -2 on SLA class I promoters experimentally, Max cells transiently transfected with the promoter

Figure 5.5: Analysis of ISRE sequences from MHC class I promoters

Nucleotide sequences of ISRE regions from classical and non-classical SLA and HLA class I were aligned to highlight similarities and differences between human and porcine classical and non-classical genes.

Figure 5.6A and B: Effect of IRF-1 and IRF-2 on SLA class I promoters

Max cells transfected for 48 hrs with SLA promoter constructs or a positive control reporter for IRF-1 activity and co-transfected with IRF-1 (A) or IRF-2 (B) expression vectors. Graphs A and B show fold induction of normalised luciferase values $\pm$ standard deviation (SD) when compared to a negative control where pIRF-1/IRF-2 was substituted with pcDNA3.1. Statistical significance was calculated using Wilcoxon signed ranked tests on paired raw normalised luciferase data sets. Asterisks show significant data ($p < 0.05$).

reporter constructs pGL3-SLA-1, -3, -7, or -MIC-2 were co-transfected for 48 hours with an IRF-1 expression plasmid (pEFPlink2-IRF-1) or an IRF-2 expression plasmid (pEFPlink2-IRF-2) or with the negative control backbone plasmid pEFPlink2. In all cases cells were also transfected with the β -galactosidase reporter plasmid pJATLacZ for normalisation. A control reporter p[(AAGTGA)₄5tk Δ (-39)]lucifer was used to monitor IRF-1 activity. This plasmid contains a luciferase gene under the control of a synthetic promoter containing the herpes simplex virus thymidine kinase TATA box and 5 tandem repeats of an IRF-1/2 binding motif. Normalised luciferase values were calculated for each sample as above and statistical significance was assessed using Wilcoxon signed ranked tests on paired data. For simplicity, fold inductions were calculated by comparing mock co-transfected and IRF-1 co-transfected normalised luciferase data and the mean fold induction for each SLA promoter is shown in figure 5.6A.

Co-transfection of IRF-1 with the positive control vector HEX-4 results in a significant increase in transcription of the luciferase gene ($p=0.043$). In figure 5.6a, transcriptional activity of the SLA-1 promoter is induced by IRF-1 ($p=0.028$), a mean fold induction of 1.6 is reported here. Induction of SLA-3 by IRF-1 does not reach significance despite a mean fold induction of 1.5, perhaps due to the considerable variation in induction levels causing a substantial standard deviation. Interestingly, a 1.4 fold induction of the non-classical SLA-7 promoter by IRF-1 was observed ($p = 0.028$). Preliminary data for the MIC-2 promoter suggests that transcription of this gene is not

affected by IRF-1 although statistical analysis could not be performed on this small number of data points.

Co-transfection with the IRF-2 cDNA caused a variety of effects on classical SLA class I promoters (Figure 5.6B). Interestingly, and rather unexpectedly, a mean fold induction of 1.6 suggests that SLA-1 is weakly induced by IRF-2, these data are approaching significance ($p=0.068$). IRF-2 co-transfection caused a fold induction of 1.1 in SLA-2 promoter activity, suggesting that IRF-2 does not repress SLA-2 promoter activity. The SLA-3 promoter however was repressed by IRF-2. Transfection with IRF-2 did not affect SLA-7 or MIC-2 promoters.

Next, the effect of expression of IRF-3 on SLA class I promoters was investigated. Activated IRF-3 binds in a homodimeric complex to the same consensus sites as IRF-1 and -2. Max cells were transiently transfected with pGL3-SLA-1, -2 and -7 and co-transfected with an IRF-3 expression vector and the β -galactosidase control pJATLacZ for normalisation for 48 hrs. Normalised fold induction of SLA promoters in samples where IRF-3 was expressed were calculated by comparison with samples transfected with pcDNA3.1 control vector. In figure 5.7, IRF-3 induces SLA-1 promoter activity by nearly 3-fold ($p=0.028$). Interestingly, IRF-3 had no significant effect on SLA-2 or SLA-7 promoters despite both these promoters containing an ISRE and a reasonable sample size. This indicates that IRF-3 is a more important transcription factor for gene induction from SLA-1.

Figure 5.7: Effect of IRF-3 expression on SLA class I promoters

Fold induction of normalised luciferase readings from Max cells transfected for 48 hrs with SLA promoter constructs, IRF-3 and pJATLacZ. The standard deviation (SD) and number of repeats (*n*) is indicated for each experiment. Statistical significance of inductions was calculated using Wilcoxon signed ranked tests on paired raw normalised luciferase data sets, significant data ($p < 0.05$) are marked with an asterisk.

Figure 5.8: Effect of IRF-7 expression on SLA class I promoters

Fold induction of normalised luciferase readings from Max cells transfected for 48 hrs with SLA promoter constructs, IRF-3 and pJATLacZ. The standard deviation (SD) and number of repeats (*n*) is indicated for each experiment. Statistical significance of inductions was calculated using Wilcoxon signed ranked tests on paired raw normalised luciferase data sets, significant data ($p < 0.05$) are marked with an asterisk.

The effect of IRF-7 on SLA-1, -2, -7 and MIC-2 promoters was investigated next. Max cells were transfected with SLA promoter constructs, the control pJATLacZ and an IRF-7 expression vector pIRF-7-GFP. Normalised fold inductions were calculated as before, comparing luciferase output from cells transfected with and without pIRF-7-GFP. Figure 5.8 shows a trend that IRF-7 induces SLA-1 by a mean fold induction of 1.9 and SLA-2 by a mean fold induction of 2.3, but these observations could not be confirmed by statistical analysis. The non-classical SLA-7 promoter and the MIC-2 promoter were not induced by IRF-7.

The effect of IRF-9 expression on SLA class I promoters was determined in a similar fashion. IRF-9 stimulates transcription of ISGs only as part of a complex with STAT1 and STAT2. Therefore, a constitutively active IRF-9 fusion protein that combines IRF-9 with the transactivator domain of the HSV1 VP16 protein was co-transfected into Max cells with the promoter constructs SLA-1, SLA-7 or MIC-2 and pJatLacZ for normalisation. As in previous experiments, normalised fold inductions were calculated for each condition and statistical significance was calculated using Wilcoxon signed ranked tests on pairs of normalised luciferase data. In the experiment shown in figure 5.9, co-expression of IRF-9 induces SLA-1 promoter activity ($p=0.043$), a mean fold induction of 2.9 was observed. SLA-7 appears to be weakly induced by IRF-9, significance was not reached using these data. Preliminary indications suggest that IRF-9 has no effect on MIC-2, a larger pool of data is required to confirm this.

Figure 5.9: Effect of IRF-9 expression on SLA class I promoters

Fold induction of normalised luciferase readings from Max cells transfected for 48 hrs with SLA promoter constructs, pIRF-9vp16 and pJATLacZ. The standard deviation (SD) and number of repeats (n) is indicated for each experiment. Statistical significance of inductions was calculated using Wilcoxon signed ranked tests on paired raw normalised luciferase data sets, significant data ($p < 0.05$) are marked with an asterisk.

5.4.2 Enhancer A region (NF- κ B consensus site)

Putative NF- κ B binding sites have been identified *in silico* within the Enhancer A regions of all SLA class I promoters with the exception of MIC-2 (Chardon *et al.*, 2001, Renard *et al.*, 2001); the functions of these regions have not been tested experimentally previously. An alignment of Enhancer A consensus from human and porcine class I promoters is shown in figure 5.10. The nucleotide sequences of Enhancer A regions of classical SLA promoters are identical and highly homologous to human classical Enhancer A sequences. Non-classical Enhancer A sequences of SLA-6 and SLA-8 promoters are identical, share overall homology with their human counterparts and only differ from classical SLA Enhancer A sequences by a single trinucleotide. The Enhancer A region of SLA-7 lacks two nucleotides, this may affect the binding of NF- κ B subunits.

The effect of co-expression of NF- κ B p65 and p50 on SLA class I promoters was investigated. Max cells were transiently transfected for 48 hours with the promoter reporter constructs pGL3-SLA-1, -2, -3, -6, -7, or – MIC-2 and pJATLacZ for normalisation. For NF- κ B studies, these cells were also co-transfected with a NF- κ B p65 expression vector. A control reporter plasmid p6 κ B containing 6 tandem repeats of a consensus κ B binding sequence upstream of the firefly luciferase gene was used as a positive control to monitor NF- κ B p65 activity. For NF- κ B p50 studies cells were co-transfected with p50 expression vector or pcDNA3.1. In each experiment normalised fold inductions were calculated as before and statistical

Figure 10: Analysis of Enhancer A sequences from MHC class I promoters

Nucleotide sequences of Enhancer A regions from classical and non-classical SLA and HLA class I were aligned. Human and porcine enhancer a sequences are highly conserved and share considerable homology between species.

Figure 5.11 A and B: Effect of NF- κ B p65 and p50 on SLA class I promoters

Max cells transfected for 48 hrs with pJAT-LacZ and SLA promoter constructs or a p6kB reporter and co-transfected with p65 (A) or p50 (B) expression vectors. Graphs show fold induction of normalised luciferase values compared to a negative control where p50/p65 was substituted with pcDNA3.1 Statistical significance was calculated using Wilcoxon signed ranked tests on paired raw data sets, significant data ($p < 0.05$) are marked with an asterisk.

significance of inductions was calculated using Wilcoxon signed ranked tests on paired raw normalised luciferase data sets.

Figure 5.11A shows that the positive control, 6κB is induced by co-transfection with NF-κB p65. The classical SLA promoters SLA-1, appears to be induced by NF-κB p65, although this could not be proven statistically using this data set. SLA-2 is induced by p65 ($p=0.042$), a mean fold induction of 3.2 was observed. Data presented for SLA-3 is approaching significance ($p=0.068$) indicating that SLA-3 is also induced by p65, an average fold induction of 3.2 suggests a similar magnitude of induction to that of SLA-2 despite having identical Enhancer A nucleotide sequences. A larger standard deviation and smaller sample size is likely to account for the lack of statistical significance. Previous results from this thesis showed that SLA-1 promoter activity is induced by TNF- α (see section 5.3.2), the induction of classical SLA promoters by NF-κB p65 correlates with these results. Therefore it can be confirmed that the putative Enhancer A sequences identified within classical SLA promoters are functional and, like HLA class I promoters, are regulated by NF-κB p65. In contrast to classical SLA class I the non-classical promoters were not induced by NF-κB p65 despite having similar Enhancer A sequences to those reported in classical and HLA class I promoters. A weak induction of SLA-7 activity by TNF- α was observed, this was also seen in some, but not all of the samples tested NF-κB p50 caused different effects on each classical SLA class I promoter tested. It was not possible to show any statistically significant decrease or

increase in promoter activity correlating with p50 co-transfection. Trends emerging from this data are that p50 may induce SLA-2 and repress SLA-3 but does not effect SLA-1. Since the Enhancer A sequences of these promoters are identical, there may be other factors that influence NF- κ B p50 regulation of classical SLA class I genes. The non-classical SLA-6 and -7 and MIC-2 promoters appear to be repressed by p50.

5.4.3 S, X and Y elements within SLA class I promoters

In the human, S, X and Y elements are important in both constitutive and IFN- γ induced transactivation of HLA class I gene expression mediated by CIITA. CIITA acts as a co-activator of transcription, it does not bind DNA directly but mediates gene transcription by interacting with RFX, CREB and NF-Y complexes bound to promoter elements X1, X2 (site α) and Y respectively (Martin *et al.*, 1997). In the pig, S, X and Y boxes have been identified within classical class Ia and non-classical class Ib promoters but appear to be absent within the class Ic promoter MIC-2 (Chardon *et al.*, 2001, Renard *et al.*, 2001). The effect of CIITA on SXY regulatory domains in SLA class I genes has not been investigated previously. An alignment of human and porcine S, X, and Y motifs within classical and non-classical promoters is shown in figure 5.12. Consideration of the sequences of S, X and Y regulatory domains may help explanation any differences in CIITA co-activation between promoters.

Classical SLA S motifs are conserved, identical to HLA-C and differ from HLA-A and -B by a single base. Classical X motifs differ from each

other by a single base only and share overall homology to HLA class I S motifs. The Y motifs of SLA-1 and -2 are identical, and differ from those of classical HLA by a single base. The S regions of the non-classical SLA promoters are less conserved than in classical SLA genes. The SLA-6 S motif is identical to that of HLA-E and -G and the S motif of SLA-8 differs from HLA-F and -G by a single base. The S box of SLA-7 is most divergent. Comparison of the non-classical X motifs with those of non-classical HLA reveals a series of conserved residues, with single base variations at numerous point in the sequences. The effect of these differences upon transcription factor binding remains to be established. By contrast, the Y motifs of the non-classical SLA are all identical to that of HLA-F.

The effect of co-expression of CIITA on SLA class I promoters was investigated. The promoter reporter constructs pGL3-SLA-1, -2, -3, -6, -7, and -MIC-2 were co-transfected into Max cells with pJATLacZ and a CIITA expression vector or pcDNA3.1 for 48 hours. Normalised luciferase values and fold inductions were calculated as standard and statistical significance was calculated using Wilcoxon signed ranked tests on paired raw data sets (figure 5.13).

The classical SLA promoter SLA-1 is induced by co-transfection with CIITA ($p = 0.008$) and a strong fold induction of 4.9 was observed. Similarly, CIITA also appears to induce SLA -2 and -3, and p values on these data approach significance ($p=0.043$ and 0.043 respectively). Chapter 3 of this thesis showed that classical SLA promoter activity was induced by IFN- α . The

Figure 5.12: Analysis of S, X and Y domain sequences from MHC class I promoters

Nucleotide sequences of SXY regions from classical and non-classical SLA and HLA class I were aligned to highlight similarities and differences between human and porcine classical and non-classical genes.

Figure 5.13: Effect of CIITA on SLA class I promoters

Normalised fold induction of SLA promoter activity in Max cells following expression CIITA for 48 hrs. CIITA induces classical but not non-classical SLA promoter activity. The standard deviation (SD) and number of repeats (n) indicated below each data set. Statistical significance was calculated using Wilcoxon signed ranked tests on paired raw data sets. Asterisks indicate significant data ($p < 0.05$).

induction of classical SLA promoters by CIITA, which is induced by IFN- γ , provides supporting evidence for these results. These experiments confirm that the putative S, X, and Y regulatory modules identified within classical SLA promoters are functional and, as with human classical promoters, are regulated by CIITA.

CIITA expression did not induce the non-classical SLA promoters. For the SLA-6 promoter, this correlates with the lack of promoter response to IFN- γ stimulation reported here and may reflect differences in the sequence of the S, X, Y module, particularly in the X2/site- α region. In contrast SLA-7 promoter activity is repressed by CIITA ($p=0.028$) but not by IFN- γ , sequence differences in the S and X regions of SLA-7 may provide a partial explanation for the lack of CIITA activation. Our preliminary observations indicate that both CIITA and IFN- γ repress MIC-2 promoter activity. Since the MIC-2 promoter does not contain S, X or Y regulatory domains this comes as no surprise.

5.5 Viral disruption of SLA class I promoters

5.5.1 CSFV down-regulates the constitutive activity of SLA class I promoters.

Results from this thesis have shown that infection with CSFV down regulates cell surface MHC class I expression on porcine cells (see section 3.3.2). Since modulation of gene transcription at the level of the promoter is a common strategy used by viruses to control the expression of host cell

genes, the activity of SLA class I promoter constructs in CSFV-infected cells was investigated. Max cells were infected with CSFV Brescia for 24 hrs. CSFV-infected and uninfected Max cells were transiently transfected for 48 hrs with pGL3-SLA-1, -2, -7, or pGL3-control. After a total infection time of 72 hpi, cells were harvested and luciferase activity was measured. Previous experiments showed that the constitutive β -galactosidase activity of pJATLacZ was down regulated during CSFV infection and for this reason, luciferase values were not normalised against β -galactosidase activity in these experiments. Constitutive SLA promoter activities in uninfected and CSFV infected Max cells are shown in figure 5.14. At 72 hpi, the basal activity of SLA-1 is significantly down-regulated ($p=0.028$). In addition, the data show a trend that constitutive activity of the classical promoter SLA-2 is reduced. The basal activity of the non-classical promoters SLA-6 and SLA-7 also appear to be decreased and these data are approaching significance ($p=0.068$ for both). In addition, data from these experiments shows a decrease in the constitutive luciferase activity of pGL3-control suggesting that CSFV causes a more generalised disruption of transcription.

Figure 5.14: CSFV down regulates constitutive classical and non-classical SLA class I promoter activity.

Max cells were mock infected or infected with the Brescia strain of CSFV for 24 hrs. Cells were then transiently transfected for a further 48 hrs with pGL3-SLA-1, pGL3-SLA-2, pGL3-SLA-3, pJATLacZ or the positive control pGL3-control (luciferase synthesis driven by the SV40 promoter). Total time for CSFV infection was 72 hrs. This graph shows the luciferase activity of SLA-1, -2, -6, -7 promoters and pGL3-control in CSFV infected cells, shown as percentage expression compared with the equivalent uninfected controls. Normalised RLU were not used as CSFV down-regulated β -galactosidase expression by pJATLacZ (data not shown). Statistical significance was calculated for data (where $n \geq 2$) using Wilcoxon signed ranked tests on paired raw luciferase data. Significant data ($p < 0.05$) are marked with an asterisk.

5.6 Discussion

5.6.1 Summary of results

The results described above have characterised the responses of classical and non-classical SLA class I promoters to cytokines and transcription factors. This chapter has shown that all the classical SLA and non-classical SLA promoters are constitutively expressed and that there are locus-specific responses to cytokines and transcription factors. Table 8 shows a summary of the expected results and table 9 shows the overall results according to SLA gene. In addition, the key regions of SLA class I promoters responsible for transcriptional regulation of SLA class I genes have been analysed in detail and conclusions about their function will now be discussed. First some general but important considerations that should be taken into consideration when interpreting the results reported here will be discussed.

Table 8: Summary of the expected results

Condition	Classical SLA	Non-classical SLA
Constitutive	+++	variable/tissue specific
IFN- α	++++	variable inductions
IFN- γ	++++	variable inductions
TNF- α	+++	variable inductions
TNF- α + IFN- α	+++++	variable inductions
TNF- α + IFN- γ	+++++	variable inductions
IRF-1	++++	variable inductions
IRF-2	++++	variable inductions
IRF-3	++++	variable inductions
IRF-7	++++	variable inductions
IRF-9	++++	variable inductions
p65	++++	variable inductions
p50	-	repression
CIITA	++++	unknown

Table 9: Summary of the SLA promoter response results

Condition	Promoter response to stimulus					
	SLA-1	SLA-2	SLA-3	SLA-6	SLA-7	MIC-2
Constitutive	+++	++	+	+++	++++	++
IFN- α	++	0	nd	0	-	-
IFN- γ	++++	++	++	0	0	-
TNF- α	++	nd	nd	nd	+	nd
TNF- α + IFN- α	++	nd	nd	nd	-	nd
TNF- α + IFN- γ	++++	nd	nd	nd	-	nd
IRF-1	+	nd	+	nd	+	0
IRF-2	+	0	-	nd	0	0
IRF-3	+++	0	nd	nd	0	nd
IRF-7	++	++	nd	nd	0	-
IRF-9	+++	nd	nd	nd	+	0
p65	++	+++	+++	-	0	-
p50	0	+	-	-	-	-
CIITA	+++++	+++++	+++	0	-	-

5.6.2 General considerations

The experiments described above required the transient transfection of cells with up to 4 plasmids, this presents a series of potential technical limitations. Amine transfection reagents are toxic to cells and may affect gene expression. Transient nucleic acid transfection has been shown to induce PKR and RNase L activity and suppress gene expression from viral and non-viral based vectors (Kaufman & Murtha, 1987, Terenzi et al., 1999) and in these studies transfection may affect the expression of co-transfected expression construct genes and reporter genes. In addition there may be competition between plasmids for cellular transcription factors, this may disrupt the expression of co-transfected genes. In future studies, the use of cells stably transfected with SLA class I promoters or may help to minimise

the effects of these factors and have the added advantage that SLA promoters can be studied in a chromatised environment.

The magnitudes of SLA class I promoter inductions following cytokine treatment and expression of transcription factors were considerably lower than those reported in similar studies using human class I promoters (Gobin *et al.*, 1998, 1999). There may be several explanations for this, the most likely being that this study used long promoter regions that contain a number of silencer and enhancer motifs whose impact upon promoter activity and gene expression is not fully understood. Human studies used truncated promoter regions containing motifs such as Enhancer A or ISRE in isolation upstream of a reporter gene; similar reporter constructs should be used in future studies of SLA class I promoters. In addition, studies of HLA class I promoters were compared in a number of cell lines. As discussed previously, only a limited number of porcine cell lines are available. Studies carried out by the Singer lab demonstrated that the mechanisms controlling the regulation of PD1 are conserved in mouse and pig (Ehrlich *et al.*, 1989; Frels *et al.*, 1985), therefore murine cell lines could be an acceptable alternative for future SLA class I promoter studies.

5.6.3 Constitutive activity of SLA promoters in porcine cells

It is important to understand the constitutive activities of SLA class I promoters in order to benchmark changes in activity that take place when induction occurs. Class I promoters that are constitutively expressed at high

levels are probably most important in peptide presentation and as inhibitory receptors to prevent NK cell activation. The consistent trend observed for constitutive expression was: SLA-7 had the highest constitutive activity followed by SLA-1 and SLA-6 and then SLA-2 and MIC-2, SLA-3 had the lowest constitutive activity. Constitutive activity of the classical gene promoters is consistent with characteristic ubiquitous expression that is associated with classical genes. Despite sharing highly homologous promoter sequences, there is locus specific variation in the constitutive activities of the classical SLA class I genes. Differences in constitutive activity may arise from binding of different transcription factor and regulatory promoter domains outside of the regions studied in this thesis such as distal enhancer and silencer sites, which may account for the differences observed. A comparison of the transcription factor binding profiles of each classical gene in resting cells could be used to explore this theory. It is noteworthy that in the H01 haplotype, an interspersed repeat or "SINE" separates the enhancer from silencer of SLA-2 and lengthens the promoter of SLA 3 (Renard *et al.*, 2001). The fact that SINE is absent from the SLA-1 promoter may explain why this gene is more constitutively active than the others. It is possible that the SINE changes the configuration of the DNA and hinders transcription. Not only does the location of the SINE vary from loci to loci, but also from haplotype to haplotype as described for the H04 (d/d) pig (Maguire *et al.*, 1990).

Given that mRNA levels of PD1 expression have been shown to differ dramatically between tissues, it is almost certain that the levels of constitutive

activity of H01 classical genes would differ among tissues; SLA-1 may not always be the most highly expressed. It would therefore be interesting to compare the constitutive activity of these promoters in cells derived from several different tissues; in particular comparison of non-immune and immune tissues such as the spleen, thymus and lymph nodes would be useful. Unfortunately, only a number of porcine cell lines are available, cell lines of defined haplotypes are particularly rare and transfection of primary cell lines is technically challenging. Analysis of classical gene expression using quantitative RT-PCR on a range of cell types would compliment promoter studies.

Strong constitutive activity of the promoters of the non-classical genes SLA-6 and SLA-7 was observed and these results correspond with the constitutive mRNA expression of these genes reported in the kidney (Crew *et al.*, 2004). In the light of this recent data, it would be interesting to compare constitutive expression of these promoters in other tissues including those found to be negative for mRNA expression of SLA-6 and SLA-7. One pertinent question that remains is what is the functional equivalent of HLA-G in the pig? To address this, a study of expression of non-classical genes in cells at the materno-foetal interface is required.

5.6.4 Interferon induces classical SLA class I promoter activity

Induction of classical MHC class I by type I and II IFNs has been well documented. The data presented here indicate that IFN- α and IFN- γ also

induce classical SLA class I promoter activity and, although the data presented approach significance, induction of these genes by IFNs could not be proven. One exception is SLA-1, which was significantly induced by IFN- γ .

The experimental use of IFNs is complicated by the inherent function of these cytokines as mediators of anti-viral responses, which have been well documented. Type I and II IFNs induce the RNA-dependent protein kinase (PKR) activity; this enzyme phosphorylates the α subunit of the eukaryotic initiation factor eIF2, leading to an inhibition of protein translation. In the experimental system used in this study there is evidence that inhibition of translation is taking place; luciferase activity of the SV40 driven luciferase gene (pGL3-control vector), which should be constitutively active, was repressed in IFN- α /- γ treated cells. IFN- α is a weaker inducer of PKR (S. Goodbourn, personal communication) and accordingly, the decrease in constitutive luciferase activity of SV40-luciferase was less than that observed with IFN- γ . Induction of PKR is therefore a plausible explanation for the lack of, or low levels of IFN- α induction seen for classical SLA class I promoters and the variation in inductions that prevents this data from reaching statistical significance. In future studies, inhibition of eIF2 α could be monitored by western blot.

IFN- γ strongly induces SLA-1 promoter activity and appears to induce SLA-2 and SLA-3 albeit to a lower magnitude. This correlates to the

presence of a putative STAT binding site in each classical promoter. The fold inductions following IFN- γ treatment presented for SLA-2 and SLA-3 are smaller than those presented for SLA-1, which may explain why these data approach, but do not reach significance. Comparison of the mean fold inductions caused by each IFN species shows that IFN- γ induces SLA promoter activity more strongly than IFN- α . Both IFN- α and - γ induce the expression of families of transcription factors such as the IRFs which have the capacity to bind ISRE of SLA class I promoters and induce transcription. In addition, IFN- γ induces class I expression *via* the transcription factor CIITA, which interacts with the SXY motifs. Induction of transcription factors which bind distinct regions of the class I promoter may confer IFN- γ with an added potency for MHC class I induction. Interestingly, IFN- γ also caused a significant decrease in luciferase production in control experiments using the pGL3-control vector and this decrease is less than that observed with IFN- α . Taking this into consideration, IFN- γ could be a more potent inducer of class I in this system than IFN- α because it is a weaker inducer of PKR activity than IFN- α . A combination of increased potency and weaker induction of PKR are likely to be accountable for the differences in IFN- γ and IFN- α fold inductions reported here. Experimental quantification of PKR activity would be required to exemplify this theory.

Overall these results suggest that there are locus specific differences in IFN- α responses of classical SLA class I genes. Since the classical SLA promoters contain consensus ISRE sites one can assume that in common

with humans, this region is functional and responsible for interferon-mediated regulation of classical SLA expression.

5.6.5 Non-classical promoter activity was not induced by IFNs

This study demonstrated that the promoters of the non-classical genes SLA-6 and -7 and the promoter of the class I c gene MIC-2 did not respond to IFN- α or - γ . This is in contrast to human studies where IFN- γ induced HLA-E and HLA-F (Gobin & van den Elsen, 2000) and therefore SLA class Ib genes are either not regulated by IFN- α/γ or inductions by these IFNs could not be detected using this experimental method. As discussed above it is possible that induction of PKR and RNase L has masked the effect of IFN- α/γ . For clarification, these experiments should be repeated using truncated promoters containing only the putative ISRE elements in parallel with measuring mRNA expression by quantitative RT-PCR in IFN- α/γ treated cells. In these studies the SLA-8 promoter fused to a luciferase reporter gene would be useful as a negative control because SLA-8 it lacks an ISRE and is therefore not predicted to respond to IFNs.

In this study IFN- γ did not induce non-classical SLA promoters. *In silico* analysis of SLA-6, -7 and -8 promoters identified a single putative STAT binding site within the promoter region of SLA-6, this site requires further characterisation to confirm its function in SLA-6. No equivalent site was found in the promoters of SLA-7 and SLA-8; the lack of a STAT binding

motif may explain the lack of response to IFN- γ observed with these two promoters.

5.6.6 Regulation of SLA class I promoters by Interferon Regulatory Factors

IFN- α/γ mediated induction of SLA class I expression was explored in more detail by analysing promoter responses to co-transfection with members of the IRF family of transcription factors. As discussed previously, these transcription factors are induced by IFNs and bind to consensus sequences within the ISRE of IFN-sensitive genes. Binding of IRFs to the ISRE regions of classical HLA class I gene has been described previously (Girdlestone *et al.*, 1993, Lefebvre *et al.*, 2001, Massa & Wu, 1995). *In silico* analysis of the ISRE of SLA-1, -2 and -3 promoters revealed consensus sequences for IRF transcription factor binding (Chardon *et al.*, 2001; Renard *et al.*, 2001) and therefore, in common with their human counterparts, the ISRE of classical SLA genes should bind and be induced by IRFs. Co-transfection of the SLA-1 promoter with an IRF-1 expression vector results in an induction of SLA-1 driven luciferase production demonstrating that the ISRE of SLA-1 is functional and is induced by IRF-1. Similar studies using SLA-3 suggest that the ISRE of this promoter is also functional and induced by IRF-1 and these data approach significance.

The nucleotide sequence of the ISRE of SLA-7 differs from that of SLA-1 and HLA-A at two points; there is a C to G inversion and a C to T

insertion within in the 5' region. It was predicted that these nucleotide differences might have an adverse effect upon IRF binding and prevent transcriptional induction of this gene by these transcription factors. However, co-transfection of SLA-7 with IRF-1 caused an increase in SLA-7 promoter activity, this was in contrast to experiments with IFN- α/γ where no induction of SLA-7 was observed. As discussed above, induction of PKR may mask the effects of IFN- α/γ .

The nucleotide sequence of the ISRE region of the SLA-6 promoter also differs from those of classical and HLA class I genes and contains an insertion of two nucleotides. In future experiments, this promoter would be tested using the same experimental model to determine the effects of this insertion on IRF binding and responses to IFN- α .

In order to further examine the interaction of IRF transcription factors with the ISRE of SLA class I genes, the effect of IRF-2, -3, -7 and -9 was investigated. All these IRFs bind to the same DNA sequence and with the exception of IRF-2, these transcription factors are known transcriptional activators of ISG's. Accordingly, SLA-1 is induced by IRF-3 and IRF-9 and may be induced by IRF-7. SLA-2 was not induced by IRF-3 but appears to be induced by IRF-7 although significance was not reached by these data, perhaps due to the small sample size tested. SLA-7 was not induced by IRF-3 or IRF-7 but was, in some cases, induced by IRF-9, although these data did not reach significance. This is in contrast to IRF-1, which did induce SLA-7.

Overall, IRF-3 and IRF-7 did not behave as anticipated, they should have induced classical SLA class I transcription. One possible explanation is that IRF-3 and IRF-7 need to be activated in order to translocate to the nucleus and exert their effects on gene transcription. Co-transfection of these transcription factors alone may not be sufficient for their activation. Previous studies have shown that transfection of IRF-3 and IRF-7 cDNAs induced IFN α 4 and α 6 promoter activity, induction of these promoters was much stronger when cells were infected with virus (Marie *et al.*, 1998). Therefore it is likely that activation of IRF-3 and IRF-7 by viral infection or stimulation with IFNs or synthetic dsRNA would substantially increase SLA promoter responses to these transcription factors. In this study a constitutively active form of IRF-9 was used, and accordingly SLA-1 was significantly induced.

IRF-2 is usually a transcriptional repressor of ISG's therefore it was predicted that IRF-2 would repress SLA class I promoter activity. With the exception of SLA-3, however, there were no clear trends of transcriptional repression and rather unexpectedly, IRF-2 appeared to induce SLA-1 promoter activity. These experiments produced variable results and contained small sample sizes, this is reflected by considerable variation in the results obtained from these experiments and accordingly, statistical significance was not reached for any of the classical promoters studied

These results have described the gross effect of cytokines and TFs on full length SLA class I promoters, which have been related to putative

regulatory motifs identified within promoter regions. Future experiments should use truncated SLA promoters lacking upstream silencer elements and containing only the ISRE region. In addition, binding of IRFs to the ISRE should be confirmed using band shift assays and northern blots.

5.6.7 Transcriptional regulation by TNF- α and NF- κ B

TNF- α induces activation of NF- κ B species which bind to kB motifs located within Enhancer A promoter regions and induce transcription of MHC class I genes (Gobin *et al.*, 1998, Johnson, 2003). In this study, TNF- α appears to induce SLA-1 promoter activity and is shown to induce the SLA-7 promoter significantly, suggesting that the Enhancer A regions of these genes, which share homology to the consensus sequence found in human HLA class I promoters, are functional. Results from co-transfection of classical SLA promoters with NF- κ B p65 studies also indicated that the Enhancer A regions of these genes are functional. A significant induction of SLA-2 was obtained following co-transfection of p65, but large standard deviation prevented significance being reached in similar experiments using SLA-1 and SLA-3. Given that the sequences of the Enhancer A region of all three classical SLA are identical, it can be assumed that they are functional on the basis of the data presented.

In humans, TNF- α has been shown to synergistically induce MHC class I with IFN- α and - γ , mediated by co-operation of IRF-1 and NF- κ B p65 at the promoter level (Drew *et al.*, 1995b, Johnson, 2003). In this study, no

clear synergistic inductions of SLA class I genes were observed using these cytokines. As discussed above, induction of PKR and RNase L by interferons may mask synergistic activation.

5.6.8 CIITA induces classical SLA class I promoter activity

CIITA is a transcriptional activator of HLA class I and class II genes and mediates this effect *via* co-operative binding of RFX/CREB/NFY which are bound to the SXY region of these promoters. Until now, the effect of CIITA on SLA class I had not been examined, these data showed that CIITA is a transcriptional activator of classical SLA class I genes. Interestingly, the non-classical genes SLA-6 and SLA-7 were not induced by this transcription factor. The nucleotide sequences of the S and X1 regions of the SLA-7 promoter are noticeably different; this probably affects binding of RFX and CREB and in turn, may explain the lack of CIITA induction observed. Similarly, the X1 and X2 regions of the SLA-6 promoter contain nucleotide sequences that are divergent from those of classical SLA which may disrupt CREB binding and prevent trans-activation by CIITA. To confirm that this is the case, the SXY regions of SLA-6 and -7 should be examined in isolation to determine which transcription factors bind them and compared with classical SLA promoters. MIC-2 was not induced by CIITA, this was as expected since this promoter does not contain an SXY box.

5.6.9 Regulation of MIC-2 expression

The regulation of the MIC-2 promoter requires separate consideration since in the experiments using the MIC-2 promoter reporter described in this chapter, none of the cytokines or transcription factors used caused an increase in MIC-2 promoter activity. The porcine MIC-2 promoter contains a region that resembles the heat shock element (HSE) found in human MIC promoters, in humans this region directs transcription of the MIC-A and MIC-B genes in response to heat stress at 42 °C. In the pig, MIC-2 expression has been shown to be up-regulated in response to heat shock but this observation was not confirmed by promoter analysis (Garcia-Borges *et al.*, 2005). Our attempts to repeat these heat shock experiments in Max cells using the MIC-2 promoter reporter construct did not detect an increase in MIC-2 promoter activity (data not shown). The MIC-2 promoter used in this study contained the sequence originally identified by Chardon *et al* (2001) which was identified by comparison of porcine MIC-2 exons with those of the human MIC genes. When the inter-genetic sequences were compared they could not be aligned with the exception of a 1.2kb region upstream of the MIC-2 coding sequence. Since this was the only conserved region found, it was presumed to be important for regulation of the MIC-2 gene. In this study this putative promoter was constitutively active but failed to respond to any of the transcriptional activators tested and therefore, it is likely that this region is not the MIC-2 promoter but does contain a regulatory sequence with the capacity to direct the transcription of the luciferase reporter gene. A second,

putative promoter region has been identified which is positioned several kilo bases upstream of the MIC-2 promoter sequence predicted by Chardon *et al* (Mark Crew, unpublished data). Whether this sequence actually represents the true MIC-2 promoter remains to be established.

5.6.10 CSFV down-regulates basal SLA class I promoter activity

Down-regulation of SLA class I expression by CSFV is a novel immune evasion strategy that has not been described previously for this virus. In order to determine if this down-regulation represents a block in SLA class I gene transcription, the effect of CSFV upon SLA class I promoter activity was examined. This study has shown that CSFV significantly decreases the constitutive activity of SLA-1 and a trend which strongly suggests that CSFV down-regulates the activity of SLA-2, SLA-6 and SLA-7 promoters. In addition, a decrease in the basal activities of the luciferase and β -galactosidase control vectors appears to be caused by CSFV. Both these vectors are under the control of a constitutively active promoter and the level of their expression should not change during viral infection unless viral infection disrupts the availability of transcription factors or host cell transcriptional machinery, suggesting that CSFV disrupts host cell transcription at a global level early during infection.

Host cell shutoff and cessation of host cell transcription is a mechanism that some viruses employ, CSFV however, causes a persistent infection and does not cause cell death in culture, therefore these

observations cannot be explained by host-cell shut off. Instead, it is possible that CSFV targets several transcription factors involved in transcription. This is currently being investigated in this laboratory and preliminary experiments have shown that both NF- κ B p65 and TFIIB are down regulated during CSFV infection. In a previous study by this laboratory, CSFV was shown to prevent IRF-3 binding to the IFN- α promoter, resulting in inhibition of IFN- α synthesis (Larocca *et al.*, 2005). Alternatively, CSFV may disrupt components of the host cell transcriptional machinery such as RNA polymerase activity.

The next question to address is which viral proteins are responsible for blocking SLA class I transcription. Stable cell lines that express individual CSFV proteins including E0 and Erns are available and a Max cell line stably transfected with N^{pro} was produced as part of this study. These cells were analysed by FACS and transfection with SLA class I promoters; unfortunately no effect on SLA class I expression/transcription could be attributed to a single viral protein at this early stage (results therefore were not included in this thesis). It is important that this work continues, as understanding the viral proteins and mechanism responsible for SLA class I down-regulation will allow a greater understanding of the molecular pathogenesis of CSFV and may lead to the identification of therapeutic points of intervention.

5.6.11 Summary and future perspectives

Taking the results and discussion from this study into consideration, a model of the current understanding of the transcriptional regulation of

classical and non-classical SLA class I genes is presented (figure 5.15). In this figure, statistically significant results from this thesis are shown as coloured shapes, results that showed a trend of induction, approach significance and can be justified by similar studies in human are indicated by grey shapes. In summary, the transcriptional regulation of SLA class I genes in response to cytokines is controlled by four key promoter domains; a putative STAT binding motif which mediates responses to IFN- γ , the Enhancer A region containing two κ B binding motifs that mediate responses to TNF- α upon binding of NF- κ B subunits, the ISRE which contains IRF binding motifs and mediates responses to type I IFN and the SXY domain, which is predicted to be bound by a multiprotein complex containing RFX, CREB and NF-Y and is transactivated by CIITA. Locus specific responses to cytokines and transcription factors can be attributed, in part, to sequence variation within these four key domains.

The sequences of these promoter domains in classical SLA class I share homology with each other as well as their human counterparts and accordingly, these promoters are responsive to IFN- γ , NF- κ B, IRFs and CIITA. Responses to TNF- α and NF- κ B are presumed from trends indicated by the data presented. Given that SLA-2 and -3 contain STAT binding sites, that cell surface SLA class I substantially increases following IFN- γ treatment, that human MHC class I promoters containing STAT binding motifs respond to IFN- γ and that trends of IFN- γ induction were observed in SLA-2 and -3 promoter studies, it is presumed that these motifs are functional. Further

Figure 5.15: Model of the transcriptional regulation of SLA class I genes

Classical and non-classical gene expression was investigated at the molecular level using promoter regions upstream of three classical class I genes SLA-1, -2 and -3, two non-classical genes SLA 6 and -7 and the MHC class I chain related gene MIC-2, cloned up stream of a luciferase reporter. Four core promoter regions are shown: STAT binding motif, Enhancer A, ISRE and SXY. This figure summarises the results of this thesis and shows the transcription factors that drive SLA class I transcription in response to cytokine stimulation and the motifs which they interact with. Coloured shapes show statistically proven responses from this study, grey shapes show indicative trends of induction requiring further validation.

validation of this model could be achieved using promoters that have been dissected into individual regions and tested individually. These reagents could be used to determine the response of individual promoter elements to transcription factor and cytokine stimulation as well as to determine the physical binding of transcription factors to isolated DNA sequences. Finally, if time and money were of no object, these investigations should be repeated using cell lines stably transfected with each SLA class I promoter, in order to examine promoter responses in a chromatised environment.

Chapter 6. Conclusion

The porcine immune system has attracted a considerable amount of interest over the last decade due to the economic importance of this species in the food industry as well as the potential for pigs to provide solid organ and cell replacements for humans. The genes encoded in the major histocompatibility complex are central to both of these interests as transplantation antigens and as mediators of immune responses. Therefore, a greater understanding of their functions and how these genes are regulated is important to establish. The aim of this project was to obtain a more detailed understanding of how SLA class I gene expression is regulated. Given the importance of MHC class I in the immune response, selective pressure has maintained MHC class I function throughout evolution. The structure of MHC molecules is vital for correct function of the MHC genes and accordingly, have remained highly conserved. The expression of these genes, however, is variable. Since the majority of variation lies within the promoter region a greater understanding of how the SLA promoter operates is the key to understanding SLA class I expression and therefore in this project, the regulation of SLA class I promoters were studied in detail.

I have demonstrated that IFN- α and - γ increase cell surface SLA class I expression on d/d PAECs and Max cells, in particular, one allele or group of alleles were identified that are particularly sensitive to stimulation with IFN- α and - γ which justifies further investigation. The tools for analysing SLA class I are few in number. Additional limitations are that the identities of the alleles

recognised by each antibody is not known and cell lines available from animals that express a restricted set of SLA class I are not readily available. Furthermore, while primary cells provide a more accurate representation of *in vivo* SLA class I expression than transformed cell lines, there is continued emphasis on reducing the use of animals for experimentation (for example the 3R's initiative). As such, Max cells provide an attractive alternative to primary cells for studies of the regulation of SLA class I expression. In order to fully understand cell surface SLA class I expression more analytical tools are required. Stable transfection of cell lines with individual SLA class I genes could be used to characterise existing SLA class I antibodies and to study expression of SLA class I genes in isolation in a chromatized environment and in response to cytokine stimulation. In addition, these cell lines could be used to determine the nature of the peptides bound by each class I molecule, this would be particularly interesting for non-classical SLA, vaccine design and tetramer production. A second possibility would be to raise new antibodies against short fragments of SLA class I proteins specific to each class I protein, but this is complicated by the high homology of SLA class I alleles. Finally, as none of the existing antibodies available are able to detect class I MHC by Western blotting, generation of an antibody with this capability would be a valuable reagent for studying SLA class I protein expression in the future.

RT-PCR is commonly used to study gene expression at the mRNA level and has been used by other groups to study SLA class I expression (see Crew *et al.*, 2004). An extensive collection of tissue samples derived from

d/d pigs were collected as part of this study, including tissues from neonatal piglets, placental tissues, tissues from immune organs and tissues which have not previously been studied in the d/d pig to date. These tissues are a valuable resource for conducting a detailed tissue survey of SLA class I expression using the RT-PCR primers and conditions described by Crew *et al* (2004). Importantly, examination of non-classical SLA expression in placental tissues may allow a functional equivalent of HLA-G to be found in the pig, which remains an important question.

Studying the promoter regions of SLA class I has permitted a greater understanding of the regulation of class I genes at the molecular level and helps to explain locus specific differences in levels of class I expression. Four regulatory motifs within SLA class I promoters mediate transcriptional responses to cytokines and transcription factors. Future studies should examine which transcription factors physically bind to these motifs both constitutively and following cytokine stimulation in order to confirm the results. In addition, SLA class I promoter motifs should be cloned upstream of reporter genes and so that the transcriptional responses of individual promoter motifs to cytokine and transcription factors can be assessed. In addition, a detailed comparison of classical and non-classical SLA promoters should be made. Finally, reporter constructs should be made which contain individual regulatory motifs in isolation, and responses to cytokines and specific transcription factors analysed. In particular, the promoter region of SLA-5 has not been examined, these are important experiments because the expression of SLA-5 and the function of this gene as a *bona fide* classical

SLA is not known. Similarly, the promoter region of SLA-8 must be examined in order to gain a better understanding of the transcription of this non-classical gene. Perhaps the most interesting aspect of this study was the examination of non-classical SLA-6 and -7 promoters, as the functions of these genes in the pig are not well understood.

The additional tools described above would also be valuable for studying SLA class I expression during viral infections. In this study, the affect of CSFV infection on SLA class I was investigated. Classical swine fever is a constant threat to the UK and European herd and outbreaks of disease cause substantial economical losses; the last outbreak in Europe is estimated to have cost US\$ 2.3 billion and required the slaughter of 11 million pigs (Oleksiewicz *et al.*, 2003). It is therefore imperative that mechanisms of controlling this disease are developed. In order to achieve the ultimate goal of providing a safe, efficient and inexpensive vaccine, a greater understanding of the virus, the porcine immune system and crucially how the virus interacts with the porcine immune system, must be gained. The continuous interactions between host and pathogens during their co-evolution have not only shaped the immune system, but also the counter measures used by pathogens. Down-regulation of cell surface MHC class I expression is a common immune evasion strategy employed by viruses to avoid recognition by CTLs and selective disruption of specific MHC class I alleles allows a virus to avoid NK cell lysis (see Lilley & Ploegh, 2005 for a review). This project addressed two important questions; does CSFV disrupt SLA class I expression ? and if so, how ? Using FACS, these studies have

shown that SLA class I expression on Max cells is down-regulated early during viral infection, within 24 hours. This thesis also showed that SLA class I promoter activity was compromised during CSFV infection and conclude that down regulation of cell surface SLA class I expression is achieved by inhibition of SLA class I transcription. Many viruses target stages of the peptide presentation pathway in order to prevent MHC expression. It is very likely that CSFV targets multiple steps in the peptide presentation pathway, such as inhibition of TAP or ER retention to prevent SLA class I from reaching the cell surface, this should be explored in future studies by analysing expression, activity and trafficking of individual components of the peptide presentation pathway. The next question to answer is which viral protein allows CSFV to disrupt SLA class I ? This is an important question because targeting specific viral proteins may allow SLA class I expression to be restored therapeutically. Max cell lines stably expressing CSFV viral proteins are now being produced and studied so that the effect of individual CSFV proteins upon class I expression can be examined.

Some unexpected results from this study were that CSFV down regulates the expression of two reporter genes under the control of constitutively active promoters including SV40 driven luciferase and β -galactosidase driven by a rat β -actin promoter, indicating that CSFV modulates host cell transcription at a global level. These results support results from previous and ongoing studies in this laboratory which have shown that CSFV down-regulates the transcription of IRF-3, NF- κ B and

TFIIB. This suggests that CSFV either disrupts part of the transcriptional machinery or inhibits global transcription factors. Understanding CSFV immune evasion strategies is important work that must be continued if a solution to this virus is to be developed.

Chapter 7. References

- Abendroth, A., Lin, I., Slobedman, B., Ploegh, H. & Arvin, A. M. (2001). Varicella-zoster virus retains major histocompatibility complex class I proteins in the Golgi compartment of infected cells. *Journal of Virology* **75**, 4878-4888.
- Aki, M., Shimbara, N., Takashina, M., Akiyama, K., Kagawa, S., Tamura, T., Tanahashi, N., Yoshimura, T., Tanaka, K., Ichihara, A. (1994). Interferon-gamma induces different subunit organizations and functional diversity of proteasomes. *Journal of Biochemistry* **115**, 257-269.
- Alexopoulou, L., Holt, A. C. and Flavell, R. A. (2001). Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* **413**(6857), 732-738.
- Allan, D. S., Colonna, M., Lanier, L. L., Churakova, T. D., Abrams, J. S., Ellis, S. A., McMichael, A. J. & Braud, V. M. (1999). Tetrameric complexes of human histocompatibility leukocyte antigen (HLA)-G bind to peripheral blood myelomonocytic cells. *Journal of Experimental Medicine* **189**, 1149-1156.
- Au, W. C., Moore, P. A., Lowther, W., Juang, Y. T. & Pitha, P. M. (1995). Identification of a member of the interferon regulatory factor family that binds to the interferon-stimulated response element and activates expression of interferon-induced genes. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 11657-11661.
- Bahram, S. (2000). MIC genes: from genetics to biology. *Advances in Immunology* **76**, 1-60.
- Bastians, H., Topper, L. M., Gorbsky, G. L. & Ruderman, J. V. (1999). Cell cycle-regulated proteolysis of mitotic target proteins. *Molecular Biology of the Cell* **10**, 3927-3941.
- Bauer, S., Groh, V., Wu, J., Steinle, A., Phillips, J. H., Lanier, L. L. & Spies, T. (1999). Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science* **285**, 727-729.
- Bensaude, E., Turner, J. L., Wakeley, P. R., Sweetman, D. A., Pardieu, C., Drew, T. W., Wileman, T. & Powell, P. P. (2004). Classical swine fever virus induces proinflammatory cytokines and tissue factor expression and inhibits apoptosis and interferon synthesis during the establishment of long-term infection of porcine vascular endothelial cells. *Journal of General Virology* **85** 1029-1037.

- Bjorkman, P. J., Saper, M. A., Samraoul, B., Bennett, W. S., Strominger, J. L. & Wiley, D. C. (1987). Structure of the human class I histocompatibility antigen, HLA-A2. *Nature* **329**, 506-512.
- Bogdan, C., Mattner, J. & Schleicher, U. (2004). The role of type I interferons in non-viral infections. *Immunological Reviews* **202**, 33-48.
- Bowie, A. G. & Haga, I. R. (2005). The role of Toll-like receptors in the host response to viruses. *Molecular Immunology* **42**, 859-867.
- Braun, B. C., Glickman, M., Kraft, R., Dahlmann, B., Kloetzel, P. M., Finley, D. & Schmidt, M. (1999). The base of the proteasome regulatory particle exhibits chaperone-like activity. *Nature Cell Biology* **1**, 221-226.
- Carrasco, C. P., Rigden, R. C., Vincent, I. E., Balmelli, C., Ceppi, M., Bauhofer, O., Tache, V., Hjertner, B., McNeilly, F., van Gennip, H. G., McCullough, K. C. & Summerfield, A. (2004). Interaction of classical swine fever virus with dendritic cells. *Journal of General Virology* **85**, 1633-1641.
- Cerwenka, A., Bakker, A. B., McClanahan, T., Wagner, J., Wu, J., Phillips, J. H. & Lanier, L. L. (2000). Retinoic acid early inducible genes define a ligand family for the activating NKG2D receptor in mice. *Immunity* **12**, 721-727.
- Chang, C. H., Hammer, J., Loh, J. E., Fodor, W. L. & Flavell, R. A. (1992). The activation of major histocompatibility complex class I genes by interferon regulatory factor-1 (IRF-1). *Immunogenetics* **35**, 378-384.
- Chapman, T. L., Heikeman, A. P. & Bjorkman, P. J. (1999). The inhibitory receptor LIR-1 uses a common binding interaction to recognize class I MHC molecules and the viral homolog UL18. *Immunity* **11**, 603-613.
- Chardon, P., Renard, C., Vaiman, M. (1999). The major histocompatibility complex in swine. *Immunological Reviews* **167**, 179-182.
- Chardon, P., Renard, C., Gaillard, C. R., Vaiman, M. (2000). The porcine major histocompatibility complex and related paralogous regions: a review. *Genetics Selection Evolution* **32**, 109-128.
- Chardon, P., Rogel-Gaillard, C., Cattolico, L., Duprat, S., Vaiman, M. & Renard, C. (2001). Sequence of the swine major histocompatibility complex region containing all non-classical class I genes. *Tissue Antigens* **57**, 55-65.

- Charley, B., Lavenant, L. & Delmas, B. (1991). Glycosylation is required for coronavirus TGEV to induce an efficient production of IFN alpha by blood mononuclear cells. *Scandinavian Journal of Immunology* **33**, 435-440.
- Cosman, D., Mullberg, J., Sutherland, C. L., Chin, W., Armitage, R., Fanslow, W., Kubin, M. & Chalupny, N. J. (2001). ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the NKG2D receptor. *Immunity* **14**, 123-133.
- Crew, M. D., Phanavanh, B. & Garcia-Borges, C. N. (2004). Sequence and mRNA expression of nonclassical SLA class I genes SLA-7 and SLA-8. *Immunogenetics* **56**, 111-114.
- Davis, W. C., Marusic, S., Lewin, H. A., Splitter, G. A., Perryman, L. E., McGuire, T. C. & Gorham, J. R. (1987). The development and analysis of species specific and cross reactive monoclonal antibodies to leukocyte differentiation antigens and antigens of the major histocompatibility complex for use in the study of the immune system in cattle and other species. *Veterinary Immunology and Immunopathology* **15**, 337-76.
- Didcock, L., Young, D. F., Goodbourn, S. & Randall, R. E. (1999). Sendai virus and simian virus 5 block activation of interferon-responsive genes: importance for virus pathogenesis. *Journal of Virology* **73**, 3125-3133.
- Diefenbach, A., Jamieson, A. M., Liu, S. D., Shastri, N. & Raulet, D. H. (2000). Ligands for the murine NKG2D receptor: expression by tumor cells and activation of NK cells and macrophages. *Nature Immunology* **1**, 119-126.
- Dissen, E., Ryan, J. C., Seaman, W. E. & Fossum, S. (1996). An autosomal dominant locus, Nka, mapping to the Ly-49 region of a rat natural killer (NK) gene complex, controls NK cell lysis of allogeneic lymphocytes. *Journal of Experimental Medicine* **183**, 2197-2207.
- Dohring, C., Scheidegger, D., Samaridis, J., Cella, M. & Colonna, M. (1996). A human killer inhibitory receptor specific for HLA-A1,2. *Journal of Immunology* **156**, 3098-3101.
- Drew, P. D., Franzoso, G., Carlson, L. M., Biddison, W. E., Siebenlist, U. & Ozato, K. (1995a). Interferon regulatory factor-2 physically interacts with NF-kappa B in vitro and inhibits NF-kappa B induction of major histocompatibility class I and beta 2-microglobulin gene expression in transfected human neuroblastoma cells. *Journal of Neuroimmunology* **63**, 157-162.

- Drew, P. D., Franzoso, G., Becker, K. G., Bours, V., Carlson, L. M., Siebenlist, U. & Ozato, K. (1995b). NF kappa B and interferon regulatory factor 1 physically interact and synergistically induce major histocompatibility class I gene expression. *Journal of Interferon and Cytokine Research* **15**, 1037-1045.
- Ehrlich, R., Lifshitz, R., Pescovitz, M. D., Rudikoff, S. & Singer, D. S. (1987). Tissue-specific expression and structure of a divergent member of a class I MHC gene family. *Journal of Immunology* **139**, 593-602.
- Ehrlich, R., Maguire, J. E. & Singer, D. S. (1988). Identification of negative and positive regulatory elements associated with a class I major histocompatibility complex gene. *Molecular Cell Biology* **8**, 695-703.
- Ehrlich, R., Sharrow, S. O., Maguire, J. E. & Singer, D. S. (1989). Expression of a class I MHC transgene: effects of in vivo alpha/beta-interferon treatment. *Immunogenetics* **30**, 18-26.
- Fisk, B., Ioannides, C. G., Aggarwal, S., Wharton, J. T., O'Brian, C. A., Restifo, N. & Glisson, B. S. (1994). Enhanced expression of HLA-A,B,C and inducibility of TAP-1, TAP-2, and HLA-A,B,C by interferon-gamma in a multidrug-resistant small cell lung cancer line. *Lymphokine and Cytokine Research* **13**, 125-131.
- Frels, W. I., Bluestone, J. A., Hodes, R. J., Capecchi, M. R. & Singer, D. S. (1985). Expression of a microinjected porcine class I major histocompatibility complex gene in transgenic mice. *Science* **228**, 577-580.
- Frels, W. I., Bordallo, C., Golding, H., Rosenberg, A., Rudikoff, S. & Singer, D. S. (1990). Expression of a class I MHC transgene: regulation by a tissue-specific negative regulatory DNA sequence element. *New Biologist* **2**, 1024-1033.
- Fruh, K. & Yang, Y. (1999). Antigen presentation by MHC class I and its regulation by interferon gamma. *Current Opinion in Immunology* **11**, 76-81.
- Gagnier, L., Wilhelm, B. T. & Mager, D. L. (2003). Ly49 genes in non-rodent mammals. *Immunogenetics* **55**, 109-115.
- Garcia-Borges, C. N., Phanavanh, B., Saraswati, S., Dennis, R. A. & Crew, M. D. (2005). Molecular cloning and characterization of a porcine UL16 binding protein (ULBP)-like cDNA. *Molecular Immunology* **42**, 665-671.

- Geffrotin, C., Popescu, C. P., Cribiu, E. P., Boscher, E. P., Renard, C., Chardon, P. & Vaiman, M. (1984). Assignment of MHC in swine to chromosome 7 by in situ hybridization and serological typing. *Animal Genetics* **27**, 213-219.
- Girdlestone, J., Isamat, M., Gewert, D. & Milstein, C. (1993). Transcriptional regulation of HLA-A and -B: differential binding of members of the Rel and IRF families of transcription factors. *Proceedings of the National Academy of Sciences of the United States of America* **90**, 11568-11572.
- Gobin, S. J. & van den Elsen, P. J. (2000). Transcriptional regulation of the MHC class Ib genes HLA-E, HLA-F, and HLA-G. *Human Immunology* **61**, 1102-1107.
- Gobin, S. J., Biesta, P. & Van den Eisen, P. J. (2003). Regulation of human beta 2-microglobulin transactivation in hematopoietic cells. *Blood* **101**, 3058-3064.
- Gobin, S. J., Keijzers, V., van Zutphen, M. & van den Elsen, P. J. (1998). The role of enhancer A in the locus-specific transactivation of classical and nonclassical HLA class I genes by nuclear factor kappa B. *Journal of Immunology* **161**, 2276-2283.
- Gobin, S. J., van Zutphen, M., Woltman, A. M. & van den Elsen, P. J. (1999). Transactivation of classical and nonclassical HLA class I genes through the IFN-stimulated response element. *Journal of Immunology* **163**, 1428-1434.
- Goh, K. C., deVeer, M. J. & Williams, B. R. (2000). The protein kinase PKR is required for p38 MAPK activation and the innate immune response to bacterial endotoxin. *EMBO Journal* **19**, 4292-4297.
- Gonen-Gross, T., Achdout, H., Gazit, R., Hanna, J., Mizrahi, S., Markel, G., Goldman-Wohl, D., Yagel, S., Horejsi, V., Levy, O., Baniyash, M. & Mandelboim, O. (2003). Complexes of HLA-G protein on the cell surface are important for leukocyte Ig-like receptor-1 function. *Journal of Immunology* **171**, 1343-1351.
- Gorer, P. A. (1937). The genetic and antigenic basis of tumour transplantation. *Journal of Pathology and Bacteriology* **44**, 691-697.
- Gorer, P. A., Lyman, S. & Snell, G. D. (1948). Studies on the genetic and antigenic basis of tumour transplantation. Linkage between a histocompatibility gene and "fused" in mice. *Proceedings of the Royal Society of London - Series B: Biological Sciences* **135**, 499-505.

- Groh, V., Bahram, S., Bauer, S., Herman, A., Beauchamp, M. & Spies, T. (1996). Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 12445-12450.
- Groh, V., Rhinehart, R., Randolph-Habecker, J., Topp, M. S., Riddell, S. R. & Spies, T. (2001). Costimulation of CD8 α beta T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nature Immunology* **2**, 255-260.
- Gromme, M. & Neefjes, J. (2002). Antigen degradation or presentation by MHC class I molecules via classical and non-classical pathways. *Molecular Immunology* **39**, 181-202.
- Gumperz, J. E., Litwin, V., Phillips, J. H., Lanier, L. L. & Parham, P. (1995). The Bw4 public epitope of HLA-B molecules confers reactivity with natural killer cell clones that express NKB1, a putative HLA receptor. *Journal of Experimental Medicine* **181**, 1133-1144.
- Gustafson, K. S. & Ginder, G. D. (1996). Interferon-gamma induction of the human leukocyte antigen-E gene is mediated through binding of a complex containing STAT1 α to a distinct interferon-gamma-response element. *Journal of Biological Chemistry* **271**, 20035-20046.
- Harada, H., Fujita, T., Miyamoto, M., Kimura, Y., Maruyama, M., Furia, A., Miyata, T. & Taniguchi, T. (1989). Structurally similar but functionally distinct factors, IRF-1 and IRF-2, bind to the same regulatory elements of IFN and IFN-inducible genes. *Cell* **58**, 729-739.
- Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F., Kirschning, C., Akira, S., Lipford, G., Wagner, H. & Bauer, S. (2004). Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* **303** 1526-1529.
- Heinmeyer, T., Wingender, E., Reuter, I., Hermjakob, H., Kel, A. E., Kel, O. V., Ignatieva, E. V., Ananko, E. A., Podkolodnaya, O. A., Kolpakov, F. A., Podkolodny, N. L. & Kolchanov, N. A. (1998). Databases on Transcriptional Regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids Research* **26**, 364-370.
- Herberman, R. B., Nunn, M. E. & Lavrin, D. H. (1975). Natural cytotoxic reactivity of mouse lymphoid cells against syngeneic acid allogeneic tumors. I. Distribution of reactivity and specificity. *International Journal of Cancer* **16**, 216-229.

- Holmes, M. A., Li, P., Petersdorf, E. W. & Strong, R. K. (2002). Structural studies of allelic diversity of the MHC class I homolog MIC-B, a stress-inducible ligand for the activating immunoreceptor NKG2D. *Journal of Immunology* **169**, 1395-1400.
- Horton, R., Wilming, L., Rand, V., Lovering, R. C., Bruford, E. A., Khodiyar, V. K., Lush, M. J., Povey, S., Talbot, C. C. Jr., Wright, M. W., Wain, H. M., Trowsdale, J., Zeigler, A. & Beck, S. (2004). Gene map of the extended human MHC. *Nature Reviews Genetics* **5**: 889-899.
- Hou, S., Guan, H. & Ricciardi, R. P. (2002). In adenovirus type 12 tumorigenic cells, major histocompatibility complex class I transcription shutoff is overcome by induction of NF-kappaB and relief of COUP-TFII repression. *Journal of Virology* **76**, 3212-3220.
- Huppa, J. B. & Ploegh, H. L. (1997). The alpha chain of the T cell antigen receptor is degraded in the cytosol. *Immunity* **7**, 113-122.
- Ibrahim, E. C., Morange, M., Dausset, J., Carosella, E. D. & Paul, P. (2000). Heat shock and arsenate induce expression of the non-classical class I histocompatibility HLA-G gene in tumor cell lines. *Cell Stress Haperones* **5**, 207-218.
- Isaacs, A. & Lindenmann, J. (1957). Virus interference. I. The interferon. *Proceedings of the Royal Society of London - Series B: Biological Sciences* **147**, 258-267.
- Israel, A., Le Bail, O., Hatat, D., Piette, J., Kieran, M., Logeat, F., Wallach, D., Fellous, M. & Kourilsky, P. (1989). TNF stimulates expression of mouse MHC class I genes by inducing an NF kappa B-like enhancer binding activity which displaces constitutive factors. *EMBO Journal* **8**, 3793-800.
- Ivanoska, D., Sun, D. C. & Lunney, J. K. (1991). Production of monoclonal antibodies reactive with polymorphic and monomorphic determinants of SLA class I gene products. *Immunogenetics* **33**, 220-223.
- Jesse, T. L., LaChance, R., Iademarco, M. F. & Dean, D. C. (1998). Interferon regulatory factor-2 is a transcriptional activator in muscle where it regulates expression of vascular cell adhesion molecule-1. *Journal of Cell Biology* **140**, 1265-1276.
- Johnson, D. R. (2003). Locus-specific constitutive and cytokine-induced HLA class I gene expression. *Journal of Immunology* **170**, 1894-1902.
- Jugovic, P., Hill, A. M., Tomazin, R., Ploegh, H. & Johnson, D. C. (1998). Inhibition of major histocompatibility complex class I antigen presentation in pig and primate cells by herpes simplex virus type 1 and 2 ICP47. *Journal of Virology* **72**, 5076-5084.

- Kaufman, R. J. & Murtha, P. (1987). Translational control mediated by eucaryotic initiation factor-2 is restricted to specific mRNAs in transfected cells. *Molecular and Cellular Biology* **7**, 1568-1571.
- Kessler, B., Hong, X., Petrovic, J., Borodovsky, A., Dantuma, N. P., Bogyo, M., Overkleeft, H. S., Ploegh, H. & Glas, R. (2003). Pathways accessory to proteasomal proteolysis are less efficient in major histocompatibility complex class I antigen production. *Journal of Biological Chemistry* **278**, 10012-10021.
- King, P. & Goodbourn, S. (1994). The beta-interferon promoter responds to priming through multiple independent regulatory elements. *Journal of Biological Chemistry* **269**, 30609-30615.
- King, P. & Goodbourn, S. (1998). STAT1 is inactivated by a caspase. *Journal of Biological Chemistry* **273**, 8699-8704.
- Kisselev, A. F., Akopian, T. N., Woo, K. M. & Goldberg, A. L. (1999). The sizes of peptides generated from protein by mammalian 26 and 20 S proteasomes. Implications for understanding the degradative mechanism and antigen presentation. *Journal of Biological Chemistry* **274**, 3363-3371.
- Kotenko, S. V., Gallagher, G., Baurin, V. V., Lewis-Antes, A., Shen, M., Shah, N. K., Langer, J. A., Sheikh, F., Dickensheets, H. & Donnelly, R. P. (2003). IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. *Nature Immunology* **4**, 69-77.
- Kraus, T. A., Lau, J. F., Parisien, J. P. & Horvath, C. M. (2003). A hybrid IRF9-STAT2 protein recapitulates interferon-stimulated gene expression and antiviral response. *Journal of Biological Chemistry* **278**, 13033-13038.
- Lanier, L. L. (1998). NK cell receptors. *Annual Review of Immunology* **16**, 359-393.
- Lanier, L. L. (2003). Natural killer cell receptor signaling. *Current Opinion in Immunology* **15**, 308-314.
- Larocca, S. A., Herbert, R. J., Crooke, H., Drew, T. W., Wileman, T. E. & Powell, P. P. (2005). Loss of interferon regulatory factor 3 in cells infected with classical swine fever virus involves the N-terminal protease, Npro. *Journal of Virology* **79**, 7239-7247.
- Laude, H., Gelfi, J., Lavanant, L. & Charley, B. (1992). Single amino acid changes in the viral glycoprotein M affect induction of alpha interferon by the coronavirus transmissible gastroenteritis virus. *Journal of Virology* **66**, 743-749.

- Lefebvre, S., Berrih-Aknin, S., Adrian, F., Moreau, P., Poea, S., Gourand, L., Dausset, J., Carosella, E. D. & Paul, P. (2001). A specific interferon (IFN)-stimulated response element of the distal HLA-G promoter binds IFN-regulatory factor 1 and mediates enhancement of this nonclassical class I gene by IFN-beta. *Journal of Biological Chemistry* **276**, 6133-6139.
- Lehner, P. J. & Cresswell, P. (2004). Recent developments in MHC-class-I-mediated antigen presentation. *Current Opinion in Immunology* **16**, 82-89.
- Lilley, B. N. & Ploegh, H. L. (2005). Viral modulation of antigen presentation: manipulation of cellular targets in the ER and beyond. *Immunological Reviews* **207**, 126-144.
- Lin, R., Heylbroeck, C., Genin, P., Pitha, P. M. & Hiscott, J. (1999). Essential role of interferon regulatory factor 3 in direct activation of RANTES chemokine transcription. *Molecular and Cellular Biochemistry* **19**, 959-966.
- Ljunggren, H. G. & Karre, K. (2002). In search of the 'missing self': MHC molecules and NK cell recognition. *Immunology Today* **11**, 237-44.
- Lunney, J. K. (1994). Current status of the swine leukocyte antigen complex. *Veterinary Immunology and Immunopathology* **43**, 19-28.
- Lund, J., Sato, A., Akira, S., Medzhitov, R. & Iwasaki, A. (2003). Toll-like receptor 9-mediated recognition of Herpes simplex virus-2 by plasmacytoid dendritic cells. *Journal of Experimental Medicine* **198**, 513-520.
- Lunny, J. K. & Pescovitz, M. D. (1988). Differentiation antigens of swine lymphoid tissues. In Differentiation antigens in lymphohemopoietic tissues, pp 421-454. Edited by M. M. T. Z. New York and Basel: Marcel Dekker, INC.
- Maguire, J. E., Ehrlich, R., Frels, W. I. & Singer, D. S. (1990). Regulation of expression of a class I major histocompatibility complex transgene. *Journal of Reproduction and Fertility Supplements* **41**, 59-62.
- Malmgaard, L. (2004). Induction and regulation of IFNs during viral infections. *Journal of Interferon and Cytokine Research* **24**, 439-454.
- Marais, R., Light, Y., Paterson, H. F. & Marshall, C. J. (1995). Ras recruits Raf-1 to the plasma membrane for activation by tyrosine phosphorylation. *EMBO Journal* **14**, 3136-3145.

- Masumi, A., Yamakawa, Y., Fukazawa, H., Ozato, K. & Komuro, K. (2003). Interferon regulatory factor-2 regulates cell growth through its acetylation. *Journal of Biological Chemistry* **278**, 25401-25407.
- Marie, I., Durbin, J. E. & Levy, D. E. (1998). Differential viral induction of distinct interferon-alpha genes by positive feedback through interferon regulatory factor-7. *EMBO Journal* **17**, 6660-6669.
- Martin, B. K., Chin, K. C., Olsen, J. C., Skinner, C. A., Dey, A., Ozato, K. & Ting, J. P. (1997). Induction of MHC class I expression by the MHC class II transactivator CIITA. *Immunity* **6**, 591-600.
- Massa, P. T. & Wu, H. (1995). Interferon regulatory factor element and interferon regulatory factor 1 in the induction of major histocompatibility complex class I genes in neural cells. *Journal of Interferon and Cytokine Research* **15**, 799-810.
- Meuwissen, M. P., Horst, S. H., Huirne, R. B. & Dijkhuizen, A. A. (1999). A model to estimate the financial consequences of classical swine fever outbreaks: principles and outcomes. *Preventive Veterinary Medicine* **42**, 249-270.
- Mizushima, S. & Nagata, S. (1990). pEF-BOS, a powerful mammalian expression vector. *Nucleic Acids Research* **18**, 5322.
- Molinero, L. L., Fuertes, M. B., Girart, M. V., Fainboim, L., Rabinovich, G. A., Costas, M. A. & Zwirner, N. W. (2004). NF-kappa B regulates expression of the MHC class I-related chain A gene in activated T lymphocytes. *Journal of Immunology* **173**, 5583-5590.
- Milone, M. C. & Fitzgerald-Bocarsly, P. (1998). The mannose receptor mediates induction of IFN-alpha in peripheral blood dendritic cells by enveloped RNA and DNA viruses. *Journal of Immunology* **161**, 2391-2399.
- Moretta, A., Bottino, C., Vitale, M., Pende, D., Biassoni, R., Mingari, M. C. & Moretta, L. (1996). Receptors for HLA class-I molecules in human natural killer cells. *Annual Review of Immunology* **14**, 619-648.
- Moynagh, P. N. (2005). TLR signalling and activation of IRFs: revisiting old friends from the NFkB pathway. *Trends in Immunology* **26**, 469-476.
- Navarro, F., Llano, M., Bellon, T., Colonna, M., Geraghty, D. E. & Lopez-Botet, M. (1999). The ILT2(LIR1) and CD94/NKG2A NK cell receptors respectively recognize HLA-G1 and HLA-E molecules co-expressed on target cells. *European Journal of Immunology* **29**, 277-283.

- Oleksiewicz, M. B., Rasmussen, T. B., Normann, P. & Uttenhal, A. (2003). Determination of the sequence of the complete open reading frame of the Paderborn isolate of classical swine fever virus. *Veterinary Microbiology* **92**, 311-325.
- Palmer, E. & Cresswell, P. (1998). Mechanisms of MHC class I-restricted antigen processing. *Annual Review of Immunology* **16**, 323-358.
- Parham, P. (2005). MHC class I molecules and KIRs in human history, health and survival. *Nature Reviews Immunology* **5**, 201-214.
- Pauly, T., Elbers, K., Konig, M., Lengsfeld, T., Saalmuller, A. & Thiel, H. J. (1995). Classical swine fever virus-specific cytotoxic T lymphocytes and identification of a T cell epitope. *Journal of General Virology* **76**, 3039-3049.
- Pennington, L. R., Lunney, J. K. & Sachs, D. H. (1981). Transplantation in miniature swine. VIII. Recombination within the major histocompatibility complex of miniature swine. *Transplantation* **31**, 66-71.
- Pescovitz, M. D., Lunney, J. K. & Sachs, D. H. (1984). Preparation and characterization of monoclonal antibodies reactive with porcine PBL. *Journal of Immunology* **133**, 368-375.
- Pestka, S., Krause, C. D. & Walter, M. R. (2004). Interferons, interferon-like cytokines, and their receptors. *Immunological Reviews* **202**, 8-32.
- Piguet, V., Wan, L., Borel, C., Mangasarian, A., Demareux, N., Thomas, G. & Trono, D. (2000). HIV-1 Nef protein binds to the cellular protein PACS-1 to downregulate class I major histocompatibility complexes. *Nature Cell Biology* **2**, 163-167.
- Pizzato, N., Derrien, M. & Lenfant, F. (2004). The short cytoplasmic tail of HLA-G determines its resistance to HIV-1 Nef-mediated cell surface downregulation. *Human Immunology* **65**, 1389-1396.
- Posch, P. E., Borrego, F., Brooks, A. G. & Coligan, J. E. (1998). HLA-E is the ligand for the natural killer cell CD94/NKG2 receptors. *Journal of Biomedical Science* **5**, 321-331.
- Rabin, M., Fries, R., Singer, D. & Ruddle, F. H. (1985). Assignment of the porcine major histocompatibility complex to chromosome 7 by in situ hybridization. *Cytogenetics and Cell Genetics* **39**, 206-209.
- Radaev, S., Kattah, M., Zou, Z., Colonna, M. & Sun, P. D. (2002). Making sense of the diverse ligand recognition by NKG2D. *Journal of Immunology* **169**, 6279-6285.

- Rady, P. L., Cadet, P., Bui, T. K., Tying, S. K., Baron, S., Stanton, G. J. & Hughes, T. K. (1995). Production of interferon gamma messenger RNA by cells of non-immune origin. *Cytokine* **7**, 793-798.
- Rajagopalan, S. & Long, E. O. (1999). A human histocompatibility leukocyte antigen (HLA)-G-specific receptor expressed on all natural killer cells. *Journal of Experimental Medicine* **189**, 1093-1100.
- Reid, L. E., Brasnett, A. H., Gilbert, C. S., Porter, A. C., Gewert, D. R., Stark, G. R. & Kerr, I. M. (1989). A single DNA response element can confer inducibility by both alpha- and gamma-interferons. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 840-844.
- Reis, L. F., Harada, H., Wolchok, J. D., Taniguchi, T. & Vilcek, J. (1992). Critical role of a common transcription factor, IRF-1, in the regulation of IFN-beta and IFN-inducible genes. *EMBO Journal* **11**, 185-193.
- Reith, W. & Mach, B. (2001). The bare lymphocyte syndrome and the regulation of MHC expression. *Annual Review of Immunology* **19**, 331-373.
- Reits, E. A., Vos, J. C., Gromme, M. & Neefjes, J. (2000). The major substrates for TAP in vivo are derived from newly synthesized proteins. *Nature* **404**, 774-778.
- Reits, E., Neijssen, J., Herberts, C., Benckhuijsen, W., Janssen, L., Drijfhout, J. W. & Neefjes, J. (2004). A major role for TPPII in trimming proteasomal degradation products for MHC class I antigen presentation. *Immunity* **20**, 495-506.
- Renard, C., Vaiman, M., Chiannilkulchai, N., Cattolico, L., Robert, C. & Chardon, P. (2001). Sequence of the pig major histocompatibility region containing the classical class I genes. *Immunogenetics* **53**, 490-500.
- Rogel-Gaillard, C., Bourgeaux, N., Save, J. C., Renard, C., Coullion, P., Pinton, P., Yerle, M., Vaiman, M. & Chardon, P. (1997). Construction of a swine YAC library allowing an efficient recovery of unique and centromeric repeated sequences. *Mammalian Genome* **3**, 186-192.
- Rogel-Gaillard, C., Bourgeaux, N., Billault, A., Vaiman, M. & Chardon, P. (1999). Construction of a swine BAC library: application to the characterization and mapping of porcine type C endoviral elements. *Cytogenetics and Cell Genetics* **85**, 205-211.

- Rousseau, P., Masternak, K., Krawczyk, M., Reith, W., Dausset, J., Carosella, E. D. & Moreau, P. (2004). In vivo, RFX5 binds differently to the human leukocyte antigen-E, -F and -G promoters and participates in HLA class I protein expression in a cell type-dependent manner. *Immunology* **1**, 53-65.
- Rousseau, P., Paul, P., O'Brien, M., Dausset, J., Carosella, E. D. & Moreau, P. (2000) The X1 box of HLA-G promoter is a target site for RFX and Sp1 factors. *Human Immunology* **61**, 1132-1137.
- Rozen S & Skaletsky, H. (2000). Primer3 on the www for general users and for biologist programmers. In: Krawetz, S, Misener S (eds) *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa, NJ, pp 365-386.
- Sadasivan, B., Lehner, P. J., Ortmann, B., Spies, T. & Cresswell, P. (1996). Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* **5**, 103-114.
- Salter, R. D., Norment, A. M., Chen, B. P., Clayberger, C., Krensky, A. M., Littman, D. R. & Parham, P. (1989). Polymorphism in the alpha 3 domain of HLA-A molecules affects binding to CD8. *Nature* **338**, 345-347.
- Saric, T., Chang, S. C., Hattori, A., York, I. A., Markant, S., Rock, K. L., Tsujimoto, M. & Goldberg, A. L. (2002). An IFN-gamma-induced aminopeptidase in the ER, ERAP1, trims precursors to MHC class I-presented peptides. *Nature Immunology* **3**, 1169-1176.
- Satz, M. L. & Singer, D. S. (1984). Effect of mouse interferon on the expression of a porcine major histocompatibility gene introduced into mouse L cells. *Journal of Immunology* **132**, 496-501.
- Satz, M. L., Wang, L. C., Singer, D. S. & Rudikoff, S. (1985). Structure and expression of two porcine genomic clones encoding class I MHC antigens. *Journal of Immunology* **135**, 2167-2175.
- Schubert, U., Anton, L. C., Gibbs, J., Norbury, C. C., Yewdell, J. W. & Bennink, J. R. (2000). Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* **404**, 770-774.
- Serwold, T., Gonzalez, F., Kim, J., Jacob, R. & Shastri, N. (2002). ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* **419**, 480-483.

- Sheppard, P., Kindsvogel, W., Xu, W., Henderson, K., Schlutsmeyer, S., Whitmore, T. E., Kuestner, R., Garrigues, U., Birks, C., Roraback, J., Ostrander, C., Dong, D., Shin, J., Presnell, S., Fox, B., Haldeman, B., Cooper, E., Taft, D., Gilbert, T., Grant, F. J., Tackett, M., Krivan, W., McKnight, G., Clegg, C., Foster, D. & Klucher, K. M. (2003). IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nature Immunology* **4**, 63-68.
- Sica, A., Dorman, L., Viggiano, V., Cippitelli, M., Ghosh, P., Rice, N. & Young, H. A. (1997). Interaction of NF-kappaB and NFAT with the interferon-gamma promoter. *Journal of Biological Chemistry* **272**, 30412-30420.
- Singer, D. S., Ehrlich, R., Satz, L., Frels, W., Bluestone, J., Hodes, R. & Rudikoff, S. (1987). Structure and expression of class I MHC genes in the miniature swine. *Veterinary Immunology and Immunopathology* **17**, 211-221.
- Smith, D. M., Lunney, J. K., Martens, G. W., Ando, A., Lee, J. H., Ho, C. S., Schook, L., Renard, C. & Chardon, P. (2005). Nomenclature for factors of the SLA class-I system, 2004. *Tissue Antigens* **65**, 136-49.
- Stryer, L. (2001). Programmed protein destruction. In *Biochemistry* 5th Edition pp 942-943. Edited by Tynoczka, J.L. & Berg, J.M. New York and Basingstoke: WH Freemans and Company.
- Suzuka, I., Shimizu, N., Sekiguchi, K., Hoshino, H., Kodama, M. & Shimotohno, K. (1986). Molecular cloning of unintegrated closed circular DNA of porcine retrovirus. *FEBS Letters* **198**, 339-343.
- Szabo, S. J., Kim, S. T., Costa, G. L., Zhang, X., Fathman, C. G. & Glimcher, L. H. (2000). A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* **100**, 655-669.
- Tabeta, K., Georgel, P., Janssen, E., Du, X., Hoebe, K., Crozat, K., Mudd, S., Shamel, L., Sovath, S., Goode, J., Alexopoulou, L. & Flavell, R. A. (2004). Toll-like receptors 9 and 3 are essential components of innate immune defense against mouse cytomegalovirus infection. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 3516-3521.
- Takeda, K. & Akira, S. (2005). Toll-like receptors in innate immunity. *International Immunology* **17**, 1-14.
- Terenzi, F., deVeer, M. J., Ying, H., Restifo, N. P., Williams, B. R. & Silverman, R. H. (1999). The antiviral enzymes PKR and RNase L suppress gene expression from viral and non-viral based vectors. *Nucleic Acids Research* **27**, 4369-4375.

- Trowsdale, J. (1995). "Both man & bird & beast": comparative organization of MHC genes. *Immunogenetics* **41**, 1-17.
- Vaiman, M., Renard, C., LaFage, P., Ameteau, J. & Nizza, P. (1970). Evidence for a histocompatibility system in swine (SL-A). *Transplantation* **10**, 155-164.
- Vallee, I., Tait, S. W. & Powell, P. P. (2001). African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. *Journal of Virology* **75**, 10372-10382.
- van den Elsen, P. J., Holling, T. M., Kuipers, H. F. & van der Stoep, N. (2004). Transcriptional regulation of antigen presentation. *Current Opinion in Immunology* **16**, 67-75.
- van den Elsen, P. J., Peijnenburg, A., van Eggermond, M. C. & Gobin, S. J. (1998). Shared regulatory elements in the promoters of MHC class I and class II genes. *Immunology Today* **19**, 308-312.
- Vance, R. E., Kraft, J. R., Altman, J. D., Jensen, P. E. & Raulet, D. H. (1998). Mouse CD94/NKG2A is a natural killer cell receptor for the nonclassical major histocompatibility complex (MHC) class I molecule Qa-1(b). *Journal of Experimental Medicine* **188**, 1841-1848.
- Van Gennip, H. G., Vlot, A. C., Hulst, M. M., De Smit, A. J. & Moormann, R. J. (2004). Determinants of virulence of classical swine fever virus strain Brescia. *Journal of Virology* **78**, 8812-8823.
- Velten, F., Rogel-Gaillard, C., Renard, C., Pontarotti, P., Tazi-Ahnini, R., Vaiman, M. & Chardon, P. (1998). A first map of the porcine major histocompatibility complex class I region. *Tissue Antigens* **51**, 183-194.
- Velten, F. W., Renard, C., Rogel-Gaillard, C., Vaiman, M. & Chardon, P. (1999). Spatial arrangement of pig MHC class I sequences. *Immunogenetics* **49**, 919-930.
- Viza, D., Sugar, J. R. & Binns, R. M. (1970). Lymphocyte stimulation in pigs: evidence for the existence of a single major histocompatibility locus, PL-A. *Nature* **227**, 949-950.
- Wainwright, S. D., Biro, P. A. & Holmes, C. H. (2000). HLA-F is a predominantly empty, intracellular, TAP-associated MHC class Ib protein with a restricted expression pattern. *The Journal of Immunology* **164**, 319-328.

- Weissman, J. D. & Singer, D. S. (1991). A complex regulatory DNA element associated with a major histocompatibility complex class I gene consists of both a silencer and an enhancer. *Molecular Cell Biology* **11**, 4217-4227.
- Wheelock, E. F. (1965). Interferon-like virus inhibitor induced in human leukocytes by phytohaemagglutinin. *Science* **149**, 310.
- White, L. C., Wright, K. L., Felix, N. J., Ruffner, H., Reis, L. F., Pine, R. & Ting, J. P. (1996). Regulation of LMP2 and TAP1 genes by IRF-1 explains the paucity of CD8⁺ T cells in IRF-1^{-/-} mice. *Immunity* **5**, 365-376.
- Whitelegg, A. M. E (2001). Analysis of peptides bound to porcine MHC class I molecules: PhD thesis. 2pp. University of Bristol.
- Xi, H. & Blanck, G. (2003). The IRF-2 DNA binding domain facilitates the activation of the class II transactivator (CIITA) type IV promoter by IRF-1. *Molecular Immunology* **39**, 677-684.
- Ye, J., Cipitelli, M., Dorman, L., Ortaldo, J. R. & Young, H. A. (1996). The nuclear factor YY1 suppresses the human gamma interferon promoter through two mechanisms: inhibition of AP1 binding and activation of a silencer element. *Molecular and Cellular Biology* **16**, 4744-4753.
- Zhang, L. & Pagano, J. S. Structure and function of IRF-7. (2002). *Journal of Interferon Cytokine Research* **22**, 95-101.
- Zwirner, N. W., Fernandez-Vina, M. A. & Stastny, P. (1998). MICA, a new polymorphic HLA-related antigen, is expressed mainly by keratinocytes, endothelial cells, and monocytes. *Immunogenetics* **47**, 139-148.